



wwPDB EM Validation Summary Report ⓘ

Sep 9, 2026 – 02:19 PM JST

PDB ID : 21RN / pdb_000021rn
EMDB ID : EMD-67945
Title : Cryo-EM structure of mouse myeloperoxidase in complex with Fab fragments of antibodies mAb-A24 and mAb-A46
Authors : Fujii, T.; Irie, M.; Torizawa, T.
Deposited on : 2025-12-23
Resolution : 2.95 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev133
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 2.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	13114 (2.45 - 3.45)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	224	5% 42% 9% 48%
2	B	213	11% 46% • 50%
3	C	224	12% 41% 9% 49%
4	D	213	11% 42% 8% 50%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	E	114	17% 68% 20% • 11%
6	F	476	6% 78% 14% • 8%

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 7871 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called mAb-A46 Fab Heavy chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	117	Total	C	N	O	S	0	0
			924	588	148	185	3		

- Molecule 2 is a protein called mAb-A46 Fab Light chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	106	Total	C	N	O	S	0	0
			817	522	135	158	2		

- Molecule 3 is a protein called mAb-A24 Fab Heavy chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	C	115	Total	C	N	O	S	0	0
			888	557	152	176	3		

- Molecule 4 is a protein called mAb-A24 Fab Light chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	D	106	Total	C	N	O	S	0	0
			819	521	135	160	3		

- Molecule 5 is a protein called Myeloperoxidase light chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	E	102	Total	C	N	O	S	0	0
			824	520	152	148	4		


- Molecule 6 is a protein called Myeloperoxidase.


Mol	Chain	Residues	Atoms					AltConf	Trace
6	F	440	Total	C	N	O	S	0	0
			3527	2240	646	619	22		

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	719	HIS	-	expression tag	UNP Q6RFG4
F	720	HIS	-	expression tag	UNP Q6RFG4
F	721	HIS	-	expression tag	UNP Q6RFG4
F	722	HIS	-	expression tag	UNP Q6RFG4
F	723	HIS	-	expression tag	UNP Q6RFG4
F	724	HIS	-	expression tag	UNP Q6RFG4
F	725	HIS	-	expression tag	UNP Q6RFG4
F	726	HIS	-	expression tag	UNP Q6RFG4
F	727	HIS	-	expression tag	UNP Q6RFG4
F	728	HIS	-	expression tag	UNP Q6RFG4

- Molecule 7 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).





Mol	Chain	Residues	Atoms				AltConf
8	F	1	Total	C	N	O	0
			14	8	1	5	
8	F	1	Total	C	N	O	0
			14	8	1	5	

- Molecule 9 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		AltConf
9	F	1	Total	Ca	0
			1	1	

PHE PRO
ALA VAL
LEU LEU
GLN SER
ASP ASP
LEU LEU
TYR THR
THR THR
SER SER
SER SER
SER SER
VAL VAL
THR THR
THR THR
SER SER
SER SER
PRO PRO
TRP TRP
SER SER
GLN GLN
SER SER
ILE ILE
THR THR
CYS CYS
ASN ASN
VAL VAL
ALA ALA
HIS HIS
PRO PRO
ALA ALA
SER SER
SER SER
THR THR
THR THR
VAL VAL
LYS LYS
ASP ASP
LYS LYS
ILE ILE
GLU GLU
PRO PRO
ARG ARG
GLY GLY
PRO PRO
THR THR
ILE ILE
LYS LYS
PRO PRO

• Molecule 4: mAb-A24 Fab Light chain

Chain D: 11% 42% 8% 50%

D1 M4 T5 O6 S7 S12 A13 S14 L15 G16 G17 K18 V19 T20 I21 K31 Y32 I33 A34 W35 Y36 Q37 G41 L47 P56 G57 I58 P59 S60 R61 F73 S74 I75 S76 N77 L78 E79 P80 E81 D82 C88 E104 L105 K106 ARG ALA ASP ALA PRO THR

VAL SER ILE PHE PRO THR PRO SER SER LEU GLN THR THR GLY GLY ALA VAL VAL CYS PHE LEU ASN ASN PHE TYR PRO LYS ASP ILE ASN VAL LYS TRP PHE ILE ASP ARG ASN GLY SER GLU ARG ASN GLN GLY VAL LEU ASN SER TRP THR ASP GLN ASP SER ASP LYS SER THR TYR SER

MET SER SER LEU THR LEU THR LYS ASP TYR ARG HIS SER TYR THR CYS ALA THR HIS LYS THR SER THR ASP PRO ILE VAL LYS SER PHE ASN ARG ASN GLY CYS

• Molecule 5: Myeloperoxidase light chain

Chain E: 17% 68% 20% 11%

VAL THR CYS P142 P143 M144 R158 G163 R171 A175 W187 T188 P189 G190 V191 N192 V204 V209 R210 F211 P212 N213 D214 Q215 L216 T217 K218 D219 Q220 E221 A223 L224 M225 F226 M227 G230 Q231 F232 L233 D234 H235 D236 I237 T238 L239 P243 ALA THR ARG PHE

SER PHE THR GLY

• Molecule 6: Myeloperoxidase

Chain F: 6% 78% 14% 8%

LEU ASN CYS THR SER CYS LEU GLN GLN PRO P264 I270 P271 P272 N273 D274 P275 R276 I277 K278 N279 Q280 K281 D282 C283 F287 A292 CYS THR ARG ASN ILE ILE R301 L307 T308 S309 D312 D321 P322 L323 L327 M351 D354 SER LEU HIS ASP

ASP P360 T364 N365 R366 R369 M383 P384 E385 L386 H390 T391 T392 R410 W411 L416 E419 I423 M427 Y436 V440 P443 K447 Q452 I464 M470 R473 Y474 G475 H476 Q480 V499 P500 K503 V504 F505 D518 P519

M525 P528 A529 K530 V539 D540 E541 I542 R543 E544 R545 F547 E548 Q549 V550 M551 R552 I553 G554 L555 N561 M562 V588 E599 R602 M605 P611 I614 M618 R630 W653 W654 E655 K662 Q663 P674 I677 T685 V686 S687 K688 ASN M690

Y697 P698 R699 D700 F701 N705 E717 THR HIS HIS HIS HIS HIS HIS HIS HIS HIS HIS

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	1062641	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS GLACIOS	Depositor
Voltage (kV)	200	Depositor
Electron dose ($e^-/\text{\AA}^2$)	60.0	Depositor
Minimum defocus (nm)	1500	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	FEI FALCON IV (4k x 4k)	Depositor
Maximum map value	3.214	Depositor
Minimum map value	-2.935	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.036	Depositor
Recommended contour level	0.49	Depositor
Map size (Å)	354.80002, 354.80002, 354.80002	wwPDB
Map dimensions	400, 400, 400	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.887, 0.887, 0.887	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, NAG, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.13	0/949	0.32	0/1293
2	B	0.14	0/838	0.35	0/1136
3	C	0.12	0/906	0.25	0/1224
4	D	0.13	0/839	0.30	0/1136
5	E	0.14	0/849	0.32	0/1155
6	F	0.14	0/3609	0.30	0/4888
All	All	0.14	0/7990	0.31	0/10832

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	924	0	871	12	0
2	B	817	0	805	3	0
3	C	888	0	858	12	0
4	D	819	0	805	6	0
5	E	824	0	794	22	0
6	F	3527	0	3559	43	0
7	F	43	0	30	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	F	28	0	26	0	0
9	F	1	0	0	0	0
All	All	7871	0	7748	85	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 85 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:30:THR:HG23	3:C:54:ARG:HB3	1.67	0.77
3:C:91:SER:HA	3:C:114:VAL:O	1.83	0.77
5:E:224:LEU:HD21	6:F:480:GLN:HG3	1.78	0.65
2:B:37:GLN:HB2	2:B:47:LEU:HD11	1.79	0.65
5:E:237:ILE:HG13	5:E:238:THR:HG22	1.78	0.65

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	115/224 (51%)	109 (95%)	6 (5%)	0	100	100
2	B	104/213 (49%)	95 (91%)	9 (9%)	0	100	100
3	C	111/224 (50%)	108 (97%)	3 (3%)	0	100	100
4	D	104/213 (49%)	100 (96%)	4 (4%)	0	100	100
5	E	100/114 (88%)	97 (97%)	3 (3%)	0	100	100
6	F	432/476 (91%)	422 (98%)	10 (2%)	0	100	100
All	All	966/1464 (66%)	931 (96%)	35 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	101/195 (52%)	97 (96%)	4 (4%)	28	53
2	B	90/188 (48%)	88 (98%)	2 (2%)	45	69
3	C	95/190 (50%)	90 (95%)	5 (5%)	20	45
4	D	91/189 (48%)	86 (94%)	5 (6%)	19	43
5	E	89/99 (90%)	87 (98%)	2 (2%)	45	69
6	F	387/423 (92%)	378 (98%)	9 (2%)	44	67
All	All	853/1284 (66%)	826 (97%)	27 (3%)	35	59

5 of 27 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	D	19	VAL
5	E	220	GLN
6	F	542	ILE
5	E	216	LEU
6	F	308	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
6	F	534	GLN
6	F	680	ASN
4	D	49	HIS
5	E	156	ASN
5	E	213	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
8	NAG	F	802	6	14,14,15	0.37	0	17,19,21	0.61	0
8	NAG	F	803	6	14,14,15	0.74	0	17,19,21	0.86	0
7	HEM	F	801	6,5	50,50,50	0.72	1 (2%)	66,82,82	0.67	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	NAG	F	802	6	-	4/6/23/26	0/1/1/1
8	NAG	F	803	6	-	0/6/23/26	0/1/1/1
7	HEM	F	801	6,5	-	7/14/54/54	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	F	801	HEM	CBB-CAB	2.23	1.41	1.30

There are no bond angle outliers.

There are no chirality outliers.

5 of 11 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
7	F	801	HEM	C2C-C3C-CAC-CBC
7	F	801	HEM	C4C-C3C-CAC-CBC
8	F	802	NAG	C8-C7-N2-C2
8	F	802	NAG	O7-C7-N2-C2
8	F	802	NAG	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	F	801	HEM	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-67945. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 200




Y Index: 200




Z Index: 200


6.2.2 Raw map



X Index: 200



Y Index: 200




Z Index: 200


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 201




Y Index: 229




Z Index: 177


6.3.2 Raw map



X Index: 201



Y Index: 229




Z Index: 177


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.49. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_67945_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 34 nm^3 ; this corresponds to an approximate mass of 31 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.339 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.339 $\text{\AA}^{-1}$

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.95	-	-
Author-provided FSC curve	2.95	3.31	2.98
Unmasked-calculated*	3.35	3.55	3.36

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.35 differs from the reported value 2.95 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-67945 and PDB model 21RN. Per-residue inclusion information can be found in [section 3](#) on [page 7](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.49 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.49).

9.4 Atom inclusion [i](#)



At the recommended contour level, 74% of all backbone atoms, 74% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.49) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7400	0.5900
A	0.7320	0.5880
B	0.6870	0.5870
C	0.6720	0.5760
D	0.6580	0.5640
E	0.7080	0.5840
F	0.7970	0.6030

1.0

0.0

<0.0