wwPDB X-ray Structure Validation Summary Report (i)

Oct 22, 2026 – 02:14 PM JST

PDB ID	:	22OX / pdb_000022ox
Title	:	Crystal structure of ATRX ATPase domain
Authors	:	Wang, X.; Chen, Y.
Deposited on	:	2026-01-19
Resolution	:	2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.2
Xtriage (Phenix)	:	2.2
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	747	% 55% 34% · 10%

2 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 5444 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcriptional regulator ATRX.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	675	5444	3450	943	1018	33	0	0	0

There are 70 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	SER	deletion	UNP P46100
A	?	-	LEU	deletion	UNP P46100
A	?	-	SER	deletion	UNP P46100
A	?	-	SER	deletion	UNP P46100
A	?	-	ASP	deletion	UNP P46100
A	?	-	ASP	deletion	UNP P46100
A	?	-	TYR	deletion	UNP P46100
A	?	-	THR	deletion	UNP P46100
A	?	-	LYS	deletion	UNP P46100
A	?	-	LYS	deletion	UNP P46100
A	?	-	LYS	deletion	UNP P46100
A	?	-	LYS	deletion	UNP P46100
A	?	-	LYS	deletion	UNP P46100
A	?	-	GLY	deletion	UNP P46100
A	?	-	LYS	deletion	UNP P46100
A	?	-	LYS	deletion	UNP P46100
A	?	-	GLY	deletion	UNP P46100
A	?	-	LYS	deletion	UNP P46100
A	?	-	LYS	deletion	UNP P46100
A	?	-	ASP	deletion	UNP P46100
A	?	-	SER	deletion	UNP P46100
A	?	-	SER	deletion	UNP P46100
A	?	-	SER	deletion	UNP P46100
A	?	-	SER	deletion	UNP P46100
A	?	-	GLY	deletion	UNP P46100
A	?	-	SER	deletion	UNP P46100
A	?	-	GLY	deletion	UNP P46100

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	SER	deletion	UNP P46100
A	?	-	ASP	deletion	UNP P46100
A	?	-	ASN	deletion	UNP P46100
A	?	-	ASP	deletion	UNP P46100
A	?	-	VAL	deletion	UNP P46100
A	?	-	GLU	deletion	UNP P46100
A	?	-	VAL	deletion	UNP P46100
A	?	-	ILE	deletion	UNP P46100
A	?	-	LYS	deletion	UNP P46100
A	?	-	VAL	deletion	UNP P46100
A	?	-	TRP	deletion	UNP P46100
A	?	-	ASN	deletion	UNP P46100
A	?	-	SER	deletion	UNP P46100
A	?	-	ARG	deletion	UNP P46100
A	?	-	SER	deletion	UNP P46100
A	?	-	ARG	deletion	UNP P46100
A	?	-	GLY	deletion	UNP P46100
A	?	-	GLY	deletion	UNP P46100
A	?	-	GLY	deletion	UNP P46100
A	?	-	GLU	deletion	UNP P46100
A	?	-	GLY	deletion	UNP P46100
A	?	-	ASN	deletion	UNP P46100
A	?	-	VAL	deletion	UNP P46100
A	?	-	ASP	deletion	UNP P46100
A	?	-	GLU	deletion	UNP P46100
A	?	-	THR	deletion	UNP P46100
A	?	-	GLY	deletion	UNP P46100
A	?	-	ASN	deletion	UNP P46100
A	?	-	ASN	deletion	UNP P46100
A	?	-	PRO	deletion	UNP P46100
A	?	-	SER	deletion	UNP P46100
A	?	-	VAL	deletion	UNP P46100
A	?	-	SER	deletion	UNP P46100
A	?	-	LEU	deletion	UNP P46100
A	?	-	LYS	deletion	UNP P46100
A	?	-	LEU	deletion	UNP P46100
A	?	-	GLU	deletion	UNP P46100
A	?	-	GLU	deletion	UNP P46100
A	?	-	SER	deletion	UNP P46100
A	?	-	LYS	deletion	UNP P46100
A	?	-	ALA	deletion	UNP P46100
A	?	-	THR	deletion	UNP P46100

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	SER	deletion	UNP P46100

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Transcriptional regulator ATRX



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α, β, γ	159.53Å 60.40Å 94.14Å 90.00° 100.57° 90.00°	Depositor
Resolution (Å)	49.61 – 2.90 49.61 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.4 (49.61-2.90) 99.4 (49.61-2.90)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.51 (at 2.91Å)	Xtriage
Refinement program	PHENIX (1.18.2_3874: ???)	Depositor
R, R_{free}	0.232 , 0.278 0.239 , 0.279	Depositor DCC
R_{free} test set	939 reflections (4.74%)	wwPDB-VP
Wilson B-factor (Å^2)	89.8	Xtriage
Anisotropy	0.424	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.32 , 78.1	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	5444	wwPDB-VP
Average B, all atoms (Å^2)	115.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.75% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| > 5$	RMSZ	$\#\|Z\| > 5$
1	A	0.39	0/5543	0.69	0/7470

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5444	0	5401	210	0
All	All	5444	0	5401	210	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 210 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2141:PRO:HG3	1:A:2179:GLN:HG2	1.46	0.98
1:A:2079:ARG:HG2	1:A:2080:ASN:ND2	1.85	0.91
1:A:1591:ILE:HG13	1:A:1765:ILE:HD13	1.59	0.85
1:A:1656:LEU:HD11	1:A:1669:LEU:HD11	1.62	0.82
1:A:2020:MET:HE1	1:A:2050:ILE:HG12	1.64	0.79

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	663/747 (89%)	637 (96%)	25 (4%)	1 (0%)	43 72

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1711	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	589/673 (88%)	565 (96%)	24 (4%)	27 61

5 of 24 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2069	LEU
1	A	2095	ARG
1	A	2091	THR
1	A	2096	LYS
1	A	1867	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2156	GLN
1	A	2168	GLN
1	A	1877	GLN
1	A	2080	ASN
1	A	2093	GLN

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	675/747 (90%)	0.21	10 (1%)	72	64	56, 109, 191, 238	0

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1881	ARG	3.5
1	A	1523	ILE	3.2
1	A	2207	LEU	2.9
1	A	2213	ASP	2.6
1	A	1900	GLY	2.4

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no oligosaccharides in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.