wwPDB X-ray Structure Validation Summary Report (i)

Oct 19, 2026 – 04:08 PM EDT

PDB ID	:	12YZ / pdb_000012yz
Title	:	Crystal structure of human K293st geranylgeranyl diphosphate synthase with 5 geranylgeranyl diphosphate bound and 1 magnesium only subunits
Authors	:	Pham, A.C.; Borgstahl, G.E.O.
Deposited on	:	2026-04-23
Resolution	:	1.82 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.2
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.2
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1112 (1.82-1.82)
Clashscore	190562	1148 (1.82-1.82)
Ramachandran outliers	187476	1140 (1.82-1.82)
Sidechain outliers	187428	1140 (1.82-1.82)
RSRZ outliers	180081	1112 (1.82-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	294	4% 88% 8% ·
1	B	294	3% 90% 5% 5%
1	C	294	% 90% 6% ·
1	D	294	% 93% · ·
1	E	294	% 92% 5% ·

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	294	

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 16173 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Geranylgeranyl pyrophosphate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	283	Total	C	N	O	S	0	0	0
			2307	1485	388	428	6			
1	B	279	Total	C	N	O	S	0	0	0
			2278	1468	382	422	6			
1	C	284	Total	C	N	O	S	0	0	0
			2310	1486	388	430	6			
1	D	283	Total	C	N	O	S	0	0	0
			2310	1488	389	427	6			
1	E	286	Total	C	N	O	S	0	0	0
			2332	1499	391	435	7			
1	F	286	Total	C	N	O	S	0	0	0
			2336	1503	391	436	6			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP O95749
B	0	SER	-	expression tag	UNP O95749
C	0	SER	-	expression tag	UNP O95749
D	0	SER	-	expression tag	UNP O95749
E	0	SER	-	expression tag	UNP O95749
F	0	SER	-	expression tag	UNP O95749

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Mg	0	0
			2	2		
2	B	2	Total	Mg	0	0
			2	2		
2	C	2	Total	Mg	0	0
			2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	D	2	Total Mg 2 2	0	0
2	E	2	Total Mg 2 2	0	0
2	F	2	Total Mg 2 2	0	0

- Molecule 3 is GERANYLGERANYL DIPHOSPHATE (CCD ID: GRG) (formula: $C_{20}H_{36}O_7P_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O P 29 20 7 2	0	0
3	C	1	Total C O P 29 20 7 2	0	0
3	D	1	Total C O P 29 20 7 2	0	0
3	E	1	Total C O P 29 20 7 2	0	0
3	F	1	Total C O P 29 20 7 2	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	313	Total O 313 313	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	288	Total O 288 288	0	0
4	C	376	Total O 376 376	0	0
4	D	406	Total O 406 406	0	0
4	E	416	Total O 416 416	0	0
4	F	344	Total O 344 344	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Geranylgeranyl pyrophosphate synthase

Chain A:




• Molecule 1: Geranylgeranyl pyrophosphate synthase

Chain B:




• Molecule 1: Geranylgeranyl pyrophosphate synthase

Chain C:




• Molecule 1: Geranylgeranyl pyrophosphate synthase


Chain D:



• Molecule 1: Geranylgeranyl pyrophosphate synthase


Chain E:





• Molecule 1: Geranylgeranyl pyrophosphate synthase

Chain F:





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	86.19Å 135.65Å 146.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.54 – 1.82 39.54 – 1.82	Depositor EDS
% Data completeness (in resolution range)	99.9 (39.54-1.82) 96.1 (39.54-1.82)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.42 (at 1.82Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.169 , 0.202 0.170 , 0.204	Depositor DCC
R_{free} test set	2009 reflections (1.31%)	wwPDB-VP
Wilson B-factor (Å^2)	21.4	Xtriage
Anisotropy	0.239	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.28 , 33.2	EDS
L-test for twinning[2]	$< \|L\| > = 0.48$, $< L^2 > = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	16173	wwPDB-VP
Average B, all atoms (Å^2)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.50% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, GRG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| > 5$	RMSZ	$\#\|Z\| > 5$
1	A	0.16	0/2357	0.34	0/3190
1	B	0.18	0/2328	0.39	0/3152
1	C	0.18	0/2360	0.38	0/3196
1	D	0.18	0/2360	0.38	0/3193
1	E	0.18	0/2382	0.37	0/3223
1	F	0.17	0/2387	0.34	0/3231
All	All	0.17	0/14174	0.37	0/19185

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2307	0	2308	20	0
1	B	2278	0	2280	9	0
1	C	2310	0	2304	13	0
1	D	2310	0	2316	10	0
1	E	2332	0	2336	12	0
1	F	2336	0	2333	12	0
2	A	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
2	E	2	0	0	0	0
2	F	2	0	0	0	0
3	A	29	0	33	7	0
3	C	29	0	33	2	0
3	D	29	0	33	2	0
3	E	29	0	33	1	0
3	F	29	0	33	2	0
4	A	313	0	0	5	0
4	B	288	0	0	2	0
4	C	376	0	0	5	0
4	D	406	0	0	4	0
4	E	416	0	0	2	0
4	F	344	0	0	5	0
All	All	16173	0	14042	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 71 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:117:ARG:NH2	4:D:405:HOH:O	2.29	0.64
1:A:10:ARG:NH1	4:A:1008:HOH:O	2.32	0.63
1:C:170:LYS:NZ	4:C:406:HOH:O	2.31	0.63
1:A:6:GLU:OE2	1:A:17:LYS:NZ	2.33	0.60
1:B:9:GLN:NE2	4:B:404:HOH:O	2.31	0.60

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	279/294 (95%)	276 (99%)	3 (1%)	0	100	100
1	B	275/294 (94%)	274 (100%)	1 (0%)	0	100	100
1	C	280/294 (95%)	278 (99%)	2 (1%)	0	100	100
1	D	279/294 (95%)	276 (99%)	3 (1%)	0	100	100
1	E	282/294 (96%)	280 (99%)	2 (1%)	0	100	100
1	F	282/294 (96%)	279 (99%)	3 (1%)	0	100	100
All	All	1677/1764 (95%)	1663 (99%)	14 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	254/266 (96%)	253 (100%)	1 (0%)	84	81
1	B	251/266 (94%)	250 (100%)	1 (0%)	84	81
1	C	254/266 (96%)	253 (100%)	1 (0%)	84	81
1	D	254/266 (96%)	254 (100%)	0	100	100
1	E	258/266 (97%)	258 (100%)	0	100	100
1	F	258/266 (97%)	258 (100%)	0	100	100
All	All	1529/1596 (96%)	1526 (100%)	3 (0%)	87	86

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	121	GLU
1	B	27	VAL
1	C	27	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such

sidechains are listed below:

Mol	Chain	Res	Type
1	F	236	GLN
1	F	262	ASN
1	E	124	GLN
1	E	274	GLN
1	F	58	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 17 ligands modelled in this entry, 12 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	GRG	F	303	2	27,28,28	4.14	9 (33%)	33,37,37	1.35	4 (12%)
3	GRG	C	303	2	27,28,28	4.11	9 (33%)	33,37,37	1.33	5 (15%)
3	GRG	A	303	2	27,28,28	4.13	9 (33%)	33,37,37	1.37	4 (12%)
3	GRG	E	303	2	27,28,28	4.10	9 (33%)	33,37,37	1.35	4 (12%)
3	GRG	D	303	2	27,28,28	4.07	9 (33%)	33,37,37	1.20	3 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GRG	F	303	2	-	17/31/31/31	-
3	GRG	C	303	2	-	12/31/31/31	-
3	GRG	A	303	2	-	13/31/31/31	-
3	GRG	E	303	2	-	17/31/31/31	-
3	GRG	D	303	2	-	16/31/31/31	-

The worst 5 of 45 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	303	GRG	C7-C8	9.17	1.54	1.33
3	A	303	GRG	C7-C8	9.17	1.54	1.33
3	C	303	GRG	C7-C8	9.10	1.54	1.33
3	D	303	GRG	C7-C8	9.07	1.53	1.33
3	F	303	GRG	C7-C8	9.05	1.53	1.33

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	A	303	GRG	C4-C3-C5	3.28	120.92	115.23
3	E	303	GRG	C4-C3-C5	3.26	120.88	115.23
3	C	303	GRG	C4-C3-C5	3.24	120.85	115.23
3	F	303	GRG	C4-C3-C5	3.23	120.84	115.23
3	A	303	GRG	C14-C13-C15	3.09	120.59	115.23

There are no chirality outliers.

5 of 75 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	303	GRG	C11-C12-C13-C14
3	A	303	GRG	C11-C12-C13-C15
3	A	303	GRG	C6-C7-C8-C10
3	A	303	GRG	C1-C2-C3-C4
3	D	303	GRG	C1-C2-C3-C5


There are no ring outliers.


5 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	F	303	GRG	2	0
3	C	303	GRG	2	0
3	A	303	GRG	7	0
3	E	303	GRG	1	0
3	D	303	GRG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	283/294 (96%)	-0.02	11 (3%)	43	49	13, 24, 47, 75	0
1	B	279/294 (94%)	-0.04	9 (3%)	50	57	15, 24, 46, 73	0
1	C	284/294 (96%)	-0.27	4 (1%)	73	79	13, 20, 40, 57	0
1	D	283/294 (96%)	-0.25	4 (1%)	73	79	13, 21, 39, 70	0
1	E	286/294 (97%)	-0.28	2 (0%)	84	88	13, 21, 35, 48	0
1	F	286/294 (97%)	-0.01	7 (2%)	59	66	14, 24, 53, 72	0
All	All	1701/1764 (96%)	-0.14	37 (2%)	62	69	13, 22, 45, 75	0

The worst 5 of 37 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	8	VAL	4.8
1	C	203	SER	4.0
1	B	291	LEU	3.5
1	B	8	VAL	3.2
1	A	20	LEU	3.2

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no oligosaccharides in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	MG	B	302	1/1	0.87	0.09	34,34,34,34	0
2	MG	B	301	1/1	0.92	0.06	37,37,37,37	0
3	GRG	A	303	29/29	0.92	0.12	17,39,56,58	0
3	GRG	C	303	29/29	0.92	0.12	16,33,58,72	0
3	GRG	F	303	29/29	0.94	0.11	16,35,51,56	0
3	GRG	E	303	29/29	0.95	0.11	14,31,43,50	0
3	GRG	D	303	29/29	0.95	0.11	13,32,47,50	0
2	MG	F	302	1/1	0.97	0.03	18,18,18,18	0
2	MG	E	301	1/1	0.97	0.03	16,16,16,16	0
2	MG	F	301	1/1	0.97	0.03	18,18,18,18	0
2	MG	C	301	1/1	0.99	0.01	18,18,18,18	0
2	MG	C	302	1/1	0.99	0.02	16,16,16,16	0
2	MG	D	301	1/1	0.99	0.03	16,16,16,16	0
2	MG	A	301	1/1	0.99	0.04	18,18,18,18	0
2	MG	E	302	1/1	0.99	0.02	15,15,15,15	0
2	MG	A	302	1/1	0.99	0.04	19,19,19,19	0
2	MG	D	302	1/1	1.00	0.03	15,15,15,15	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around GRG A 303:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)










Electron density around GRG C 303:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)










Electron density around GRG F 303:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)










Electron density around GRG E 303:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)








Electron density around GRG D 303:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



6.5 Other polymers

There are no such residues in this entry.