



wwPDB EM Validation Summary Report ⓘ

Oct 19, 2026 – 04:14 PM EDT

PDB ID : 12ZR / pdb_000012zr
EMDB ID : EMD-76894
Title : Cryo-EM structure of hexameric apo human K293st geranylgeranyl diphosphate synthase
Authors : Pham, A.C.; Borgstahl, E.O.
Deposited on : 2026-04-24
Resolution : 2.30 Å(reported)
Based on initial model : 6R4V

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev133
MolProbity : 4-5-2 with Phenix2.2
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.51

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	4254 (1.80 - 2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	293	
1	B	293	
1	C	293	
1	D	293	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	293	<div><div></div><div>7%</div><div>57%</div><div>16%</div><div>27%</div></div>
1	F	293	<div><div></div><div>86%</div><div>12%</div><div></div></div>

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 12740 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Geranylgeranyl pyrophosphate synthase.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	285	Total	C	N	O	S	0	0
			2332	1500	391	435	6		
1	B	212	Total	C	N	O	S	0	0
			1718	1115	286	313	4		
1	C	288	Total	C	N	O	S	0	0
			2328	1497	392	433	6		
1	D	286	Total	C	N	O	S	0	0
			2312	1489	389	428	6		
1	E	214	Total	C	N	O	S	0	0
			1731	1122	288	317	4		
1	F	287	Total	C	N	O	S	0	0
			2319	1492	390	431	6		


There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP O95749
B	0	SER	-	expression tag	UNP O95749
C	0	SER	-	expression tag	UNP O95749
D	0	SER	-	expression tag	UNP O95749
E	0	SER	-	expression tag	UNP O95749
F	0	SER	-	expression tag	UNP O95749


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Geranylgeranyl pyrophosphate synthase



- Molecule 1: Geranylgeranyl pyrophosphate synthase



- Molecule 1: Geranylgeranyl pyrophosphate synthase



- Molecule 1: Geranylgeranyl pyrophosphate synthase

Chain F: 



Position	Conservation (bits)	Most Frequent Residue
1	1.8	GLN
2	1.8	GLN
3	1.8	GLN
4	1.8	GLN
5	1.8	GLN
6	1.8	GLN
7	1.8	GLN
8	1.8	GLN
9	1.8	GLN
10	1.8	GLN
11	0.2	GLN
12	0.2	GLN
13	0.2	GLN
14	0.2	GLN
15	0.2	GLN
16	0.2	GLN
17	0.2	GLN
18	0.2	GLN
19	0.2	GLN
20	0.2	GLN
21	0.2	GLN
22	0.2	GLN
23	0.2	GLN
24	0.2	GLN
25	0.2	GLN
26	0.2	GLN
27	0.2	GLN
28	0.2	GLN
29	0.2	GLN
30	0.2	GLN
31	0.2	GLN
32	0.2	GLN
33	0.2	GLN
34	0.2	GLN
35	0.2	GLN
36	0.2	GLN
37	0.2	GLN
38	0.2	GLN
39	0.2	GLN
40	0.2	GLN
41	0.2	GLN
42	0.2	GLN
43	0.2	GLN
44	0.2	GLN
45	0.2	GLN
46	0.2	GLN
47	0.2	GLN
48	0.2	GLN
49	0.2	GLN
50	0.2	GLN
51	0.2	GLN
52	0.2	GLN
53	0.2	GLN
54	0.2	GLN
55	0.2	GLN
56	0.2	GLN
57	0.2	GLN
58	0.2	GLN
59	0.2	GLN
60	0.2	GLN
61	0.2	GLN
62	0.2	GLN
63	0.2	GLN
64	0.2	GLN
65	0.2	GLN
66	0.2	GLN
67	0.2	GLN
68	0.2	GLN
69	0.2	GLN
70	0.2	GLN
71	0.2	GLN
72	0.2	GLN
73	0.2	GLN
74	0.2	GLN
75	0.2	GLN
76	0.2	GLN
77	0.2	GLN
78	0.2	GLN
79	0.2	GLN
80	0.2	GLN
81	0.2	GLN
82	0.2	GLN
83	0.2	GLN
84	0.2	GLN
85	0.2	GLN
86	0.2	GLN
87	0.2	GLN
88	0.2	GLN
89	0.2	GLN
90	0.2	GLN
91	0.2	GLN
92	0.2	GLN
93	0.2	GLN
94	0.2	GLN
95	0.2	GLN
96	0.2	GLN
97	0.2	GLN
98	0.2	GLN
99	0.2	GLN
100	0.2	GLN
101	0.2	GLN
102	0.2	GLN
103	0.2	GLN
104	0.2	GLN
105	0.2	GLN
106	0.2	GLN
107	0.2	GLN
108	0.2	GLN
109	0.2	GLN
110	0.2	GLN
111	0.2	GLN
112	0.2	GLN
113	0.2	GLN
114	0.2	GLN
115	0.2	GLN
116	0.2	GLN
117	0.2	GLN
118	0.2	GLN
119	0.2	GLN
120	0.2	GLN
121	0.2	GLN
122	0.2	GLN
123	0.2	GLN
124	0.2	GLN
125	0.2	GLN
126	0.2	GLN
127	0.2	GLN
128	0.2	GLN
129	0.2	GLN
130	0.2	GLN
131	0.2	GLN
132	0.2	GLN
133	0.2	GLN
134	0.2	GLN
135	0.2	GLN
136	0.2	GLN
137	0.2	GLN
138	0.2	GLN
139	0.2	GLN
140	0.2	GLN
141	0.2	GLN
142	0.2	GLN
143	0.2	GLN

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C2	Depositor
Number of particles used	1142326	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	47	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	1900	Depositor
Magnification	165000	Depositor
Image detector	TFS FALCON 4i (4k x 4k)	Depositor
Maximum map value	0.565	Depositor
Minimum map value	-0.208	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.012	Depositor
Recommended contour level	0.09	Depositor
Map size (Å)	300.5952, 300.5952, 300.5952	wwPDB
Map dimensions	412, 412, 412	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.7296, 0.7296, 0.7296	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.31	0/2384	0.48	0/3227
1	B	0.19	0/1753	0.35	0/2372
1	C	0.26	0/2379	0.42	0/3224
1	D	0.20	0/2363	0.35	0/3203
1	E	0.28	0/1766	0.41	0/2390
1	F	0.18	0/2370	0.32	0/3212
All	All	0.24	0/13015	0.39	0/17628

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	2
1	C	0	1
1	E	0	2
1	F	0	3
All	All	0	11

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 11 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	10	ARG	Sidechain
1	A	187	ARG	Sidechain
1	A	74	ARG	Sidechain
1	B	117	ARG	Sidechain
1	B	74	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2332	0	2328	23	0
1	B	1718	0	1755	20	0
1	C	2328	0	2308	24	0
1	D	2312	0	2292	20	0
1	E	1731	0	1761	29	0
1	F	2319	0	2300	21	0
All	All	12740	0	12744	130	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 130 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:22:LEU:HB3	1:A:74:ARG:HD2	1.75	0.68
1:E:41:VAL:HG21	1:E:163:MET:HB3	1.76	0.67
1:C:41:VAL:HG21	1:C:163:MET:HB3	1.80	0.64
1:C:237:ARG:HH11	1:C:237:ARG:HG3	1.64	0.62
1:F:41:VAL:HG21	1:F:163:MET:HB3	1.82	0.61

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	283/293 (97%)	274 (97%)	9 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	208/293 (71%)	205 (99%)	3 (1%)	0	100	100
1	C	286/293 (98%)	282 (99%)	4 (1%)	0	100	100
1	D	284/293 (97%)	278 (98%)	6 (2%)	0	100	100
1	E	210/293 (72%)	204 (97%)	6 (3%)	0	100	100
1	F	285/293 (97%)	279 (98%)	6 (2%)	0	100	100
All	All	1556/1758 (88%)	1522 (98%)	34 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	257/265 (97%)	256 (100%)	1 (0%)	84	92
1	B	188/265 (71%)	186 (99%)	2 (1%)	65	81
1	C	253/265 (96%)	253 (100%)	0	100	100
1	D	250/265 (94%)	250 (100%)	0	100	100
1	E	189/265 (71%)	188 (100%)	1 (0%)	81	90
1	F	252/265 (95%)	250 (99%)	2 (1%)	73	86
All	All	1389/1590 (87%)	1383 (100%)	6 (0%)	81	92

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	73	ARG
1	F	27	VAL
1	F	289	LYS
1	B	103	VAL
1	A	70	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	21	GLN
1	C	118	GLN
1	F	33	GLN
1	D	150	GLN
1	E	69	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-76894. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y



Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 206




Y Index: 206




Z Index: 206


6.2.2 Raw map



X Index: 206



Y Index: 206




Z Index: 206


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 171




Y Index: 208




Z Index: 184


6.3.2 Raw map



X Index: 171



Y Index: 204




Z Index: 183


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.09. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_76894_msk_1.map [i](#)



X




Y




Z


6.6.2 emd_76894_msk_2.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 76 nm³; this corresponds to an approximate mass of 69 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.435 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.435 $\text{\AA}^{-1}$

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.30	-	-
Author-provided FSC curve	2.30	2.71	2.35
Unmasked-calculated*	2.91	3.37	2.96

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 2.91 differs from the reported value 2.3 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-76894 and PDB model 12ZR. Per-residue inclusion information can be found in section [3](#) on page [5](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.09 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.09).

9.4 Atom inclusion ⓘ



At the recommended contour level, 94% of all backbone atoms, 87% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.09) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div><div></div></div> 0.8740	<div><div></div></div> 0.5810
A	<div><div></div></div> 0.8850	<div><div></div></div> 0.5870
B	<div><div></div></div> 0.7790	<div><div></div></div> 0.5070
C	<div><div></div></div> 0.9290	<div><div></div></div> 0.6190
D	<div><div></div></div> 0.9400	<div><div></div></div> 0.6270
E	<div><div></div></div> 0.7690	<div><div></div></div> 0.5090
F	<div><div></div></div> 0.8920	<div><div></div></div> 0.5990

1.0

0.0

<0.0