wwPDB X-ray Structure Validation Summary Report (i)

Oct 19, 2026 – 06:56 PM JST

PDB ID	:	26BE / pdb_000026be
Title	:	X-ray crystal structure of human NBS1 FHA-BRCT1-BRCT2 domain
Authors	:	Ite, K.; Kitanishi, K.; Unno, M.
Deposited on	:	2026-04-24
Resolution	:	2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.2
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	2.2
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	334	4% 84% 9% 6%
1	B	334	2% 88% 6% • •
1	C	334	4% 86% 8% • 5%
1	D	334	2% 87% 7% • •

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 10441 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nibrin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	313	Total	C	N	O	S	0	1	0
			2460	1571	414	460	15			
1	B	320	Total	C	N	O	S	0	1	0
			2503	1597	420	471	15			
1	C	316	Total	C	N	O	S	0	3	0
			2499	1597	419	468	15			
1	D	319	Total	C	N	O	S	0	0	0
			2488	1587	418	468	15			

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	327	LEU	-	expression tag	UNP O60934
A	328	GLU	-	expression tag	UNP O60934
A	329	HIS	-	expression tag	UNP O60934
A	330	HIS	-	expression tag	UNP O60934
A	331	HIS	-	expression tag	UNP O60934
A	332	HIS	-	expression tag	UNP O60934
A	333	HIS	-	expression tag	UNP O60934
A	334	HIS	-	expression tag	UNP O60934
B	327	LEU	-	expression tag	UNP O60934
B	328	GLU	-	expression tag	UNP O60934
B	329	HIS	-	expression tag	UNP O60934
B	330	HIS	-	expression tag	UNP O60934
B	331	HIS	-	expression tag	UNP O60934
B	332	HIS	-	expression tag	UNP O60934
B	333	HIS	-	expression tag	UNP O60934
B	334	HIS	-	expression tag	UNP O60934
C	327	LEU	-	expression tag	UNP O60934
C	328	GLU	-	expression tag	UNP O60934
C	329	HIS	-	expression tag	UNP O60934
C	330	HIS	-	expression tag	UNP O60934
C	331	HIS	-	expression tag	UNP O60934

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
C	332	HIS	-	expression tag	UNP O60934
C	333	HIS	-	expression tag	UNP O60934
C	334	HIS	-	expression tag	UNP O60934
D	327	LEU	-	expression tag	UNP O60934
D	328	GLU	-	expression tag	UNP O60934
D	329	HIS	-	expression tag	UNP O60934
D	330	HIS	-	expression tag	UNP O60934
D	331	HIS	-	expression tag	UNP O60934
D	332	HIS	-	expression tag	UNP O60934
D	333	HIS	-	expression tag	UNP O60934
D	334	HIS	-	expression tag	UNP O60934

- Molecule 2 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O P 5 4 1	0	0
2	B	1	Total O P 5 4 1	0	0
2	C	1	Total O P 5 4 1	0	0
2	D	1	Total O P 5 4 1	0	0

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0
3	D	1	Total O S 5 4 1	0	0

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	D	1	Total C O 4 2 2	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	88	Total O 88 88	0	0
5	B	135	Total O 135 135	0	0
5	C	107	Total O 107 107	0	0
5	D	92	Total O 92 92	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Nibrin

Chain A:




- Molecule 1: Nibrin

Chain B:




- Molecule 1: Nibrin


Chain C:



- Molecule 1: Nibrin

Chain D:




K320
N321
Y322
C323
L327
GLU
HIS
HIS
HIS
HIS
HIS
HIS

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	69.26Å 115.17Å 124.58Å 90.00° 104.66° 90.00°	Depositor
Resolution (Å)	48.29 – 2.50 48.29 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.29-2.50) 99.9 (48.29-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.13 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.8.0430	Depositor
R, R_{free}	0.194 , 0.237 0.200 , 0.238	Depositor DCC
R_{free} test set	3290 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å^2)	48.3	Xtriage
Anisotropy	0.041	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.32 , 33.5	EDS
L-test for twinning[2]	$< \|L\| > = 0.51$, $< L^2 > = 0.34$	Xtriage
Estimated twinning fraction	0.013 for h,-k,-h-l	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	10441	wwPDB-VP
Average B, all atoms (Å^2)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.08% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, EDO, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|$ >5	RMSZ	#$\|Z\|$ >5
1	A	0.66	0/2507	1.10	7/3387 (0.2%)
1	B	0.67	1/2554 (0.0%)	1.12	6/3452 (0.2%)
1	C	0.66	0/2547	1.14	8/3439 (0.2%)
1	D	0.66	0/2536	1.11	2/3427 (0.1%)
All	All	0.66	1/10144 (0.0%)	1.12	23/13705 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	3
1	D	0	1
All	All	0	5

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	299	ARG	NE-CZ	5.25	1.38	1.33

The worst 5 of 23 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	D	26	VAL	N-CA-CB	6.97	119.36	110.99
1	B	299	ARG	NE-CZ-NH2	6.38	124.94	119.20
1	D	244	PHE	CA-CB-CG	6.17	119.97	113.80
1	C	258	GLU	N-CA-CB	5.95	118.72	109.97
1	C	36	GLU	CB-CG-CD	5.83	122.51	112.60

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	43	ARG	Sidechain
1	B	250	ARG	Sidechain
1	B	28	ARG	Sidechain
1	B	53	SER	Peptide
1	D	28	ARG	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2460	0	2490	16	0
1	B	2503	0	2533	13	0
1	C	2499	0	2527	12	0
1	D	2488	0	2511	15	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
2	D	5	0	0	0	0
3	A	5	0	0	0	0
3	B	5	0	0	1	0
3	C	10	0	0	1	0
3	D	5	0	0	0	0
4	A	4	0	6	0	0
4	B	16	0	24	0	0
4	D	4	0	6	0	0
5	A	88	0	0	1	0
5	B	135	0	0	4	0
5	C	107	0	0	1	0
5	D	92	0	0	2	0
All	All	10441	0	10097	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 55 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:289:TRP:CZ2	1:B:293:ILE:HD11	2.18	0.79
1:A:35:ILE:O	1:A:41:ILE:CD1	2.36	0.74
1:A:38:ASP:O	1:A:41:ILE:HG12	1.91	0.71
1:A:70:ASP:HB2	1:A:83:MET:HE3	1.73	0.69
1:D:101:VAL:HG12	1:D:101:VAL:O	1.93	0.67

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	308/334 (92%)	297 (96%)	10 (3%)	1 (0%)	36	55
1	B	315/334 (94%)	306 (97%)	8 (2%)	1 (0%)	36	55
1	C	313/334 (94%)	303 (97%)	10 (3%)	0	100	100
1	D	313/334 (94%)	301 (96%)	11 (4%)	1 (0%)	36	55
All	All	1249/1336 (94%)	1207 (97%)	39 (3%)	3 (0%)	43	63

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	321	ASN
1	B	321	ASN
1	D	321	ASN

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	276/292 (94%)	269 (98%)	7 (2%)	42	69
1	B	281/292 (96%)	271 (96%)	10 (4%)	31	58
1	C	280/292 (96%)	267 (95%)	13 (5%)	24	48
1	D	278/292 (95%)	270 (97%)	8 (3%)	37	65
All	All	1115/1168 (96%)	1077 (97%)	38 (3%)	32	60

5 of 38 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	258	GLU
1	D	219	LYS
1	C	298	GLN
1	D	92	LYS
1	D	250	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	220	GLN
1	D	134	GLN
1	C	216	GLN
1	D	30	ASN
1	C	134	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	PO4	D	401	-	4,4,4	1.46	1 (25%)	6,6,6	0.47	0
4	EDO	D	403	-	3,3,3	0.37	0	2,2,2	0.41	0
2	PO4	B	401	-	4,4,4	1.06	0	6,6,6	0.66	0
3	SO4	B	402	-	4,4,4	0.30	0	6,6,6	0.37	0
3	SO4	D	402	-	4,4,4	0.30	0	6,6,6	0.13	0
3	SO4	C	402	-	4,4,4	0.25	0	6,6,6	0.26	0
4	EDO	B	403	-	3,3,3	0.25	0	2,2,2	0.40	0
3	SO4	A	402	-	4,4,4	0.33	0	6,6,6	0.09	0
4	EDO	B	405	-	3,3,3	0.45	0	2,2,2	0.58	0
4	EDO	B	406	-	3,3,3	0.36	0	2,2,2	0.18	0
4	EDO	A	403	-	3,3,3	0.29	0	2,2,2	0.42	0
3	SO4	C	403	-	4,4,4	0.34	0	6,6,6	0.07	0
2	PO4	A	401	-	4,4,4	1.30	0	6,6,6	0.47	0
4	EDO	B	404	-	3,3,3	0.33	0	2,2,2	0.13	0
2	PO4	C	401	-	4,4,4	0.87	0	6,6,6	0.43	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	B	403	-	-	0/1/1/1	-
4	EDO	B	405	-	-	1/1/1/1	-
4	EDO	B	406	-	-	0/1/1/1	-
4	EDO	A	403	-	-	1/1/1/1	-
4	EDO	D	403	-	-	1/1/1/1	-
4	EDO	B	404	-	-	0/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	401	PO4	P-O1	2.20	1.56	1.50

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	403	EDO	O1-C1-C2-O2
4	D	403	EDO	O1-C1-C2-O2
4	B	405	EDO	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	402	SO4	1	0
3	C	403	SO4	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	313/334 (93%)	0.11	12 (3%) 44 39	36, 53, 85, 104	1 (0%)
1	B	320/334 (95%)	-0.22	7 (2%) 62 58	23, 45, 75, 113	1 (0%)
1	C	316/334 (94%)	0.15	13 (4%) 41 37	21, 49, 92, 130	3 (0%)
1	D	319/334 (95%)	-0.12	7 (2%) 62 58	35, 49, 80, 112	0
All	All	1268/1336 (94%)	-0.02	39 (3%) 51 47	21, 49, 85, 130	5 (0%)

The worst 5 of 39 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	327	LEU	5.0
1	D	60	THR	4.4
1	C	327	LEU	4.2
1	B	10	PRO	4.0
1	B	54	VAL	3.7

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	SO4	A	402	5/5	0.51	0.11	138,140,154,171	0
4	EDO	B	403	4/4	0.53	0.30	73,95,96,97	0
3	SO4	C	403	5/5	0.57	0.10	123,142,151,153	0
3	SO4	C	402	5/5	0.71	0.11	82,84,113,121	0
4	EDO	D	403	4/4	0.80	0.18	58,77,77,80	0
4	EDO	A	403	4/4	0.83	0.19	69,71,75,84	0
4	EDO	B	404	4/4	0.85	0.22	77,78,78,86	0
4	EDO	B	405	4/4	0.90	0.19	66,74,78,81	0
4	EDO	B	406	4/4	0.93	0.14	54,66,66,72	0
3	SO4	D	402	5/5	0.93	0.08	66,67,93,95	0
3	SO4	B	402	5/5	0.96	0.07	53,56,73,82	0
2	PO4	B	401	5/5	0.97	0.09	41,47,48,50	0
2	PO4	C	401	5/5	0.98	0.04	41,45,47,56	0
2	PO4	A	401	5/5	0.98	0.06	45,45,52,60	0
2	PO4	D	401	5/5	0.99	0.05	38,42,50,58	0

6.5 Other polymers (i)

There are no such residues in this entry.