



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 8, 2026 – 04:32 PM EDT

PDB ID : 38HJ / pdb_000038hj
Title : Crystal structure of the transpeptidase domain of a Y422F mutant of PBP2 from *Neisseria gonorrhoeae* strain H041 acylated by cefoperazone
Authors : Bala, S.; Stratton, C.M.; Davies, C.
Deposited on : 2026-08-27
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	330	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2491 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Probable peptidoglycan D,D-transpeptidase PenA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	324	Total	C	N	O	S	0	0	0
			2452	1562	425	456	9			

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	232	GLY	-	expression tag	UNP F2Z7K9
A	233	SER	-	expression tag	UNP F2Z7K9
A	234	GLY	-	expression tag	UNP F2Z7K9
A	235	GLY	-	expression tag	UNP F2Z7K9
A	236	ALA	-	expression tag	UNP F2Z7K9
A	297	GLY	ALA	conflict	UNP F2Z7K9
A	?	-	TYR	deletion	UNP F2Z7K9
A	?	-	GLU	deletion	UNP F2Z7K9
A	?	-	PRO	deletion	UNP F2Z7K9
A	?	-	ASN	deletion	UNP F2Z7K9
A	?	-	LYS	deletion	UNP F2Z7K9
A	?	-	PRO	deletion	UNP F2Z7K9
A	?	-	GLY	deletion	UNP F2Z7K9
A	?	-	GLN	deletion	UNP F2Z7K9
A	?	-	ALA	deletion	UNP F2Z7K9
A	?	-	ASP	deletion	UNP F2Z7K9
A	?	-	SER	deletion	UNP F2Z7K9
A	?	-	GLU	deletion	UNP F2Z7K9
A	?	-	GLN	deletion	UNP F2Z7K9
A	?	-	ARG	deletion	UNP F2Z7K9
A	422	PHE	TYR	engineered mutation	UNP F2Z7K9

- Molecule 2 is (2R)-2-[(1R)-1-{[(2R)-2-[(4-ethyl-2,3-dioxopiperazine-1-carbonyl)amino]-2-(4-hydroxyphenyl)acetyl]amino}-2-oxoethyl]-5-methyl-3,6-dihydro-2H-1,3-thiazine-4-carboxylic acid (CCD ID: A1DZ4) (formula: C₂₃H₂₇N₅O₈S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			37	23	5	8	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	O	0	0
			2	2		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Probable peptidoglycan D,D-transpeptidase PenA



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	50.60Å 61.60Å 110.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.10 – 2.60 39.10 – 2.60	Depositor EDS
% Data completeness (in resolution range)	92.6 (39.10-2.60) 92.6 (39.10-2.60)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.30 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.8.0430	Depositor
R, R_{free}	0.227 , 0.292 0.230 , 0.292	Depositor DCC
R_{free} test set	538 reflections (4.84%)	wwPDB-VP
Wilson B-factor (Å ²)	39.8	Xtriage
Anisotropy	0.028	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 14.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	2491	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: A1DZ4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.63	0/2502	1.18	9/3395 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	536	ASP	CA-CB-CG	6.63	119.23	112.60
1	A	569	THR	CA-CB-OG1	-6.34	100.08	109.60
1	A	327	ASP	CA-CB-CG	6.24	118.84	112.60
1	A	406	ASN	CA-CB-CG	-6.01	106.59	112.60
1	A	449	PHE	CB-CA-C	5.59	119.61	110.17
1	A	272	THR	CA-CB-OG1	-5.34	101.59	109.60
1	A	330	ASP	CA-CB-CG	5.34	117.94	112.60
1	A	315	PHE	CA-CB-CG	-5.33	108.47	113.80
1	A	327	ASP	CB-CA-C	5.19	120.42	109.79

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	528	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2452	0	2502	30	0
2	A	37	0	0	0	0
3	A	2	0	0	0	0
All	All	2491	0	2502	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:264:THR:HG21	1:A:301:ALA:HB2	1.79	0.65
1:A:394:SER:OG	1:A:399:GLU:OE2	2.07	0.61
1:A:264:THR:CG2	1:A:301:ALA:HB2	2.32	0.58
1:A:244:GLN:NE2	1:A:279:VAL:HG12	2.19	0.58
1:A:502:ARG:HG2	1:A:502:ARG:HH11	1.72	0.54
1:A:330:ASP:O	1:A:354:ASP:HA	2.08	0.53
1:A:435:TYR:CE1	1:A:519:ILE:HD13	2.44	0.53
1:A:526:ASN:O	1:A:528:ARG:NH1	2.41	0.52
1:A:470:VAL:O	1:A:474:MET:HG3	2.11	0.51
1:A:322:ASP:HB2	1:A:466:THR:HG21	1.96	0.48
1:A:246:LEU:HD21	1:A:559:GLY:HA3	1.95	0.47
1:A:428:LEU:HD13	1:A:534:THR:HG23	1.95	0.47
1:A:504:LEU:HD13	1:A:509:TYR:CZ	2.51	0.46
1:A:437:VAL:O	1:A:442:GLY:HA2	2.18	0.43
1:A:466:THR:O	1:A:470:VAL:HG23	2.18	0.43
1:A:278:LEU:HD23	1:A:278:LEU:C	2.44	0.43
1:A:510:VAL:HB	1:A:512:TYR:CE2	2.54	0.42
1:A:551:PRO:O	1:A:555:GLN:HG2	2.20	0.42
1:A:338:LYS:HA	1:A:342:ALA:O	2.20	0.42
1:A:435:TYR:CD1	1:A:519:ILE:HD13	2.55	0.42
1:A:518:PHE:HB2	1:A:549:THR:OG1	2.19	0.41
1:A:359:MET:HE3	1:A:474:MET:HG2	2.02	0.41
1:A:428:LEU:HD13	1:A:534:THR:CG2	2.49	0.41
1:A:501:ALA:O	1:A:514:HIS:HA	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:382:PHE:CE2	1:A:386:LEU:HD11	2.56	0.41
1:A:259:GLN:HA	1:A:259:GLN:HE21	1.86	0.41
1:A:440:HIS:O	1:A:440:HIS:CG	2.70	0.41
1:A:355:VAL:HA	1:A:358:ILE:HD12	2.01	0.41
1:A:254:ALA:HB2	1:A:552:VAL:HG22	2.02	0.40
1:A:558:GLY:O	1:A:559:GLY:C	2.64	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	322/330 (98%)	309 (96%)	12 (4%)	1 (0%)	36 58

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	512	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	262/265 (99%)	257 (98%)	5 (2%)	50 75

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	281	THR
1	A	447	VAL
1	A	459	LYS
1	A	508	ARG
1	A	529	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	258	HIS
1	A	259	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	A1DZ4	A	601	1	36,39,39	3.85	17 (47%)	42,55,55	2.29	22 (52%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1DZ4	A	601	1	-	3/26/62/62	0/2/3/3

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	A1DZ4	C06-S07	-11.09	1.55	1.82
2	A	601	A1DZ4	C23-N24	8.70	1.62	1.47
2	A	601	A1DZ4	C08-C09	-7.42	1.42	1.51
2	A	601	A1DZ4	C09-C04	7.29	1.43	1.34
2	A	601	A1DZ4	C22-N21	6.44	1.58	1.47
2	A	601	A1DZ4	C10-C09	6.20	1.60	1.50
2	A	601	A1DZ4	C23-C22	-4.98	1.33	1.51
2	A	601	A1DZ4	C15-N14	4.86	1.44	1.34
2	A	601	A1DZ4	C19-N18	4.32	1.44	1.35
2	A	601	A1DZ4	C04-N05	3.38	1.45	1.37
2	A	601	A1DZ4	C04-C02	3.32	1.53	1.48
2	A	601	A1DZ4	C27-N24	-2.77	1.32	1.35
2	A	601	A1DZ4	C29-C27	2.70	1.57	1.53
2	A	601	A1DZ4	C19-N21	2.65	1.46	1.41
2	A	601	A1DZ4	C29-N21	-2.38	1.36	1.40
2	A	601	A1DZ4	O28-C27	-2.09	1.18	1.23
2	A	601	A1DZ4	C17-N18	-2.08	1.42	1.46

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	A1DZ4	C08-S07-C06	5.46	104.36	94.36
2	A	601	A1DZ4	C12-C11-N14	-4.48	102.74	109.80
2	A	601	A1DZ4	C25-N24-C27	3.73	124.06	119.59
2	A	601	A1DZ4	C36-C37-C31	-3.63	117.56	121.18
2	A	601	A1DZ4	C10-C09-C08	3.01	117.84	113.13
2	A	601	A1DZ4	C31-C17-C15	2.89	115.24	108.42
2	A	601	A1DZ4	C37-C31-C32	2.89	121.89	118.30
2	A	601	A1DZ4	C23-C22-N21	2.85	113.47	109.06
2	A	601	A1DZ4	C22-C23-N24	2.81	116.00	110.42
2	A	601	A1DZ4	C23-N24-C27	-2.75	115.31	120.82
2	A	601	A1DZ4	C09-C08-S07	-2.62	113.57	117.00
2	A	601	A1DZ4	C06-C11-N14	2.59	115.41	109.93

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	A1DZ4	C11-N14-C15	-2.51	120.20	123.08
2	A	601	A1DZ4	O01-C02-O03	-2.50	117.95	123.90
2	A	601	A1DZ4	O13-C12-C11	-2.48	118.29	124.86
2	A	601	A1DZ4	O20-C19-N21	2.47	122.44	119.59
2	A	601	A1DZ4	O01-C02-C04	2.40	120.53	116.73
2	A	601	A1DZ4	C08-C09-C04	-2.30	120.08	123.60
2	A	601	A1DZ4	C15-C17-N18	-2.29	101.50	108.55
2	A	601	A1DZ4	O30-C29-N21	2.25	123.65	119.99
2	A	601	A1DZ4	C32-C31-C17	-2.24	117.20	120.78
2	A	601	A1DZ4	C23-N24-C25	2.23	120.60	116.63

There are no chirality outliers.


All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	A1DZ4	O01-C02-C04-N05
2	A	601	A1DZ4	C12-C11-N14-C15
2	A	601	A1DZ4	O01-C02-C04-C09

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	324/330 (98%)	0.72	22 (6%) 23 19	26, 40, 63, 88	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	343	THR	4.3
1	A	342	ALA	4.1
1	A	505	VAL	3.3
1	A	506	ASN	3.3
1	A	527	PRO	2.9
1	A	412	PRO	2.8
1	A	573	THR	2.7
1	A	345	GLN	2.5
1	A	348	HIS	2.5
1	A	453	ALA	2.4
1	A	486	ALA	2.4
1	A	572	LEU	2.3
1	A	315	PHE	2.3
1	A	378	GLU	2.1
1	A	456	PRO	2.1
1	A	414	GLU	2.1
1	A	450	GLU	2.1
1	A	507	GLY	2.1
1	A	481	GLY	2.0
1	A	489	VAL	2.0
1	A	543	TYR	2.0
1	A	395	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	A1DZ4	A	601	37/37	0.85	0.12	31,43,68,69	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.