



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 22, 2026 – 07:25 PM EDT

PDB ID : 2A1Y / pdb_00002a1y
Title : Crystal Structure of GuaC-GMP complex from Bacillus anthracis at 2.26 Å Resolution.
Authors : Grenha, R.; Levnikov, V.M.; Blagova, E.V.; Fogg, M.J.; Brannigan, J.A.; Wilkinson, A.J.; Wilson, K.S.; Structural Proteomics in Europe (SPINE)
Deposited on : 2005-06-21
Resolution : 2.27 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	9078 (2.30-2.26)
Clashscore	190562	9802 (2.30-2.26)
Ramachandran outliers	187476	9690 (2.30-2.26)
Sidechain outliers	187428	9691 (2.30-2.26)
RSRZ outliers	180081	9085 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	336	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	5GP	A	1344	X	-	-	-

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2545 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called GMP reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	301	2347	1491	396	448	12	0	2	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-8	GLY	-	cloning artifact	UNP Q81JJ9
A	-7	SER	-	cloning artifact	UNP Q81JJ9
A	-6	SER	-	cloning artifact	UNP Q81JJ9
A	-5	HIS	-	expression tag	UNP Q81JJ9
A	-4	HIS	-	expression tag	UNP Q81JJ9
A	-3	HIS	-	expression tag	UNP Q81JJ9
A	-2	HIS	-	expression tag	UNP Q81JJ9
A	-1	HIS	-	expression tag	UNP Q81JJ9
A	0	HIS	-	expression tag	UNP Q81JJ9

- Molecule 2 is GUANOSINE-5'-MONOPHOSPHATE (CCD ID: 5GP) (formula: C₁₀H₁₄N₅O₈P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			24	10	5	8	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	174	Total	O	0	0
			174	174		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: GMP reductase



4 Data and refinement statistics

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, α , β , γ	95.61Å 95.61Å 67.67Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.26 – 2.27 30.26 – 2.27	Depositor EDS
% Data completeness (in resolution range)	97.5 (30.26-2.27) 97.6 (30.26-2.27)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.38 (at 2.26Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.167 , 0.246 0.190 , 0.251	Depositor DCC
R_{free} test set	696 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	42.2	Xtriage
Anisotropy	0.042	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 45.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.037 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2545	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 5GP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.38	11/2393 (0.5%)	1.30	15/3229 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	5

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	208	GLY	N-CA	8.33	1.53	1.45
1	A	220	ILE	CA-CB	-6.65	1.47	1.54
1	A	303	THR	CA-CB	6.08	1.62	1.54
1	A	49	ILE	CA-CB	5.67	1.62	1.54
1	A	320	ILE	CA-CB	5.42	1.61	1.54
1	A	150	THR	CA-CB	5.41	1.61	1.54
1	A	45	MET	C-O	5.37	1.30	1.23
1	A	216	VAL	C-O	-5.13	1.18	1.24
1	A	90	ILE	CA-CB	5.10	1.62	1.54
1	A	119	ILE	CA-CB	5.08	1.60	1.54
1	A	171	PRO	CA-C	5.01	1.58	1.52

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	170	GLY	CA-C-N	-7.30	112.35	120.45
1	A	170	GLY	C-N-CA	-7.30	112.35	120.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	4	VAL	CB-CA-C	6.37	119.50	110.84
1	A	279	LYS	N-CA-C	6.17	121.57	113.30
1	A	68	ARG	CB-CA-C	-6.16	100.30	110.72
1	A	13	ILE	CA-C-N	5.59	125.35	119.76
1	A	13	ILE	C-N-CA	5.59	125.35	119.76
1	A	67	HIS	N-CA-C	-5.58	103.33	110.53
1	A	320	ILE	N-CA-C	5.50	120.78	109.34
1	A	170	GLY	N-CA-C	5.40	123.36	112.34
1	A	239	SER	CA-C-N	5.19	125.19	119.89
1	A	239	SER	C-N-CA	5.19	125.19	119.89
1	A	144	ILE	CA-C-N	-5.16	114.57	122.62
1	A	144	ILE	C-N-CA	-5.16	114.57	122.62
1	A	197	ALA	N-CA-C	5.08	116.82	111.28

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	244	ILE	Peptide
1	A	249	LYS	Peptide
1	A	275	PHE	Peptide
1	A	318	ASN	Peptide
1	A	319	SER	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2347	0	2342	102	0
2	A	24	0	12	3	0
3	A	174	0	0	22	0
All	All	2545	0	2354	102	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 22.

All (102) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:43:ALA:CB	1:A:45:MET:HE2	1.64	1.24
1:A:132:GLN:HG2	3:A:1461:HOH:O	1.39	1.21
1:A:312:ASP:HB2	3:A:1485:HOH:O	1.44	1.14
1:A:244:ILE:HD11	1:A:275:PHE:CE1	1.86	1.11
1:A:43:ALA:HB3	1:A:45:MET:HE2	1.20	1.06
1:A:159:ASN:HB3	3:A:1459:HOH:O	1.58	1.04
1:A:236:HIS:HE1	1:A:285:THR:OG1	1.38	1.03
1:A:249:LYS:N	1:A:249:LYS:HE3	1.78	0.99
1:A:194:ARG:HD3	3:A:1486:HOH:O	1.61	0.97
1:A:70:GLN:HG3	3:A:1489:HOH:O	1.68	0.94
1:A:43:ALA:HB3	1:A:45:MET:CE	2.01	0.91
1:A:244:ILE:CD1	1:A:275:PHE:CE1	2.58	0.87
1:A:244:ILE:CD1	1:A:275:PHE:HE1	1.85	0.87
1:A:236:HIS:CE1	1:A:285:THR:OG1	2.29	0.82
1:A:202:LYS:HE3	3:A:1493:HOH:O	1.80	0.81
1:A:45:MET:HE3	1:A:48:ILE:CG1	2.13	0.79
1:A:121:HIS:HD2	3:A:1354:HOH:O	1.66	0.78
1:A:276:VAL:HG12	1:A:277:GLU:N	2.01	0.76
1:A:276:VAL:CG1	1:A:277:GLU:N	2.53	0.72
1:A:55:THR:OG1	1:A:81:MET:HE1	1.90	0.72
1:A:43:ALA:HB1	1:A:45:MET:HE2	1.65	0.71
1:A:244:ILE:HD11	1:A:275:PHE:HE1	1.35	0.71
1:A:136:LYS:HG2	3:A:1458:HOH:O	1.92	0.69
1:A:45:MET:HE3	1:A:48:ILE:HG12	1.75	0.68
1:A:187:GLY:HA2	3:A:1518:HOH:O	1.94	0.68
1:A:104:GLN:HG3	3:A:1481:HOH:O	1.94	0.68
1:A:45:MET:HE3	1:A:48:ILE:HG13	1.75	0.67
1:A:234:ALA:O	1:A:279:LYS:HE2	1.97	0.64
1:A:245:GLU:HB2	1:A:249:LYS:O	1.97	0.64
1:A:229:ILE:CG2	1:A:232:LEU:HB2	2.28	0.64
1:A:22:ARG:HG3	1:A:23:SER:N	2.12	0.64
1:A:175:CYS:SG	2:A:1344:5GP:N2	2.72	0.63
1:A:136:LYS:CG	3:A:1458:HOH:O	2.47	0.62
1:A:45:MET:SD	2:A:1344:5GP:H8	2.41	0.61
1:A:45:MET:CE	1:A:48:ILE:HG13	2.30	0.60
1:A:48:ILE:HD11	1:A:254:TYR:CE1	2.37	0.60
1:A:98:GLU:O	1:A:102:VAL:HG13	2.02	0.59
1:A:237[B]:GLU:HG2	1:A:278:HIS:HE1	1.68	0.59
1:A:26:ASP:HA	3:A:1371:HOH:O	2.03	0.59
1:A:79:ARG:HG3	1:A:110:LEU:HD11	1.84	0.58
1:A:150:THR:OG1	1:A:152[A]:GLU:HG2	2.02	0.58
1:A:159:ASN:CB	3:A:1459:HOH:O	2.30	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:142:PHE:CE1	1:A:163:ASP:HB3	2.40	0.57
1:A:251:TYR:HD2	1:A:276:VAL:O	1.87	0.57
1:A:102:VAL:HG21	1:A:133:HIS:CD2	2.39	0.56
1:A:175:CYS:SG	2:A:1344:5GP:C2	2.93	0.56
1:A:244:ILE:HD12	1:A:275:PHE:HE1	1.68	0.56
1:A:22:ARG:HG3	1:A:23:SER:H	1.69	0.56
1:A:95:LYS:HA	1:A:96:GLU:HG3	1.87	0.56
1:A:111:THR:HB	3:A:1517:HOH:O	2.05	0.55
1:A:3:ASN:HD22	1:A:3:ASN:N	2.05	0.53
1:A:176:ILE:HG23	1:A:180:LYS:HG2	1.91	0.53
1:A:321:PHE:C	3:A:1494:HOH:O	2.52	0.53
1:A:275:PHE:O	1:A:276:VAL:HG23	2.09	0.53
1:A:102:VAL:CG2	1:A:133:HIS:CD2	2.92	0.52
1:A:98:GLU:HA	1:A:101:PHE:HB3	1.92	0.52
1:A:198:LYS:HD2	3:A:1380:HOH:O	2.11	0.51
1:A:50:ASP:OD1	1:A:50:ASP:C	2.54	0.51
1:A:118:ASP:HB3	3:A:1424:HOH:O	2.10	0.51
1:A:185:THR:HA	1:A:188:TRP:NE1	2.26	0.51
1:A:276:VAL:CG1	1:A:277:GLU:H	2.23	0.51
1:A:229:ILE:HG21	1:A:232:LEU:HB2	1.93	0.50
1:A:29:VAL:HG22	1:A:30:THR:H	1.76	0.50
1:A:205:ILE:HG12	1:A:226:MET:HB3	1.92	0.50
1:A:55:THR:HG22	1:A:59:GLU:OE1	2.12	0.50
1:A:68:ARG:HD2	3:A:1387:HOH:O	2.12	0.49
1:A:194:ARG:HG2	3:A:1486:HOH:O	2.12	0.49
1:A:68:ARG:HG3	1:A:91:SER:HB2	1.95	0.49
1:A:64:TYR:CD1	1:A:64:TYR:C	2.91	0.48
1:A:97:ASP:O	1:A:100:GLU:HG2	2.13	0.48
1:A:34:HIS:HB2	1:A:114:TYR:OH	2.13	0.48
1:A:102:VAL:HG22	1:A:133:HIS:HE2	1.79	0.48
1:A:110:LEU:HD23	1:A:110:LEU:HA	1.65	0.48
1:A:77:PHE:CE1	1:A:81:MET:HE3	2.48	0.48
1:A:102:VAL:HG22	1:A:133:HIS:NE2	2.29	0.47
1:A:317:LYS:NZ	3:A:1441:HOH:O	2.46	0.47
1:A:249:LYS:NZ	1:A:250:LEU:H	2.13	0.47
1:A:77:PHE:CZ	1:A:81:MET:HG3	2.51	0.46
1:A:193:LEU:O	1:A:194:ARG:C	2.59	0.46
1:A:18:ILE:HD11	1:A:303:THR:HG23	1.97	0.45
1:A:150:THR:HB	1:A:152[A]:GLU:CD	2.43	0.44
1:A:94:VAL:HG11	1:A:126:ALA:HB1	1.99	0.44
1:A:243:THR:OG1	1:A:252:LYS:HE3	2.19	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:304:LYS:CE	3:A:1457:HOH:O	2.67	0.43
1:A:148:VAL:CG1	1:A:153:ALA:HB1	2.49	0.43
1:A:13:ILE:HA	1:A:14:PRO:HD3	1.73	0.42
1:A:148:VAL:HB	1:A:153:ALA:HB3	2.02	0.42
1:A:285:THR:O	1:A:289:MET:HG3	2.19	0.42
1:A:125:ASN:O	1:A:126:ALA:C	2.62	0.42
1:A:220:ILE:O	1:A:221:ARG:C	2.62	0.42
1:A:3:ASN:HD22	1:A:3:ASN:H	1.68	0.42
1:A:10:ILE:HD12	1:A:10:ILE:HG23	1.92	0.41
1:A:46:GLN:H	1:A:46:GLN:HG3	1.56	0.41
1:A:176:ILE:HG12	1:A:179:ILE:HD12	2.01	0.41
1:A:51:GLU:HG3	1:A:77:PHE:CD1	2.56	0.41
1:A:66:MET:HE3	1:A:67:HIS:CD2	2.56	0.41
1:A:100:GLU:HA	1:A:103:GLN:HB2	2.02	0.41
1:A:251:TYR:CD2	1:A:276:VAL:O	2.71	0.41
1:A:4:VAL:HG23	1:A:188:TRP:HB2	2.02	0.41
1:A:50:ASP:HA	1:A:66:MET:HE1	2.03	0.41
1:A:70:GLN:CG	3:A:1489:HOH:O	2.47	0.40
1:A:249:LYS:HZ1	1:A:250:LEU:H	1.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	297/336 (88%)	284 (96%)	10 (3%)	3 (1%)	12 13

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	320	ILE
1	A	23	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	94	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	253/280 (90%)	222 (88%)	31 (12%)	4 4

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	ASN
1	A	4	VAL
1	A	22	ARG
1	A	30	THR
1	A	37	LYS
1	A	46	GLN
1	A	52	ARG
1	A	70	GLN
1	A	72	GLU
1	A	75	ILE
1	A	84	ARG
1	A	98	GLU
1	A	102	VAL
1	A	108	GLU
1	A	140	GLU
1	A	152[A]	GLU
1	A	152[B]	GLU
1	A	173	LYS
1	A	176	ILE
1	A	194	ARG
1	A	195	TRP
1	A	198	LYS
1	A	202	LYS
1	A	232	LEU
1	A	242	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	244	ILE
1	A	245	GLU
1	A	249	LYS
1	A	275	PHE
1	A	277	GLU
1	A	320	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	3	ASN
1	A	46	GLN
1	A	61	ASN
1	A	82	GLN
1	A	109	HIS
1	A	121	HIS
1	A	189	GLN
1	A	236	HIS
1	A	278	HIS
1	A	318	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	5GP	A	1344	-	26,26,26	1.06	1 (3%)	39,40,40	2.00	14 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	5GP	A	1344	-	1/1/5/5	2/10/26/26	0/3/3/3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1344	5GP	C6-N1	-2.03	1.35	1.38

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1344	5GP	C5-C4-N3	-5.68	119.35	128.39
2	A	1344	5GP	C2-N3-C4	4.48	120.01	112.30
2	A	1344	5GP	N9-C4-N3	3.28	132.51	125.95
2	A	1344	5GP	O4'-C1'-C2'	-2.73	100.78	106.62
2	A	1344	5GP	C4'-O4'-C1'	2.58	115.15	109.47
2	A	1344	5GP	O4'-C1'-N9	2.54	114.12	108.36
2	A	1344	5GP	O6-C6-C5	-2.52	119.89	126.53
2	A	1344	5GP	C4-C5-N7	-2.40	106.87	110.67
2	A	1344	5GP	C2-N1-C6	-2.30	120.94	125.11
2	A	1344	5GP	O4'-C4'-C5'	2.29	116.66	109.33
2	A	1344	5GP	C3'-C2'-C1'	2.27	105.77	101.46
2	A	1344	5GP	C5-C6-N1	2.27	119.03	113.25
2	A	1344	5GP	N9-C8-N7	-2.26	109.22	113.40
2	A	1344	5GP	N2-C2-N1	2.04	121.07	116.76

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	1344	5GP	C4'

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1344	5GP	O4'-C4'-C5'-O5'
2	A	1344	5GP	C3'-C4'-C5'-O5'

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1344	5GP	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	301/336 (89%)	-0.15	7 (2%) 61 63	20, 44, 70, 86	2 (0%)

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	95	LYS	4.4
1	A	96	GLU	3.9
1	A	97	ASP	3.5
1	A	94	VAL	3.2
1	A	99	TYR	2.3
1	A	98	GLU	2.2
1	A	3	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	5GP	A	1344	24/24	0.97	0.09	29,47,49,50	19

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.