



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 09:07 AM UTC

PDB ID : 1AHP / pdb_00001ahp
Title : OLIGOSACCHARIDE SUBSTRATE BINDING IN ESCHERICHIA COLI
MALTODEXTRIN PHSPHORYLASE
Authors : O'Reilly, M.; Watson, K.A.; Schinzel, R.; Palm, D.; Johnson, L.N.
Deposited on : 1997-04-10
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	797	34% 53% 41% 5%
1	B	797	56% 54% 41% 5%
2	C	2	100%
2	D	2	100%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 12830 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called E.COLI MALTODEXTRIN PHOSPHORYLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	796	Total	C	N	O	S	0	0	0
			6366	4064	1126	1156	20			
1	B	796	Total	C	N	O	S	0	0	0
			6366	4064	1126	1156	20			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	112	ALA	ASN	engineered mutation	UNP P00490
A	184	ALA	LYS	conflict	UNP P00490
A	497	ASP	GLN	conflict	UNP P00490
A	547	ARG	HIS	conflict	UNP P00490
A	681	LYS	GLU	conflict	UNP P00490
A	795	ALA	LYS	conflict	UNP P00490
B	112	ALA	ASN	engineered mutation	UNP P00490
B	184	ALA	LYS	conflict	UNP P00490
B	497	ASP	GLN	conflict	UNP P00490
B	547	ARG	HIS	conflict	UNP P00490
B	681	LYS	GLU	conflict	UNP P00490
B	795	ALA	LYS	conflict	UNP P00490

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	C	2	Total	C	O	0	0	0
			23	12	11			
2	D	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is PYRIDOXAL-5'-PHOSPHATE (CCD ID: PLP) (formula: $C_8H_{10}NO_6P$).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
5	B	1	Total	C	N	O	P	0	0
			15	8	1	5	1		

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: E.COLI MALTODEXTRIN PHOSPHORYLASE





● Molecule 1: E.COLI MALTODEXTRIN PHOSPHORYLASE





● Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain C: 100%



● Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain D: 100%



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	173.20Å 112.40Å 121.70Å 90.00° 119.70° 90.00°	Depositor
Resolution (Å)	10.00 – 3.00 10.00 – 3.00	Depositor EDS
% Data completeness (in resolution range)	73.0 (10.00-3.00) 71.5 (10.00-3.00)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.39 (at 2.99Å)	Xtriage
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.232 , 0.278 0.390 , 0.395	Depositor DCC
R_{free} test set	608 reflections (2.09%)	wwPDB-VP
Wilson B-factor (Å ²)	34.3	Xtriage
Anisotropy	0.277	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 52.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.57	EDS
Total number of atoms	12830	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, PLP, SO4, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.49	0/6514	1.00	30/8837 (0.3%)
1	B	0.49	0/6514	1.00	30/8837 (0.3%)
All	All	0.49	0/13028	1.00	60/17674 (0.3%)

There are no bond length outliers.

The worst 5 of 60 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	166	TYR	CA-C-N	8.70	130.71	119.84
1	A	166	TYR	C-N-CA	8.70	130.71	119.84
1	B	166	TYR	CA-C-N	8.69	130.70	119.84
1	B	166	TYR	C-N-CA	8.69	130.70	119.84
1	A	345	HIS	N-CA-C	8.53	123.78	113.12

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6366	0	6326	413	11
1	B	6366	0	6326	407	9
2	C	23	0	20	5	0
2	D	23	0	20	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	5	0	0	1	0
3	B	5	0	0	0	0
4	A	6	0	8	0	0
4	B	6	0	8	0	0
5	A	15	0	7	1	0
5	B	15	0	7	1	0
All	All	12830	0	12722	653	12

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 26.

The worst 5 of 653 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:170:ARG:CG	1:B:20:ARG:O	1.67	1.40
1:A:158:PRO:HG2	1:B:228:PHE:O	1.49	1.11
1:A:253:LYS:HE3	1:B:242:GLU:OE2	1.54	1.07
1:A:20:ARG:HG3	1:B:47:ARG:CZ	1.84	1.07
1:A:19:GLN:CD	1:B:47:ARG:O	2.02	1.01

The worst 5 of 12 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:600:LEU:CD2	1:A:714:LYS:CE[2_555]	0.84	1.36
1:A:600:LEU:CD2	1:A:714:LYS:NZ[2_555]	1.14	1.06
1:A:85:GLN:OE1	1:B:474:LYS:NZ[4_546]	1.22	0.98
1:A:85:GLN:CD	1:B:474:LYS:NZ[4_546]	1.50	0.70
1:A:689:LYS:NZ	1:B:685:ALA:CA[1_545]	1.61	0.59

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	794/797 (100%)	716 (90%)	68 (9%)	10 (1%)	9	38
1	B	794/797 (100%)	717 (90%)	68 (9%)	9 (1%)	11	43
All	All	1588/1594 (100%)	1433 (90%)	136 (9%)	19 (1%)	10	40

5 of 19 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	2	GLN
1	B	2	GLN
1	A	210	ARG
1	A	390	GLU
1	A	391	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	664/665 (100%)	610 (92%)	54 (8%)	11	38
1	B	664/665 (100%)	610 (92%)	54 (8%)	11	38
All	All	1328/1330 (100%)	1220 (92%)	108 (8%)	11	38

5 of 108 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	92	LYS
1	B	302	VAL
1	B	652	LEU
1	B	123	LEU
1	B	204	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 39 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	221	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	531	GLN
1	B	224	HIS
1	B	437	HIS
1	B	589	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	GLC	C	1	2	12,12,12	1.75	4 (33%)	17,17,17	2.40	7 (41%)
2	GLC	C	2	2	11,11,12	2.00	4 (36%)	15,15,17	3.13	9 (60%)
2	GLC	D	1	2	12,12,12	1.75	4 (33%)	17,17,17	2.40	7 (41%)
2	GLC	D	2	2	11,11,12	2.02	4 (36%)	15,15,17	3.13	9 (60%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	C	1	2	-	0/2/22/22	0/1/1/1
2	GLC	C	2	2	-	2/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	D	1	2	-	0/2/22/22	0/1/1/1
2	GLC	D	2	2	-	2/2/19/22	0/1/1/1

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	2	GLC	C4-C5	3.61	1.60	1.53
2	C	2	GLC	C4-C5	3.58	1.60	1.53
2	D	1	GLC	O2-C2	-3.49	1.34	1.43
2	C	1	GLC	O2-C2	-3.48	1.34	1.43
2	C	1	GLC	C6-C5	3.08	1.62	1.51

The worst 5 of 32 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	2	GLC	C6-C5-C4	-6.15	97.91	113.02
2	C	2	GLC	C6-C5-C4	-6.14	97.94	113.02
2	C	2	GLC	C1-C2-C3	5.89	118.22	109.64
2	D	2	GLC	C1-C2-C3	5.85	118.17	109.64
2	C	1	GLC	O5-C1-C2	-5.33	100.92	110.30

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	2	GLC	O5-C5-C6-O6
2	D	2	GLC	O5-C5-C6-O6
2	C	2	GLC	C4-C5-C6-O6
2	D	2	GLC	C4-C5-C6-O6


There are no ring outliers.

4 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1	GLC	1	0
2	D	2	GLC	3	0
2	C	2	GLC	4	0
2	D	1	GLC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	PLP	B	999	1	15,15,16	2.01	3 (20%)	21,22,23	1.55	5 (23%)
4	GOL	A	998	-	5,5,5	0.33	0	5,5,5	0.77	0
4	GOL	B	998	-	5,5,5	0.32	0	5,5,5	0.78	0
3	SO4	B	997	-	4,4,4	0.56	0	6,6,6	0.89	0
5	PLP	A	999	1	15,15,16	2.00	3 (20%)	21,22,23	1.55	5 (23%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	A	997	-	4,4,4	0.56	0	6,6,6	0.89	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	998	-	-	0/4/4/4	-
5	PLP	A	999	1	-	3/6/6/8	0/1/1/1
5	PLP	B	999	1	-	3/6/6/8	0/1/1/1
4	GOL	B	998	-	-	0/4/4/4	-

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	999	PLP	C3-C2	-6.38	1.34	1.41
5	A	999	PLP	C3-C2	-6.36	1.34	1.41
5	A	999	PLP	C5-C4	-2.16	1.38	1.40
5	B	999	PLP	C5-C4	-2.13	1.38	1.40
5	A	999	PLP	P-O2P	2.12	1.62	1.54

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	999	PLP	C4A-C4-C5	-3.30	117.54	120.94
5	A	999	PLP	C4A-C4-C5	-3.27	117.57	120.94
5	B	999	PLP	O3P-P-O1P	2.90	122.14	110.83
5	A	999	PLP	O3P-P-O1P	2.90	122.14	110.83
5	A	999	PLP	C5-C6-N1	-2.64	119.53	123.83

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	999	PLP	C5A-O4P-P-O1P
5	A	999	PLP	C5A-O4P-P-O2P
5	A	999	PLP	C5A-O4P-P-O3P
5	B	999	PLP	C5A-O4P-P-O1P
5	B	999	PLP	C5A-O4P-P-O2P

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	999	PLP	1	0
5	A	999	PLP	1	0
3	A	997	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	796/797 (99%)	1.75	271 (34%) 1 1	2, 20, 61, 115	0
1	B	796/797 (99%)	2.43	447 (56%) 0 0	2, 20, 61, 115	0
All	All	1592/1594 (99%)	2.09	718 (45%) 0 0	2, 20, 61, 115	0

The worst 5 of 718 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	127	ALA	11.6
1	B	780	SER	8.6
1	A	561	ALA	8.2
1	B	760	ASP	7.9
1	B	410	CYS	7.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GLC	D	2	11/12	0.37	0.32	35,35,35,35	0
2	GLC	D	1	12/12	0.59	0.22	35,35,35,35	0
2	GLC	C	2	11/12	0.66	0.22	35,35,35,35	0
2	GLC	C	1	12/12	0.70	0.20	35,35,35,35	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GOL	B	998	6/6	0.47	0.31	91,91,91,91	0
4	GOL	A	998	6/6	0.53	0.27	91,91,91,91	0
3	SO4	B	997	5/5	0.63	0.23	77,77,77,77	0
5	PLP	B	999	15/16	0.68	0.30	25,25,25,25	0
5	PLP	A	999	15/16	0.81	0.21	25,25,25,25	0
3	SO4	A	997	5/5	0.83	0.22	77,77,77,77	0

6.5 Other polymers [i](#)

There are no such residues in this entry.