



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 01:17 PM UTC

PDB ID : 6BMR / pdb_00006bmr
Title : Non-receptor Protein Tyrosine Phosphatase SHP2 in Complex with Allosteric Inhibitor SHP244
Authors : Stams, T.; Fodor, M.
Deposited on : 2017-11-15
Resolution : 2.21 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	526	
1	B	526	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PO4	A	606	-	X	-	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8707 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tyrosine-protein phosphatase non-receptor type 11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	489	Total	C	N	O	S	0	2	0
			3961	2496	702	744	19			
1	B	481	Total	C	N	O	S	0	1	0
			3902	2462	694	727	19			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP Q06124
B	0	SER	-	expression tag	UNP Q06124

- Molecule 2 is 4-[(2-chlorophenyl)methyl]-1-(2-hydroxy-3-methoxyphenyl)[1,2,4]triazolo[4,3-a]quinazolin-5(4H)-one (CCD ID: DZV) (formula: C₂₃H₁₇ClN₄O₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Cl	N	O	0	0
			31	23	1	4	3		
2	B	1	Total	C	Cl	N	O	0	0
			31	23	1	4	3		

- Molecule 3 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		
3	A	1	Total	O	P	0	0
			5	4	1		
3	A	1	Total	O	P	0	0
			5	4	1		
3	A	1	Total	O	P	0	0
			5	4	1		
3	A	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	378	Total	O	0	1
			379	379		
5	B	357	Total	O	0	0
			357	357		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosine-protein phosphatase non-receptor type 11



- Molecule 1: Tyrosine-protein phosphatase non-receptor type 11



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	46.23Å 214.66Å 56.03Å 90.00° 96.96° 90.00°	Depositor
Resolution (Å)	107.00 – 2.21 107.00 – 2.21	Depositor EDS
% Data completeness (in resolution range)	99.0 (107.00-2.21) 99.7 (107.00-2.21)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.26 (at 2.20Å)	Xtriage
Refinement program	BUSTER 2.11.7	Depositor
R, R_{free}	0.167 , 0.224 0.175 , 0.233	Depositor DCC
R_{free} test set	2760 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å ²)	29.3	Xtriage
Anisotropy	0.602	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 58.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8707	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.37% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, DZV, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.87	0/4047	1.28	10/5454 (0.2%)
1	B	0.87	0/3984	1.26	6/5368 (0.1%)
All	All	0.87	0/8031	1.27	16/10822 (0.1%)

There are no bond length outliers.

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	140	SER	CA-C-N	7.21	134.68	121.70
1	A	140	SER	C-N-CA	7.21	134.68	121.70
1	B	106	ASP	CA-CB-CG	6.30	118.90	112.60
1	B	64	ASP	CA-CB-CG	5.92	118.52	112.60
1	A	296	ASP	CA-CB-CG	5.78	118.38	112.60
1	A	106	ASP	CA-CB-CG	5.72	118.32	112.60
1	A	198	LYS	N-CA-C	-5.69	104.69	111.69
1	A	358	LYS	N-CA-C	-5.35	103.33	110.55
1	B	105	ALA	CA-C-N	5.24	128.33	120.83
1	B	105	ALA	C-N-CA	5.24	128.33	120.83
1	A	273	ASN	CA-C-N	5.11	127.44	120.54
1	A	273	ASN	C-N-CA	5.11	127.44	120.54
1	A	437	ASP	CA-C-N	5.05	127.01	120.44
1	A	437	ASP	C-N-CA	5.05	127.01	120.44
1	B	437	ASP	CA-C-N	5.00	126.94	120.44
1	B	437	ASP	C-N-CA	5.00	126.94	120.44

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3961	0	3902	21	0
1	B	3902	0	3855	19	0
2	A	31	0	0	0	0
2	B	31	0	0	0	0
3	A	25	0	0	0	0
3	B	15	0	0	1	0
4	B	6	0	8	1	0
5	A	379	0	0	0	0
5	B	357	0	0	1	0
All	All	8707	0	7765	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:140:SER:HA	1:A:141:GLN:HB3	1.58	0.85
1:A:272:GLU:CB	1:A:301:VAL:HG11	2.21	0.70
1:B:357:THR:HG22	5:B:713:HOH:O	1.93	0.68
1:A:294:ASP:HB2	1:A:343:ARG:HE	1.59	0.68
1:B:272:GLU:HG3	1:B:301:VAL:HG21	1.77	0.65
1:A:140:SER:CA	1:A:141:GLN:HB3	2.36	0.54
1:A:271:GLN:HA	1:A:274:LYS:HD2	1.90	0.54
1:A:294:ASP:HB2	1:A:343:ARG:NE	2.21	0.53
1:B:394:HIS:O	3:B:604:PO4:O1	2.27	0.52
1:A:272:GLU:HB3	1:A:301:VAL:HG11	1.92	0.51
1:B:125:LEU:HB3	1:B:216:LEU:HD21	1.92	0.51
1:A:224:ALA:HB2	1:A:484:VAL:HG13	1.93	0.51
1:A:272:GLU:CG	1:A:301:VAL:HG11	2.41	0.50
1:A:133:GLY:HA3	1:A:213:LYS:HB2	1.96	0.48
1:A:382:VAL:HG23	1:A:383:MET:HE2	1.95	0.47
1:B:391:SER:HB3	4:B:605:GOL:H11	1.96	0.47
1:B:28:SER:HA	1:B:100:TYR:O	2.15	0.47
1:A:290:VAL:HG11	1:A:344:MET:HG3	1.96	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:309:ILE:HD13	1:A:328:ILE:HG12	1.96	0.46
1:B:522:ILE:O	1:B:525:LEU:HB2	2.16	0.46
1:B:432:PRO:HG3	1:B:516:MET:HG2	1.97	0.46
1:A:90:GLU:HG2	1:A:96:ILE:HD11	1.97	0.46
1:B:149:LEU:HB2	1:B:172:ILE:HD11	1.98	0.45
1:B:222:ASN:O	1:B:225:GLU:HG2	2.16	0.45
1:B:126:LEU:HB3	1:B:152:ARG:HG3	1.99	0.45
1:B:126:LEU:HD22	1:B:152:ARG:HB2	1.98	0.44
1:A:20:LEU:HD23	1:A:102:LEU:HD13	1.99	0.44
1:A:390:GLU:HG2	1:A:399:ARG:HG2	1.99	0.43
1:B:81:TYR:CD1	1:B:88:LEU:HB2	2.53	0.43
1:A:140:SER:HA	1:A:141:GLN:CB	2.38	0.43
1:B:88:LEU:HG	1:B:96:ILE:HG23	1.99	0.43
1:B:149:LEU:HD21	1:B:212:LEU:HD21	2.01	0.43
1:B:329:ALA:HB3	1:B:501:ARG:HD3	2.01	0.42
1:B:47:ARG:HD3	1:B:96:ILE:HD12	2.02	0.42
1:B:203:VAL:HG22	1:B:209:VAL:HG22	2.02	0.41
1:B:356:THR:OG1	1:B:459:CYS:HB3	2.21	0.41
1:A:194:VAL:O	1:A:198:LYS:HB2	2.20	0.41
1:A:418:TYR:HB3	1:A:438:PHE:CE1	2.56	0.41
1:A:366:LYS:HA	1:A:366:LYS:HD3	1.91	0.40
1:A:385:VAL:HA	1:A:402:LYS:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	477/526 (91%)	465 (98%)	10 (2%)	2 (0%)	30	34
1	B	468/526 (89%)	454 (97%)	13 (3%)	1 (0%)	43	51
All	All	945/1052 (90%)	919 (97%)	23 (2%)	3 (0%)	36	42

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	295	GLY
1	A	505	VAL
1	B	505	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	436/468 (93%)	408 (94%)	28 (6%)	16	19
1	B	429/468 (92%)	408 (95%)	21 (5%)	22	29
All	All	865/936 (92%)	816 (94%)	49 (6%)	18	23

All (49) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	57	GLN
1	A	87	GLN
1	A	94	ASP
1	A	96	ILE
1	A	102	LEU
1	A	111	ARG
1	A	118	SER
1	A	131	LYS
1	A	140	SER
1	A	151	VAL
1	A	152	ARG
1	A	165	SER
1	A	198	LYS
1	A	221	ILE
1	A	228	SER
1	A	236	LEU
1	A	245	GLN
1	A	249	GLU
1	A	325	LYS
1	A	335	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	378	LYS
1	A	402	LYS
1	A	408	GLN
1	A	439	LEU
1	A	477	ASP
1	A	481	GLU
1	A	484	VAL
1	A	508	GLU
1	B	47	ARG
1	B	55	LYS
1	B	90	GLU
1	B	96	ILE
1	B	129	LYS
1	B	131	LYS
1	B	152	ARG
1	B	166	LYS
1	B	171	MET
1	B	199	LYS
1	B	221	ILE
1	B	272	GLU
1	B	275	ASN
1	B	325	LYS
1	B	368	VAL
1	B	405	LYS
1	B	421	ARG
1	B	426	HIS
1	B	481	GLU
1	B	484	VAL
1	B	508	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	114	HIS
1	A	141	GLN
1	A	196	HIS
1	A	222	ASN
1	A	275	ASN
1	B	116	HIS
1	B	200	ASN
1	B	211	GLN
1	B	446	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	DZV	B	601	-	35,35,35	1.74	7 (20%)	47,51,51	2.05	7 (14%)
4	GOL	B	605	-	5,5,5	0.10	0	5,5,5	0.24	0
2	DZV	A	601	-	35,35,35	1.71	11 (31%)	47,51,51	2.04	8 (17%)
3	PO4	A	604	-	4,4,4	2.56	2 (50%)	6,6,6	0.44	0
3	PO4	B	602	-	4,4,4	2.51	2 (50%)	6,6,6	0.73	0
3	PO4	A	603	-	4,4,4	2.57	2 (50%)	6,6,6	0.42	0
3	PO4	A	605	-	4,4,4	2.12	3 (75%)	6,6,6	0.34	0
3	PO4	B	604	-	4,4,4	2.48	3 (75%)	6,6,6	1.06	0
3	PO4	A	606	-	4,4,4	2.08	4 (100%)	6,6,6	0.41	0
3	PO4	A	602	-	4,4,4	2.54	2 (50%)	6,6,6	0.70	0
3	PO4	B	603	-	4,4,4	1.96	1 (25%)	6,6,6	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DZV	A	601	-	-	1/10/10/10	0/5/5/5
2	DZV	B	601	-	-	2/10/10/10	0/5/5/5
4	GOL	B	605	-	-	2/4/4/4	-

All (37) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	603	PO4	P-O1	4.28	1.60	1.50
3	A	604	PO4	P-O1	4.27	1.60	1.50
3	A	602	PO4	P-O1	4.23	1.60	1.50
3	B	602	PO4	P-O1	4.13	1.60	1.50
3	B	604	PO4	P-O1	4.06	1.60	1.50
2	B	601	DZV	C1-N2	3.08	1.44	1.38
2	B	601	DZV	C10-C9	2.95	1.53	1.47
2	A	601	DZV	C1-N2	2.87	1.43	1.38
2	B	601	DZV	C4-C3	2.86	1.43	1.39
2	A	601	DZV	C19-C18	2.71	1.44	1.40
2	B	601	DZV	C1-N1	-2.69	1.33	1.39
2	A	601	DZV	C1-N1	-2.66	1.33	1.39
2	A	601	DZV	C8-C3	2.56	1.43	1.39
2	B	601	DZV	C12-C11	2.49	1.43	1.39
3	A	605	PO4	P-O1	2.45	1.56	1.50
2	B	601	DZV	C8-C3	2.41	1.43	1.39
2	A	601	DZV	C7-C8	2.38	1.43	1.38
3	A	606	PO4	P-O1	2.26	1.55	1.50
2	A	601	DZV	C13-C12	2.25	1.42	1.38
2	A	601	DZV	C10-C9	2.22	1.52	1.47
2	A	601	DZV	C23-C17	2.16	1.43	1.39
2	A	601	DZV	C12-C11	2.15	1.43	1.39
2	A	601	DZV	C4-C3	2.11	1.42	1.39
2	B	601	DZV	C19-C18	2.10	1.43	1.40
2	A	601	DZV	C5-C4	2.06	1.43	1.38
3	A	606	PO4	P-O2	2.03	1.60	1.54
3	B	604	PO4	P-O4	2.02	1.60	1.54
3	A	602	PO4	P-O4	2.02	1.60	1.54
3	B	604	PO4	P-O3	2.02	1.60	1.54
3	A	605	PO4	P-O4	2.02	1.60	1.54
3	A	606	PO4	P-O4	2.01	1.60	1.54
3	A	606	PO4	P-O3	2.01	1.60	1.54
3	A	603	PO4	P-O2	2.01	1.60	1.54
3	B	602	PO4	P-O2	2.01	1.60	1.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	604	PO4	P-O3	2.01	1.60	1.54
3	A	605	PO4	P-O2	2.01	1.60	1.54
3	B	603	PO4	P-O4	2.00	1.60	1.54

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601	DZV	O2-C19-C18	9.49	124.46	114.53
2	A	601	DZV	O2-C19-C18	8.96	123.90	114.53
2	B	601	DZV	C20-O2-C19	5.30	125.29	117.51
2	A	601	DZV	C3-C2-N2	4.15	118.15	113.14
2	A	601	DZV	C20-O2-C19	4.11	123.54	117.51
2	A	601	DZV	O2-C19-C21	-4.06	117.45	124.30
2	B	601	DZV	C3-C2-N2	3.99	117.97	113.14
2	B	601	DZV	O2-C19-C21	-3.64	118.17	124.30
2	A	601	DZV	C2-C3-C4	-3.37	116.13	121.67
2	B	601	DZV	C21-C19-C18	-2.97	117.23	119.98
2	A	601	DZV	O1-C9-N2	2.55	124.31	120.15
2	A	601	DZV	C17-C16-N4	-2.37	119.86	124.62
2	A	601	DZV	C2-C3-C8	2.26	124.61	120.12
2	B	601	DZV	C2-C3-C4	-2.12	118.19	121.67
2	B	601	DZV	C23-C17-C18	2.07	120.77	118.76

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	605	GOL	O2-C2-C3-O3
4	B	605	GOL	C1-C2-C3-O3
2	B	601	DZV	C21-C19-O2-C20
2	A	601	DZV	N4-C16-C17-C23
2	B	601	DZV	N4-C16-C17-C23


There are no ring outliers.


2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	605	GOL	1	0
3	B	604	PO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	489/526 (92%)	-0.19	7 (1%) 73 71	16, 38, 64, 89	2 (0%)
1	B	481/526 (91%)	-0.14	5 (1%) 79 77	20, 38, 65, 94	1 (0%)
All	All	970/1052 (92%)	-0.17	12 (1%) 76 74	16, 38, 65, 94	3 (0%)

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	236	LEU	3.6
1	B	144	PRO	3.3
1	A	154	GLY	3.2
1	B	426	HIS	3.0
1	A	177	LEU	2.6
1	A	469	PHE	2.6
1	A	144	PRO	2.5
1	A	141	GLN	2.4
1	B	153	THR	2.4
1	B	301	VAL	2.1
1	B	293	HIS	2.1
1	A	300	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	PO4	A	603	5/5	0.76	0.13	112,113,113,113	0
3	PO4	B	604	5/5	0.80	0.15	112,113,113,114	0
3	PO4	A	606	5/5	0.81	0.14	117,118,118,119	0
3	PO4	A	605	5/5	0.84	0.11	107,108,108,109	0
4	GOL	B	605	6/6	0.87	0.15	62,62,64,67	0
3	PO4	A	604	5/5	0.91	0.10	90,91,91,92	0
3	PO4	B	603	5/5	0.91	0.10	101,101,102,104	0
3	PO4	A	602	5/5	0.95	0.07	55,56,58,58	0
2	DZV	B	601	31/31	0.95	0.06	23,29,35,42	0
3	PO4	B	602	5/5	0.96	0.07	40,41,45,46	0
2	DZV	A	601	31/31	0.97	0.06	25,31,34,34	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around DZV B 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around DZV A 601:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.