



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:52 PM UTC

PDB ID : 5BVN / pdb_00005bvn
Title : Fragment-based discovery of potent and selective DDR1/2 inhibitors
Authors : Murray, C.; Berdini, V.; Buck, I.; Carr, M.; Cleasby, A.; Coyle, J.; Curry, J.; Day, J.; Hearn, K.; Iqbal, A.; Lee, L.; Martins, V.; Mortenson, P.; Munck, J.; Page, L.; Patel, S.; Roomans, S.; Kirsten, T.; Saxty, G.
Deposited on : 2015-06-05
Resolution : 2.21 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7682 (2.24-2.20)
Clashscore	190562	8402 (2.24-2.20)
Ramachandran outliers	187476	8303 (2.24-2.20)
Sidechain outliers	187428	8304 (2.24-2.20)
RSRZ outliers	180081	7683 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	324	14% 62% 20% •• 14%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2468 atoms, of which 20 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Epithelial discoidin domain-containing receptor 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	278	2245	1430	396	401	18	0	0	0


There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	590	GLY	-	expression tag	UNP Q08345
A	591	ALA	-	expression tag	UNP Q08345
A	592	MET	-	expression tag	UNP Q08345
A	593	GLY	-	expression tag	UNP Q08345
A	594	SER	-	expression tag	UNP Q08345

- Molecule 2 is IODIDE ION (CCD ID: IOD) (formula: I).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	I	0	0
			1	1		

- Molecule 3 is N-[5-({[(3-fluorophenyl)carbamoyl]amino}methyl)-2-methylphenyl]imidazo[1,2-a]pyridine-3-carboxamide (CCD ID: 4VD) (formula: C₂₃H₂₀FN₅O₂).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	F	H	N	O	0	0
			51	23	1	20	5	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	171	Total O 171 171	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Epithelial discoidin domain-containing receptor 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	61.04Å 78.50Å 74.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	31.80 – 2.21 31.80 – 2.21	Depositor EDS
% Data completeness (in resolution range)	96.0 (31.80-2.21) 95.9 (31.80-2.21)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.56 (at 2.22Å)	Xtriage
Refinement program	BUSTER BUSTER 2.11.6	Depositor
R, R_{free}	0.252 , 0.328 0.258 , 0.326	Depositor DCC
R_{free} test set	918 reflections (5.17%)	wwPDB-VP
Wilson B-factor (Å ²)	18.8	Xtriage
Anisotropy	0.580	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 63.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.027 for -h,l,k	Xtriage
F_o, F_c correlation	0.84	EDS
Total number of atoms	2468	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: IOD, 4VD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.99	0/2295	1.42	23/3101 (0.7%)

There are no bond length outliers.

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	868	VAL	CB-CA-C	6.93	120.94	110.62
1	A	708	ASP	CA-CB-CG	6.41	119.01	112.60
1	A	884	MET	CA-C-N	6.38	128.73	120.44
1	A	884	MET	C-N-CA	6.38	128.73	120.44
1	A	611	ARG	CA-C-N	6.25	130.27	121.02
1	A	611	ARG	C-N-CA	6.25	130.27	121.02
1	A	859	GLU	CB-CG-CD	6.22	123.17	112.60
1	A	757	LEU	CA-C-N	5.75	128.25	120.38
1	A	757	LEU	C-N-CA	5.75	128.25	120.38
1	A	891	GLU	CA-C-N	5.70	127.91	120.28
1	A	891	GLU	C-N-CA	5.70	127.91	120.28
1	A	904	PHE	CA-CB-CG	-5.51	108.29	113.80
1	A	667	ASN	CA-C-N	5.47	127.93	120.54
1	A	667	ASN	C-N-CA	5.47	127.93	120.54
1	A	776	GLU	CA-C-N	5.29	131.23	121.70
1	A	776	GLU	C-N-CA	5.29	131.23	121.70
1	A	835	GLU	CA-C-N	5.21	127.13	120.56
1	A	835	GLU	C-N-CA	5.21	127.13	120.56
1	A	744	LEU	CA-C-N	5.06	127.31	120.38
1	A	744	LEU	C-N-CA	5.06	127.31	120.38
1	A	881	TYR	CA-C-N	5.03	127.27	120.38
1	A	881	TYR	C-N-CA	5.03	127.27	120.38
1	A	790	ASN	CA-CB-CG	5.01	117.61	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2245	0	2226	34	0
2	A	1	0	0	1	0
3	A	31	20	20	0	0
4	A	171	0	0	0	0
All	All	2448	20	2246	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:859:GLU:HG3	1:A:866:ARG:HB2	1.74	0.70
1:A:776:GLU:HA	1:A:778:PHE:N	2.18	0.59
1:A:776:GLU:HA	1:A:778:PHE:H	1.70	0.56
1:A:837:LEU:HD11	1:A:880:LEU:HD21	1.86	0.56
1:A:693:GLN:H	1:A:693:GLN:CD	2.14	0.55
1:A:883:LEU:HD12	1:A:886:ARG:HH21	1.72	0.54
1:A:852:GLN:HA	1:A:855:GLU:HG2	1.90	0.52
1:A:706:ASN:HB2	1:A:774:VAL:HB	1.93	0.49
1:A:805:LEU:HB3	1:A:807:ILE:HG23	1.96	0.48
1:A:811:ALA:HA	1:A:827:TRP:CE2	2.49	0.48
1:A:833:LEU:HG	1:A:884:MET:HE2	1.97	0.47
1:A:814:CYS:SG	1:A:824:SER:HB2	2.54	0.47
1:A:621:PHE:CZ	1:A:668:ASP:HB3	2.51	0.46
1:A:692:VAL:H	1:A:693:GLN:NE2	2.14	0.46
1:A:679:LEU:HB3	1:A:684:ILE:HG21	1.98	0.46
1:A:884:MET:O	1:A:887:CYS:HB2	2.16	0.45
1:A:860:PHE:HA	1:A:867:GLN:HG2	2.00	0.44
1:A:811:ALA:HA	1:A:827:TRP:CD2	2.52	0.43
1:A:796:TYR:HB3	1:A:804:VAL:HG12	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:606:PRO:HB2	1:A:609:ARG:HG3	2.01	0.42
1:A:606:PRO:HG2	1:A:609:ARG:HH11	1.84	0.42
1:A:816:LEU:HB2	1:A:817:MET:HE2	2.00	0.42
1:A:620:GLN:HB2	1:A:655:LYS:NZ	2.34	0.42
1:A:626:LEU:HD11	1:A:651:LEU:HB3	2.01	0.42
1:A:845:PHE:HA	2:A:1001:IOD:I	2.90	0.42
1:A:687:LEU:HD11	1:A:699:MET:HG2	2.02	0.41
1:A:876:CYS:SG	1:A:884:MET:HE1	2.60	0.41
1:A:675:ILE:O	1:A:678:ARG:HB2	2.20	0.41
1:A:755:ARG:O	1:A:759:THR:HG23	2.20	0.41
1:A:827:TRP:CE3	1:A:888:TRP:HA	2.56	0.41
1:A:898:PHE:HA	1:A:901:LEU:HD12	2.03	0.41
1:A:622:GLY:HA2	1:A:658:ARG:HB2	2.03	0.40
1:A:693:GLN:H	1:A:693:GLN:NE2	2.20	0.40
1:A:796:TYR:HB3	1:A:804:VAL:CG1	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	272/324 (84%)	255 (94%)	15 (6%)	2 (1%)	18	18

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	776	GLU
1	A	618	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	242/272 (89%)	220 (91%)	22 (9%)	9 8

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	610	LEU
1	A	657	LEU
1	A	663	LYS
1	A	679	LEU
1	A	686	ARG
1	A	690	VAL
1	A	693	GLN
1	A	698	CYS
1	A	702	ASP
1	A	774	VAL
1	A	776	GLU
1	A	787	MET
1	A	790	ASN
1	A	800	GLN
1	A	805	LEU
1	A	837	LEU
1	A	859	GLU
1	A	868	VAL
1	A	884	MET
1	A	893	GLU
1	A	894	GLN
1	A	903	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	683	ASN
1	A	911	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	4VD	A	1002	-	34,34,34	0.30	0	46,47,47	0.71	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	4VD	A	1002	-	-	0/17/17/17	0/4/4/4

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	278/324 (85%)	1.16	45 (16%) 4 3	10, 26, 51, 77	0

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	617	GLY	8.3
1	A	631	SER	5.2
1	A	605	PHE	4.8
1	A	721	LYS	4.7
1	A	695	ASP	4.4
1	A	618	GLU	4.3
1	A	698	CYS	4.3
1	A	648	HIS	3.7
1	A	911	ASN	3.6
1	A	864	GLN	3.5
1	A	650	LEU	3.4
1	A	694	ASP	3.2
1	A	622	GLY	3.2
1	A	692	VAL	3.1
1	A	693	GLN	3.1
1	A	649	PRO	3.0
1	A	910	LEU	3.0
1	A	740	TYR	3.0
1	A	604	ASP	3.0
1	A	607	ARG	2.9
1	A	611	ARG	2.8
1	A	619	GLY	2.8
1	A	697	LEU	2.7
1	A	800	GLN	2.7
1	A	823	ALA	2.6
1	A	809	TRP	2.6
1	A	630	ASP	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	606	PRO	2.5
1	A	656	ILE	2.5
1	A	609	ARG	2.5
1	A	720	ASP	2.5
1	A	790	ASN	2.5
1	A	777	ASN	2.5
1	A	718	LEU	2.4
1	A	620	GLN	2.4
1	A	776	GLU	2.4
1	A	753	GLY	2.4
1	A	719	GLU	2.4
1	A	778	PHE	2.2
1	A	818	GLY	2.2
1	A	608	SER	2.2
1	A	714	SER	2.2
1	A	773	LEU	2.2
1	A	629	VAL	2.1
1	A	612	PHE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	4VD	A	1002	31/31	0.91	0.11	10,25,31,32	0
2	IOD	A	1001	1/1	0.92	0.13	89,89,89,89	0

6.5 Other polymers [i](#)

There are no such residues in this entry.