



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 02:48 AM UTC

PDB ID : 3BZM / pdb_00003bzm
Title : Crystal Structure of Open form of Menaquinone-Specific Isochorismate Synthase, MenF
Authors : Parsons, J.F.; Shi, K.M.; Ladner, J.E.
Deposited on : 2008-01-18
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	431	9% 67% 24% 7%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3546 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Menaquinone-specific isochorismate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	429	Total	C	N	O	S	0	3	0
			3432	2163	625	631	13			

- Molecule 2 is CITRIC ACID (CCD ID: CIT) (formula: $C_6H_8O_7$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			13	6	7		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	101	Total	O	0	0
			101	101		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Menaquinone-specific isochorismate synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	72.37Å 72.37Å 164.01Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.67 – 1.95 19.67 – 1.95	Depositor EDS
% Data completeness (in resolution range)	96.4 (19.67-1.95) 96.3 (19.67-1.95)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.58 (at 1.94Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.245 , 0.308 0.254 , 0.313	Depositor DCC
R_{free} test set	1840 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	45.3	Xtriage
Anisotropy	0.257	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 45.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.53$, $\langle L^2 \rangle = 0.37$	Xtriage
Estimated twinning fraction	0.010 for -h,-k,l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3546	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CIT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.19	5/3515 (0.1%)	1.33	22/4773 (0.5%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	28	ILE	CA-CB	6.33	1.61	1.54
1	A	108	LEU	CA-C	-5.30	1.46	1.52
1	A	93	ARG	N-CA	5.28	1.52	1.46
1	A	144	ILE	CA-C	5.28	1.59	1.52
1	A	201	PHE	CA-C	-5.17	1.46	1.52

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	185	GLU	N-CA-C	25.33	138.17	111.07
1	A	185	GLU	CB-CA-C	-10.56	94.31	110.88
1	A	189	ASP	N-CA-CB	-10.28	93.12	110.49
1	A	350	ARG	CB-CA-C	-9.87	94.35	110.74
1	A	350	ARG	N-CA-C	9.39	126.87	111.37
1	A	364	THR	CB-CA-C	-8.63	96.08	109.89
1	A	188	LEU	N-CA-C	-7.75	97.99	110.32
1	A	153	LEU	CA-C-N	-7.50	113.01	120.21
1	A	153	LEU	C-N-CA	-7.50	113.01	120.21
1	A	188	LEU	CB-CA-C	-7.11	97.36	109.51
1	A	189	ASP	N-CA-C	-6.61	96.72	110.80
1	A	183	ILE	CB-CA-C	-6.29	103.92	111.97
1	A	279	ASP	N-CA-C	-6.17	101.28	110.23
1	A	384	VAL	N-CA-C	-6.06	99.66	108.87
1	A	345	VAL	CB-CA-C	5.64	116.18	110.65
1	A	427	LEU	N-CA-C	-5.46	106.62	113.28

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	302	THR	N-CA-C	5.30	118.16	110.42
1	A	405	VAL	N-CA-C	5.28	120.33	109.34
1	A	96	GLY	N-CA-C	5.21	116.96	110.29
1	A	215	ARG	N-CA-C	5.14	117.63	111.71
1	A	339	LEU	N-CA-C	5.13	119.59	113.23
1	A	209	ALA	N-CA-C	-5.02	105.52	111.69

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3432	0	3420	96	0
2	A	13	0	5	1	0
3	A	101	0	0	6	0
All	All	3546	0	3425	96	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (96) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:405:VAL:O	1:A:406:ARG:HB3	1.54	1.07
1:A:408:SER:O	1:A:409:ASP:HB3	1.60	1.00
1:A:406:ARG:HG3	1:A:407:GLY:N	1.78	0.98
1:A:172:TRP:O	1:A:176:ILE:HG12	1.70	0.92
1:A:255:LEU:HD23	1:A:320:ARG:HB2	1.58	0.85
1:A:179:ALA:O	1:A:183:ILE:HG12	1.78	0.83
1:A:350:ARG:O	1:A:354:ARG:HG3	1.81	0.81
1:A:276:MET:O	1:A:282:GLN:NE2	2.14	0.80
1:A:384:VAL:HG12	1:A:386:LEU:HD13	1.66	0.78
1:A:53:PHE:CD1	1:A:226[B]:MET:HE3	2.18	0.78
1:A:406:ARG:HG3	1:A:407:GLY:H	1.47	0.78

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:408:SER:O	1:A:409:ASP:CB	2.30	0.77
1:A:52:GLN:HG2	1:A:67:LEU:CD1	2.17	0.75
1:A:80:GLN:NE2	1:A:333:VAL:HG13	2.05	0.72
1:A:163:GLU:HG3	1:A:199:LEU:HD12	1.74	0.70
1:A:285:ASN:OD1	1:A:321:ARG:HD2	1.93	0.69
1:A:261:ASN:OD1	1:A:312:ARG:HD2	1.94	0.68
1:A:126:THR:HG23	3:A:439:HOH:O	1.92	0.67
1:A:180:THR:HA	1:A:183:ILE:HG12	1.75	0.67
1:A:53:PHE:HD1	1:A:226[B]:MET:HE3	1.64	0.62
1:A:164:GLN:HG2	1:A:166:TRP:CZ2	2.35	0.61
1:A:162:ARG:HG2	1:A:163:GLU:N	2.15	0.61
1:A:279:ASP:O	1:A:280:LYS:HB2	1.99	0.61
1:A:193:LEU:HD12	1:A:404:ILE:HG13	1.82	0.60
1:A:364:THR:HG22	1:A:366:GLU:H	1.67	0.60
1:A:84:ARG:O	1:A:87:PRO:HD3	2.02	0.59
1:A:183:ILE:HD13	1:A:188:LEU:HD23	1.82	0.59
1:A:217:LEU:HD13	1:A:314:ARG:NH2	2.17	0.59
1:A:173:THR:HG23	1:A:410:PRO:HG3	1.86	0.58
1:A:409:ASP:O	1:A:410:PRO:C	2.47	0.57
1:A:162:ARG:NH2	1:A:164:GLN:OE1	2.34	0.56
1:A:291:ASP:OD2	1:A:295:ARG:NH1	2.36	0.56
1:A:276:MET:C	1:A:282:GLN:HE21	2.13	0.56
1:A:240:GLU:HB2	1:A:255:LEU:HD12	1.88	0.56
1:A:148:VAL:HG22	1:A:149:SER:H	1.70	0.56
1:A:405:VAL:O	1:A:406:ARG:CB	2.39	0.55
1:A:244:ARG:HD2	3:A:468:HOH:O	2.08	0.54
1:A:222:TYR:O	1:A:237:SER:HA	2.08	0.54
1:A:80:GLN:NE2	1:A:333:VAL:CG1	2.72	0.53
1:A:65:VAL:HG23	1:A:65:VAL:O	2.10	0.52
1:A:279:ASP:O	1:A:280:LYS:CB	2.57	0.52
1:A:180:THR:HA	1:A:183:ILE:CG1	2.38	0.51
1:A:36:ASP:OD1	1:A:36:ASP:N	2.44	0.51
1:A:242:LEU:HD13	1:A:243:TRP:CE3	2.45	0.51
1:A:162:ARG:HG2	1:A:163:GLU:H	1.76	0.51
1:A:188:LEU:HD21	1:A:191:VAL:HB	1.92	0.51
1:A:44:LEU:HG	1:A:65:VAL:HB	1.93	0.51
1:A:80:GLN:HE22	1:A:333:VAL:HG13	1.77	0.50
1:A:259:VAL:HG22	1:A:275:LEU:HD22	1.94	0.50
1:A:371:SER:HB2	1:A:382:PHE:HB3	1.94	0.50
1:A:293:CYS:HB3	1:A:303:LEU:HD13	1.94	0.49
1:A:59:ASN:ND2	2:A:432:CIT:H41	2.28	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:262:ASN:O	1:A:268:ALA:HB2	2.12	0.48
1:A:163:GLU:HG3	1:A:199:LEU:CD1	2.41	0.48
1:A:54:TYR:HD1	1:A:65:VAL:CG1	2.26	0.48
1:A:384:VAL:CG1	1:A:386:LEU:HD13	2.41	0.48
1:A:52:GLN:HG2	1:A:67:LEU:HD12	1.94	0.47
1:A:261:ASN:OD1	1:A:312:ARG:CD	2.61	0.47
1:A:351:ASP:O	1:A:355:GLN:HG3	2.14	0.47
1:A:406:ARG:CG	1:A:407:GLY:H	2.24	0.47
1:A:143:PHE:O	1:A:146:THR:HB	2.15	0.47
1:A:148:VAL:HG22	1:A:149:SER:N	2.30	0.47
1:A:106:ASN:HB2	1:A:367:TRP:CH2	2.50	0.47
1:A:278:ASP:OD2	1:A:278:ASP:C	2.58	0.46
1:A:286:MET:HA	1:A:289:VAL:HG13	1.95	0.46
1:A:306:LEU:HG	1:A:322:CYS:HB2	1.97	0.46
1:A:41:LEU:HD12	1:A:215:ARG:CZ	2.45	0.46
1:A:350:ARG:O	1:A:354:ARG:CG	2.59	0.46
1:A:409:ASP:HA	1:A:410:PRO:HD2	1.65	0.46
1:A:54:TYR:HD1	1:A:65:VAL:HG12	1.82	0.45
1:A:244:ARG:CD	3:A:468:HOH:O	2.65	0.45
1:A:112:LEU:HD13	1:A:125:LEU:HD12	1.98	0.45
1:A:366:GLU:HG2	1:A:367:TRP:CD1	2.51	0.45
1:A:242:LEU:HD13	1:A:243:TRP:CZ3	2.53	0.44
1:A:406:ARG:CG	1:A:407:GLY:N	2.63	0.44
1:A:301:GLN:O	1:A:302:THR:C	2.60	0.44
1:A:296:LEU:HD11	1:A:339:LEU:HD21	1.98	0.44
1:A:95:TRP:CE3	1:A:374:TYR:HB3	2.52	0.44
1:A:350:ARG:HG2	3:A:505:HOH:O	2.18	0.43
1:A:366:GLU:OE2	1:A:390:LYS:NZ	2.47	0.43
1:A:57:GLN:OE1	1:A:59:ASN:HB2	2.18	0.43
1:A:265:ASP:O	1:A:269:GLN:HB3	2.19	0.42
1:A:40:ALA:HB1	1:A:65:VAL:HG21	2.01	0.42
1:A:345:VAL:HG21	1:A:369:ALA:HB1	2.01	0.42
1:A:1:MET:HE3	1:A:1:MET:HB2	1.95	0.42
1:A:40:ALA:HB1	1:A:65:VAL:CG2	2.50	0.42
1:A:61:ASP:O	1:A:118:GLY:N	2.53	0.42
1:A:195:ARG:CB	1:A:417:ILE:HD12	2.50	0.42
1:A:216:ARG:NE	3:A:494:HOH:O	2.23	0.42
1:A:366:GLU:HG2	1:A:367:TRP:H	1.84	0.41
1:A:341:PRO:CG	1:A:354:ARG:HD3	2.49	0.41
1:A:165:HIS:ND1	1:A:195:ARG:HD2	2.36	0.41
1:A:423:GLY:HA3	3:A:532:HOH:O	2.20	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:172:TRP:CE2	1:A:176:ILE:HD11	2.55	0.40
1:A:195:ARG:HB3	1:A:417:ILE:HD12	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	430/431 (100%)	403 (94%)	22 (5%)	5 (1%)	10 4

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	280	LYS
1	A	406	ARG
1	A	408	SER
1	A	410	PRO
1	A	409	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	364/363 (100%)	317 (87%)	47 (13%)	4 1

All (47) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	GLU
1	A	25	ILE
1	A	36	ASP
1	A	65	VAL
1	A	66	VAL
1	A	71	THR
1	A	78	GLN
1	A	108	LEU
1	A	109	LEU
1	A	125	LEU
1	A	126	THR
1	A	146	THR
1	A	153	LEU
1	A	156	LEU
1	A	158	LEU
1	A	161	THR
1	A	180	THR
1	A	181	LYS
1	A	199	LEU
1	A	242	LEU
1	A	244	ARG
1	A	250	LEU
1	A	255	LEU
1	A	267	GLN
1	A	269	GLN
1	A	276	MET
1	A	278	ASP
1	A	279	ASP
1	A	287	LEU
1	A	289	VAL
1	A	306	LEU
1	A	309	GLN
1	A	316	VAL
1	A	326	SER
1	A	333	VAL
1	A	336	LEU
1	A	345	VAL
1	A	348	LEU
1	A	350	ARG
1	A	364	THR
1	A	377	LEU
1	A	386	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	390	LYS
1	A	405	VAL
1	A	406	ARG
1	A	424	LEU
1	A	427	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	10	ASN
1	A	56	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	CIT	A	432	-	12,12,12	1.27	1 (8%)	17,17,17	2.00	5 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CIT	A	432	-	-	10/16/16/16	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	432	CIT	O4-C5	-2.36	1.23	1.30

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	432	CIT	O6-C6-C3	3.76	120.36	113.14
2	A	432	CIT	O4-C5-O3	-3.40	114.58	123.33
2	A	432	CIT	C3-C2-C1	-3.23	105.09	113.92
2	A	432	CIT	C2-C3-C6	2.47	115.50	110.03
2	A	432	CIT	O4-C5-C4	2.23	121.42	114.35

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	432	CIT	C2-C3-C4-C5
2	A	432	CIT	O7-C3-C4-C5
2	A	432	CIT	C6-C3-C4-C5
2	A	432	CIT	C2-C3-C6-O6
2	A	432	CIT	C2-C3-C6-O5
2	A	432	CIT	C4-C3-C6-O5
2	A	432	CIT	C4-C3-C6-O6
2	A	432	CIT	C1-C2-C3-C4
2	A	432	CIT	O7-C3-C6-O5
2	A	432	CIT	O7-C3-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	432	CIT	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	429/431 (99%)	0.69	38 (8%) 15 17	20, 46, 73, 81	3 (0%)

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	118	GLY	4.3
1	A	268	ALA	3.6
1	A	274	TRP	3.6
1	A	405	VAL	3.4
1	A	173	THR	3.4
1	A	277	ALA	3.1
1	A	117	CYS	3.1
1	A	34	LEU	3.1
1	A	272	GLY	3.0
1	A	188	LEU	2.8
1	A	184	ALA	2.7
1	A	183	ILE	2.7
1	A	259	VAL	2.7
1	A	172	TRP	2.7
1	A	410	PRO	2.7
1	A	409	ASP	2.6
1	A	403	GLY	2.6
1	A	348	LEU	2.5
1	A	178	LEU	2.5
1	A	267	GLN	2.5
1	A	270	GLN	2.4
1	A	175	LEU	2.4
1	A	65	VAL	2.4
1	A	148	VAL	2.3
1	A	417	ILE	2.2
1	A	287	LEU	2.2
1	A	37	ALA	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	404	ILE	2.1
1	A	131	SER	2.1
1	A	166	TRP	2.1
1	A	24	GLY	2.1
1	A	284	GLU	2.1
1	A	179	ALA	2.1
1	A	187	GLU	2.1
1	A	414	TRP	2.1
1	A	158	LEU	2.1
1	A	271	LEU	2.1
1	A	292	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CIT	A	432	13/13	0.79	0.14	56,61,67,68	0

6.5 Other polymers [i](#)

There are no such residues in this entry.