



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 06:47 PM UTC

PDB ID : 9CBT / pdb_00009cbt
Title : Crystal structure of human sirtuin 3 fragment (residues 118-399) bound to intermediates from reaction with NAD and inhibitor NH6-10
Authors : Fenwick, M.K.; Young, H.J.; Lin, H.
Deposited on : 2024-06-20
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	282	
1	B	282	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4855 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called NAD-dependent protein deacetylase sirtuin-3, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	270	Total	C	N	O	S	0	3	0
			2113	1362	362	380	9			
1	B	274	Total	C	N	O	S	0	10	0
			2211	1423	383	396	9			

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		

- Molecule 3 is 2-([(2S)-6-[(Z)-(1-[(2R,3R,4R,5R)-5-([(R)-[(R)-[(2R,3S,4R,5R)-5-(6-amino-9H-purin-9-yl)-3,4-dihydroxyoxolan-2-yl]methoxy}(hydroxy)phosphoryl]oxy}(hydroxy)phosphoryl]oxy)methyl)-4-hydroxy-2-sulfanyloxolan-3-yl]oxy}tetradecylidene)amino]-2-[(benzyloxy)carbonyl]amino}hexanoyl]amino}-N,N,N-triethylethan-1-aminium (non-preferred name) (CCD ID: 5IA) (formula: C₅₁H₈₆N₉O₁₆P₂S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	S	0	0
			79	51	9	16	2	1		


- Molecule 4 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: C₈H₁₈O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			13	8	5		

- Molecule 5 is (phenylmethyl) {N}-[(2 {S})-6-[[(2 {R},3 {a} {R},5 {R},6 {R},6 {a} {R})-5-[[[(2 {R},3 {S},4 {R},5 {R})-5-(6-aminopurin-9-yl)-3,4-bis(oxidanyl)oxolan-2-yl]methoxy-oxidanyl-phosphoryl]oxy-oxidanyl-phosphoryl]oxymethyl]-6-oxidanyl-2-tridecyl-3 {a},5,6,6 {

a}-tetrahydrofuro[2,3-d][1,3]oxathiol-2-yl]amino]-1-oxidanylidene-1-[2-(triethyl- $\text{l}^{\{4\}}$ -azan yl)ethylamino]hexan-2-yl]carbamate (CCD ID: 5I7) (formula: $\text{C}_{51}\text{H}_{86}\text{N}_9\text{O}_{16}\text{P}_2\text{S}$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	N	O	P	S		
5	B	1	79	51	9	16	2	1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	128	Total	O	0	1
			128	128		
6	B	230	Total	O	0	14
			230	230		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: NAD-dependent protein deacetylase sirtuin-3, mitochondrial



- Molecule 1: NAD-dependent protein deacetylase sirtuin-3, mitochondrial



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	39.43Å 63.19Å 114.61Å 90.00° 90.79° 90.00°	Depositor
Resolution (Å)	39.43 – 1.95 39.43 – 1.95	Depositor EDS
% Data completeness (in resolution range)	97.2 (39.43-1.95) 97.2 (39.43-1.95)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.51 (at 1.95Å)	Xtriage
Refinement program	PHENIX 1.19.1_4122	Depositor
R, R_{free}	0.198 , 0.239 0.197 , 0.239	Depositor DCC
R_{free} test set	2049 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	33.7	Xtriage
Anisotropy	0.524	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 48.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.077 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4855	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, 5I7, PG4, 5IA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.14	0/2168	0.32	0/2960
1	B	0.16	0/2267	0.34	0/3089
All	All	0.15	0/4435	0.33	0/6049

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2113	0	2094	15	0
1	B	2211	0	2224	11	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	79	0	0	1	0
4	B	13	0	18	0	0
5	B	79	0	0	1	0
6	A	128	0	0	1	0
6	B	230	0	0	1	0
All	All	4855	0	4336	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 3.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:130:GLU:OE1	1:B:133:ARG:NH1	2.38	0.56
1:A:230:ILE:HD13	1:A:251:PHE:HE2	1.71	0.55
1:A:181:GLU:HA	1:A:294:PHE:HB2	1.90	0.54
1:A:292:VAL:HG11	1:A:298:LEU:HD23	1.89	0.53
1:B:132:ILE:O	1:B:135:ARG:NH1	2.38	0.52
3:A:402:5IA:S34	3:A:402:5IA:C37	2.97	0.52
1:A:168:LEU:HD12	1:A:176:PRO:HB3	1.93	0.50
1:B:156:ASP:OD2	1:B:159:SER:OG	2.25	0.50
1:A:132:ILE:O	1:A:135:ARG:NH1	2.44	0.50
1:A:288:LYS:HD2	1:A:293:PHE:CE2	2.48	0.48
1:B:200:TYR:H	1:B:201:PRO:CD	2.27	0.47
1:A:374:VAL:HG13	1:A:379:TRP:HB2	1.97	0.47
1:B:366:VAL:HG22	5:B:403:5I7:N22	2.30	0.47
1:A:196:ALA:O	1:A:200:TYR:HB2	2.15	0.46
1:A:217:HIS:CD2	1:A:243:LYS:HG3	2.50	0.45
1:B:150:THR:HA	1:B:154:ILE:O	2.17	0.45
1:B:150:THR:OG1	1:B:151:PRO:HD3	2.17	0.44
1:A:355:PRO:HD2	6:A:570:HOH:O	2.18	0.44
1:A:353:TRP:C	1:A:355:PRO:HD3	2.44	0.43
1:B:182:LEU:HD22	1:B:296:GLU:HB2	2.01	0.43
1:A:217:HIS:HD2	1:A:243:LYS:HG3	1.85	0.42
1:B:196:ALA:O	1:B:200:TYR:HB2	2.19	0.42
1:A:212:PHE:N	1:A:383:MET:HE1	2.35	0.42
1:A:150:THR:OG1	1:A:151:PRO:HD3	2.20	0.41
1:B:302:PHE:CZ	1:B:330:LEU:HD11	2.56	0.41
1:A:150:THR:HA	1:A:154:ILE:O	2.21	0.41
1:B:139:ARG:NH2	6:B:523:HOH:O	2.53	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	270/282 (96%)	261 (97%)	9 (3%)	0	100	100
1	B	279/282 (99%)	271 (97%)	7 (2%)	1 (0%)	30	21
All	All	549/564 (97%)	532 (97%)	16 (3%)	1 (0%)	43	36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	200	TYR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	230/244 (94%)	229 (100%)	1 (0%)	84	85
1	B	245/244 (100%)	243 (99%)	2 (1%)	73	74
All	All	475/488 (97%)	472 (99%)	3 (1%)	78	79

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	200	TYR
1	B	200	TYR
1	B	351	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	217	HIS
1	A	362	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	5I7	B	403	-	80,84,84	1.78	17 (21%)	100,117,117	1.49	20 (20%)
3	5IA	A	402	-	81,83,83	3.08	31 (38%)	100,113,113	1.47	14 (14%)
4	PG4	B	402	-	12,12,12	0.16	0	11,11,11	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	5I7	B	403	-	-	23/69/116/116	0/6/6/6
3	5IA	A	402	-	-	14/74/108/108	0/5/5/5
4	PG4	B	402	-	-	5/10/10/10	-

All (48) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	402	5IA	P08-O07	11.65	1.72	1.59
3	A	402	5IA	P06-O07	10.99	1.71	1.59
3	A	402	5IA	P06-O05	6.88	1.86	1.59
3	A	402	5IA	C68-N69	6.81	1.49	1.33
3	A	402	5IA	C38-C37	5.80	1.59	1.50
5	B	403	5I7	P08-O07	5.71	1.65	1.59
3	A	402	5IA	C71-C70	5.43	1.62	1.52
3	A	402	5IA	P08-O10	5.06	1.79	1.59
3	A	402	5IA	C60-C61	5.02	1.62	1.50
3	A	402	5IA	C58-N57	4.95	1.47	1.34
5	B	403	5I7	P08-O10	4.53	1.77	1.59
5	B	403	5I7	C68-N69	4.38	1.43	1.33
3	A	402	5IA	C27-N18	4.19	1.44	1.37
5	B	403	5I7	C25-C19	4.19	1.46	1.39
3	A	402	5IA	C56-C68	3.91	1.62	1.52
5	B	403	5I7	P06-O07	3.68	1.63	1.59
5	B	403	5I7	P06-O05	3.65	1.73	1.59
3	A	402	5IA	C21-N22	3.60	1.40	1.33
3	A	402	5IA	O59-C58	3.54	1.42	1.35
3	A	402	5IA	C11-C12	3.42	1.61	1.51
3	A	402	5IA	C55-C56	3.34	1.61	1.53
3	A	402	5IA	C56-N57	3.27	1.52	1.45
3	A	402	5IA	C23-N24	3.15	1.42	1.34
5	B	403	5I7	C71-C70	3.07	1.58	1.52
5	B	403	5I7	C58-N57	2.84	1.41	1.34
3	A	402	5IA	C71-N72	2.76	1.61	1.52
5	B	403	5I7	C55-C56	2.69	1.59	1.53
3	A	402	5IA	C25-C19	2.63	1.43	1.39
3	A	402	5IA	C63-C62	2.62	1.43	1.38
5	B	403	5I7	O59-C58	2.61	1.40	1.35
3	A	402	5IA	C15-C13	2.59	1.60	1.53
5	B	403	5I7	C52-N51	2.59	1.50	1.46
5	B	403	5I7	C23-N24	2.53	1.40	1.34
3	A	402	5IA	C13-C12	2.49	1.59	1.53
3	A	402	5IA	C66-C61	2.48	1.43	1.38
3	A	402	5IA	C52-N51	2.48	1.53	1.46
5	B	403	5I7	P08-O29	-2.34	1.42	1.50
5	B	403	5I7	O32-C33	-2.32	1.39	1.43
3	A	402	5IA	C65-C66	2.27	1.42	1.38
5	B	403	5I7	C27-N18	2.24	1.41	1.37
3	A	402	5IA	C53-C52	2.22	1.59	1.51
3	A	402	5IA	C77-N72	2.19	1.61	1.52
3	A	402	5IA	C25-C23	2.18	1.47	1.41

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	402	5IA	C70-N69	2.15	1.51	1.46
3	A	402	5IA	O05-C04	-2.15	1.36	1.44
5	B	403	5I7	C60-C61	2.11	1.55	1.50
5	B	403	5I7	O05-C04	-2.11	1.36	1.44
3	A	402	5IA	C54-C55	2.01	1.60	1.52

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402	5IA	O59-C58-N57	6.18	123.67	110.45
3	A	402	5IA	O36-C35-C33	4.38	122.83	110.16
5	B	403	5I7	O36-C35-C33	4.22	108.64	103.80
5	B	403	5I7	O09-P08-O07	3.97	118.00	107.27
3	A	402	5IA	O59-C58-O67	-3.85	116.87	124.26
5	B	403	5I7	O36-C35-C02	3.58	117.44	110.20
3	A	402	5IA	O67-C58-N57	-3.30	119.46	124.86
5	B	403	5I7	O59-C58-N57	3.28	117.47	110.45
5	B	403	5I7	C15-C17-N18	3.17	121.17	113.30
3	A	402	5IA	C13-C15-C17	-3.07	95.66	101.46
3	A	402	5IA	O09-P08-O07	3.02	115.43	107.27
5	B	403	5I7	O09-P08-O29	2.98	126.32	112.44
3	A	402	5IA	C60-O59-C58	2.94	122.56	115.93
3	A	402	5IA	C25-N26-C27	2.87	107.97	103.45
3	A	402	5IA	C19-C25-N26	-2.77	107.42	110.58
5	B	403	5I7	O09-P08-O10	-2.61	95.75	107.57
5	B	403	5I7	C35-C02-C03	-2.58	96.44	101.99
5	B	403	5I7	O59-C60-C61	2.51	115.50	109.40
5	B	403	5I7	C25-C19-N20	-2.51	123.26	126.72
5	B	403	5I7	C60-O59-C58	2.49	121.54	115.93
5	B	403	5I7	C19-N18-C17	2.44	132.34	126.63
3	A	402	5IA	N18-C27-N26	-2.40	110.53	113.94
3	A	402	5IA	O01-C02-C03	2.35	117.82	111.08
3	A	402	5IA	C56-C68-N69	2.30	121.48	116.54
5	B	403	5I7	C38-C39-C40	2.29	120.00	113.19
5	B	403	5I7	O28-C17-N18	-2.28	103.72	108.09
3	A	402	5IA	C15-C13-C12	2.15	106.76	102.61
5	B	403	5I7	O28-C17-C15	-2.13	102.07	106.62
5	B	403	5I7	O59-C58-O67	-2.09	120.26	124.26
5	B	403	5I7	C71-C70-N69	-2.08	106.35	111.66
3	A	402	5IA	C25-C19-N18	2.08	108.08	105.81
5	B	403	5I7	O31-P06-O07	2.05	112.82	107.27
5	B	403	5I7	O28-C12-C13	2.05	109.22	105.15

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	403	5I7	C66-C61-C62	-2.05	115.19	118.23

There are no chirality outliers.

All (42) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	402	5IA	N51-C37-O36-C35
3	A	402	5IA	C04-O05-P06-O30
5	B	403	5I7	N69-C70-C71-N72
5	B	403	5I7	C70-C71-N72-C73
3	A	402	5IA	C38-C37-O36-C35
5	B	403	5I7	C74-C73-N72-C75
3	A	402	5IA	C37-C38-C39-C40
5	B	403	5I7	N51-C52-C53-C54
5	B	403	5I7	C76-C75-N72-C71
5	B	403	5I7	C38-C39-C40-C41
5	B	403	5I7	C76-C75-N72-C77
5	B	403	5I7	C76-C75-N72-C73
4	B	402	PG4	O3-C5-C6-O4
4	B	402	PG4	O2-C3-C4-O3
5	B	403	5I7	C74-C73-N72-C77
5	B	403	5I7	C74-C73-N72-C71
4	B	402	PG4	O4-C7-C8-O5
5	B	403	5I7	C45-C46-C47-C48
3	A	402	5IA	C43-C44-C45-C46
5	B	403	5I7	C46-C47-C48-C49
5	B	403	5I7	C44-C45-C46-C47
3	A	402	5IA	C38-C39-C40-C41
3	A	402	5IA	C52-C53-C54-C55
3	A	402	5IA	C47-C48-C49-C50
5	B	403	5I7	C39-C40-C41-C42
5	B	403	5I7	C40-C41-C42-C43
5	B	403	5I7	C70-C71-N72-C75
5	B	403	5I7	C70-C71-N72-C77
3	A	402	5IA	C44-C45-C46-C47
3	A	402	5IA	C40-C41-C42-C43
4	B	402	PG4	C5-C6-O4-C7
3	A	402	5IA	C42-C43-C44-C45
5	B	403	5I7	C78-C77-N72-C75
4	B	402	PG4	C6-C5-O3-C4
5	B	403	5I7	C42-C43-C44-C45
5	B	403	5I7	C78-C77-N72-C73

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
5	B	403	5I7	C78-C77-N72-C71
3	A	402	5IA	C41-C42-C43-C44
5	B	403	5I7	C47-C48-C49-C50
5	B	403	5I7	C53-C54-C55-C56
3	A	402	5IA	C45-C46-C47-C48
3	A	402	5IA	O10-C11-C12-O28

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	403	5I7	1	0
3	A	402	5IA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	270/282 (95%)	0.50	6 (2%) 62 69	20, 48, 77, 112	2 (0%)
1	B	274/282 (97%)	0.24	7 (2%) 57 64	12, 38, 70, 105	8 (2%)
All	All	544/564 (96%)	0.37	13 (2%) 59 66	12, 44, 76, 112	10 (1%)

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	174	PRO	3.7
1	A	176	PRO	3.4
1	B	200	TYR	3.2
1	A	199	LEU	3.0
1	B	199	LEU	2.6
1	A	251	PHE	2.6
1	B	385	ASP	2.5
1	B	175	TYR	2.5
1	A	391	THR	2.4
1	B	201	PRO	2.4
1	B	394	LEU	2.2
1	B	384	ARG	2.2
1	A	173	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	PG4	B	402	13/13	0.84	0.12	39,44,53,57	0
3	5IA	A	402	79/79	0.93	0.11	30,42,66,77	0
5	5I7	B	403	79/79	0.95	0.10	25,36,57,63	0
2	ZN	A	401	1/1	0.99	0.02	36,36,36,36	0
2	ZN	B	401	1/1	1.00	0.02	46,46,46,46	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 5I7 B 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN A 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ZN B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.