



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 05:21 AM UTC

PDB ID : 5EHN / pdb_00005ehn
Title : mAChE-syn TZ2PA5 complex
Authors : Bourne, Y.; Marchot, P.
Deposited on : 2015-10-28
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	543	6% 88% 10% .
1	B	543	15% 87% 10% ..

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8683 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acetylcholinesterase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	536	Total	C	N	O	S	0	2	0
			4190	2689	727	760	14			
1	B	531	Total	C	N	O	S	0	1	0
			4148	2666	715	753	14			

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is 6-phenyl-5-[5-[3-[2-(1,2,3,4-tetrahydroacridin-9-ylamino)ethyl]-1,2,3-triazol-4-yl]pentyl]phenanthridin-5-ium-3,8-diamine (CCD ID: 5NZ) (formula: $C_{41}H_{43}N_8$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	N	0	0
			49	41	8		
3	B	1	Total	C	N	0	0
			49	41	8		

- Molecule 4 is HEXAETHYLENE GLYCOL (CCD ID: P6G) (formula: C₁₂H₂₆O₇).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			19	12	7		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	126	Total 126	O 126	0	0
5	B	88	Total 88	O 88	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Acetylcholinesterase



• Molecule 1: Acetylcholinesterase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	79.75Å 112.45Å 227.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.00 – 2.60 45.00 – 2.69	Depositor EDS
% Data completeness (in resolution range)	98.5 (45.00-2.60) 98.1 (45.00-2.69)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.24 (at 2.69Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, R_{free}	0.182 , 0.203 0.234 , 0.224	Depositor DCC
R_{free} test set	1153 reflections (2.04%)	wwPDB-VP
Wilson B-factor (Å ²)	45.1	Xtriage
Anisotropy	0.784	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 58.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	8683	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, P6G, 5NZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.84	2/4320 (0.0%)	1.27	14/5903 (0.2%)
1	B	0.83	2/4275 (0.0%)	1.26	11/5846 (0.2%)
All	All	0.84	4/8595 (0.0%)	1.26	25/11749 (0.2%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	443	MET	SD-CE	-7.29	1.61	1.79
1	B	443	MET	SD-CE	-6.75	1.62	1.79
1	B	497	SER	CA-C	5.76	1.60	1.52
1	A	85	MET	SD-CE	-5.73	1.65	1.79

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	200	PHE	CA-CB-CG	7.42	121.22	113.80
1	A	200	PHE	CA-CB-CG	7.24	121.04	113.80
1	A	74	ASP	CA-CB-CG	7.07	119.67	112.60
1	A	350	ASN	CA-CB-CG	6.44	119.04	112.60
1	A	9	LEU	CA-C-N	6.28	131.88	122.58
1	A	9	LEU	C-N-CA	6.28	131.88	122.58
1	B	350	ASN	CA-CB-CG	6.13	118.73	112.60
1	A	422	GLY	N-CA-C	6.07	122.80	114.92
1	B	422	GLY	N-CA-C	5.87	122.55	114.92
1	A	5	ASP	CA-CB-CG	5.84	118.44	112.60
1	B	292	GLU	N-CA-C	-5.70	101.21	109.59
1	A	495	SER	CA-C-N	5.60	127.72	120.44
1	A	495	SER	C-N-CA	5.60	127.72	120.44
1	A	292	GLU	N-CA-C	-5.51	101.49	109.59
1	B	525	ARG	CA-C-N	5.47	127.93	120.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	525	ARG	C-N-CA	5.47	127.93	120.54
1	B	298	SER	N-CA-C	5.18	116.73	111.14
1	A	266	ASP	CA-CB-CG	5.09	117.69	112.60
1	B	175	ASP	N-CA-C	-5.09	105.62	111.07
1	A	526	ALA	N-CA-C	5.07	117.19	111.11
1	B	491	ASP	CA-CB-CG	5.06	117.66	112.60
1	A	75	THR	CA-C-N	5.03	127.33	120.54
1	A	75	THR	C-N-CA	5.03	127.33	120.54
1	B	267	THR	CA-C-N	5.03	126.97	120.44
1	B	267	THR	C-N-CA	5.03	126.97	120.44

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4190	0	4076	24	0
1	B	4148	0	4029	30	0
2	A	14	0	13	0	0
3	A	49	0	43	3	0
3	B	49	0	43	1	0
4	B	19	0	26	3	0
5	A	126	0	0	2	0
5	B	88	0	0	0	0
All	All	8683	0	8230	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:161:LEU:HD11	1:B:269:LEU:HD22	1.59	0.83
1:B:197:VAL:H	1:B:223:HIS:HD2	1.36	0.73

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:197:VAL:H	1:A:223:HIS:HD2	1.36	0.71
1:A:537:PRO:O	1:A:540:LEU:HG	1.92	0.69
1:B:211:MET:HG2	1:B:308:LEU:HD21	1.79	0.65
1:B:224:ARG:HG2	1:B:325:GLN:HB2	1.78	0.64
1:A:337:TYR:O	1:A:340:VAL:HG22	1.99	0.63
1:A:211:MET:HG2	1:A:308:LEU:HD21	1.80	0.63
1:B:197:VAL:H	1:B:223:HIS:CD2	2.19	0.60
1:A:197:VAL:H	1:A:223:HIS:CD2	2.18	0.59
1:A:296:ARG:HH21	1:A:369[B]:GLN:HE22	1.52	0.58
1:B:326:VAL:HG12	1:B:328:VAL:HG13	1.85	0.58
1:A:68:VAL:HG11	1:A:88:PRO:HB3	1.86	0.56
1:A:369[B]:GLN:HE21	1:A:369[B]:GLN:H	1.52	0.56
1:B:161:LEU:HD12	1:B:270:ILE:HD11	1.88	0.55
1:A:68:VAL:HG23	1:A:90:ARG:HB2	1.91	0.53
1:A:296:ARG:HH21	1:A:369[B]:GLN:NE2	2.07	0.53
1:B:68:VAL:HG11	1:B:88:PRO:HB3	1.89	0.53
1:B:68:VAL:HG23	1:B:90:ARG:HB2	1.92	0.52
1:B:166:GLU:HG2	1:B:270:ILE:HD13	1.91	0.51
1:A:376:GLU:O	1:A:380:LEU:HG	2.10	0.51
1:A:74:ASP:CG	3:A:1502:5NZ:H8	2.35	0.51
1:A:210:GLY:HA3	1:A:232:PRO:HD3	1.92	0.51
1:B:211:MET:HE3	1:B:232:PRO:HA	1.93	0.50
1:B:376:GLU:O	1:B:380:LEU:HG	2.11	0.50
1:A:380:LEU:HB2	4:B:1901:P6G:H142	1.93	0.50
3:A:1502:5NZ:H29	5:A:1622:HOH:O	2.11	0.49
1:B:210:GLY:HA3	1:B:232:PRO:HD3	1.93	0.49
1:A:211:MET:HE3	1:A:232:PRO:HA	1.94	0.49
1:B:380:LEU:HB2	4:B:1901:P6G:H81	1.94	0.49
1:B:381:HIS:HA	4:B:1901:P6G:H51	1.93	0.49
1:B:104:PRO:HD2	1:B:108:PRO:HD3	1.95	0.48
1:B:528:THR:O	1:B:531:PHE:HB3	2.14	0.48
1:B:328:VAL:O	1:B:427:ALA:HA	2.14	0.47
1:A:104:PRO:HD2	1:A:108:PRO:HD3	1.96	0.47
1:B:85:MET:HE1	1:B:132:VAL:HG11	1.96	0.47
1:B:48:MET:HE1	1:B:165:ARG:HH21	1.80	0.47
1:A:224:ARG:HG2	1:A:325:GLN:HB2	1.96	0.47
1:A:528:THR:O	1:A:531:PHE:HB3	2.15	0.47
1:A:369[B]:GLN:H	1:A:369[B]:GLN:NE2	2.13	0.46
1:A:328:VAL:O	1:A:427:ALA:HA	2.16	0.46
1:A:374:ALA:HA	1:A:539:LEU:HD23	1.98	0.45
1:B:161:LEU:HD12	1:B:270:ILE:CD1	2.47	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:227:LEU:HB2	1:B:328:VAL:HG12	1.98	0.45
1:A:293:SER:HB3	1:A:368:PRO:HG3	1.99	0.45
1:A:227:LEU:HB2	1:A:328:VAL:HG12	2.00	0.43
1:B:103:THR:HG21	1:B:190:PHE:HB3	2.01	0.43
1:B:337:TYR:O	1:B:340:VAL:HG22	2.19	0.43
1:B:340:VAL:HG11	1:B:443:MET:HE1	2.00	0.43
1:A:103:THR:HG21	1:A:190:PHE:HB3	2.01	0.42
1:B:257:CYS:HA	1:B:258:PRO:HA	1.85	0.42
1:B:293:SER:HB3	1:B:368:PRO:HG3	2.01	0.42
1:B:525:ARG:HG2	1:B:528:THR:HB	2.01	0.41
3:B:1902:5NZ:C40	3:B:1902:5NZ:H30	2.50	0.41
3:A:1502:5NZ:C29	5:A:1622:HOH:O	2.67	0.41
1:B:294:ILE:HG12	1:B:365:ILE:HG22	2.03	0.40
1:B:340:VAL:HG11	1:B:443:MET:CE	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	534/543 (98%)	522 (98%)	12 (2%)	0	100	100
1	B	528/543 (97%)	517 (98%)	11 (2%)	0	100	100
All	All	1062/1086 (98%)	1039 (98%)	23 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	440/443 (99%)	428 (97%)	12 (3%)	39	67
1	B	436/443 (98%)	426 (98%)	10 (2%)	44	71
All	All	876/886 (99%)	854 (98%)	22 (2%)	42	69

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	81	GLU
1	A	132	VAL
1	A	146	LEU
1	A	161	LEU
1	A	181	GLN
1	A	216	LEU
1	A	220	SER
1	A	386	LEU
1	A	421	GLN
1	A	522	ARG
1	A	536	LEU
1	A	540	LEU
1	B	132	VAL
1	B	146	LEU
1	B	165	ARG
1	B	166	GLU
1	B	181	GLN
1	B	251	LEU
1	B	332	LYS
1	B	421	GLN
1	B	470	ARG
1	B	536	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	181	GLN
1	A	223	HIS
1	A	317	ASN
1	B	181	GLN
1	B	223	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	265	ASN
1	B	317	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	A	1501	1	14,14,15	0.42	0	17,19,21	0.84	1 (5%)
3	5NZ	A	1502	-	54,56,56	2.24	12 (22%)	70,79,79	1.72	12 (17%)
4	P6G	B	1901	-	18,18,18	2.31	6 (33%)	17,17,17	1.50	3 (17%)
3	5NZ	B	1902	-	54,56,56	2.33	12 (22%)	70,79,79	1.75	11 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	1501	1	-	2/6/23/26	0/1/1/1
3	5NZ	A	1502	-	-	5/18/25/25	0/8/8/8
4	P6G	B	1901	-	-	6/16/16/16	-
3	5NZ	B	1902	-	-	4/18/25/25	0/8/8/8

All (30) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1902	5NZ	C30-C39	7.45	1.51	1.39
3	A	1502	5NZ	C39-C37	7.05	1.51	1.40
3	A	1502	5NZ	C30-C39	6.84	1.50	1.39
3	B	1902	5NZ	C39-C37	6.70	1.51	1.40
3	B	1902	5NZ	C5-C18	5.71	1.51	1.41
3	A	1502	5NZ	C5-C18	5.34	1.51	1.41
3	A	1502	5NZ	N6-N5	4.91	1.40	1.31
3	B	1902	5NZ	N6-N5	4.85	1.40	1.31
4	B	1901	P6G	O10-C9	4.63	1.62	1.42
3	B	1902	5NZ	C10-C4	4.61	1.49	1.42
3	A	1502	5NZ	C10-C4	4.59	1.49	1.42
3	B	1902	5NZ	C31-C33	4.47	1.49	1.42
3	A	1502	5NZ	C31-C33	4.07	1.48	1.42
4	B	1901	P6G	O13-C12	4.05	1.59	1.42
4	B	1901	P6G	O16-C15	3.78	1.58	1.42
3	B	1902	5NZ	N4-N5	3.78	1.40	1.35
4	B	1901	P6G	O19-C18	3.70	1.60	1.42
3	A	1502	5NZ	C11-C10	3.69	1.50	1.43
3	B	1902	5NZ	C11-C10	3.52	1.50	1.43
4	B	1901	P6G	O4-C3	3.49	1.57	1.42
3	A	1502	5NZ	C30-C31	3.31	1.49	1.43
3	B	1902	5NZ	C30-C31	3.27	1.49	1.43
3	A	1502	5NZ	C37-N8	3.23	1.36	1.32
3	B	1902	5NZ	C37-N8	3.20	1.36	1.32
3	B	1902	5NZ	C18-N3	-2.98	1.36	1.40
4	B	1901	P6G	O7-C6	2.90	1.54	1.42
3	A	1502	5NZ	N4-N5	2.62	1.39	1.35
3	A	1502	5NZ	C18-N3	-2.40	1.37	1.40
3	B	1902	5NZ	C27-C26	2.37	1.39	1.36
3	A	1502	5NZ	C27-C26	2.35	1.39	1.36

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1902	5NZ	C37-N8-C33	6.39	125.24	117.67
3	A	1502	5NZ	C5-C18-N3	6.29	122.52	118.46
3	A	1502	5NZ	C37-N8-C33	5.63	124.34	117.67
3	B	1902	5NZ	C5-C18-N3	5.28	121.87	118.46
3	B	1902	5NZ	C39-C37-N8	-5.27	119.93	123.66
3	A	1502	5NZ	C39-C37-N8	-4.69	120.34	123.66
4	B	1901	P6G	O1-C2-C3	3.32	131.38	111.82
3	B	1902	5NZ	C28-N4-C26	3.07	132.04	129.14
3	A	1502	5NZ	C28-N4-C26	3.01	131.97	129.14
3	A	1502	5NZ	C25-C26-C27	-2.99	126.51	131.35
3	B	1902	5NZ	C25-C26-C27	-2.88	126.68	131.35
4	B	1901	P6G	O13-C12-C11	2.57	122.04	110.35
3	B	1902	5NZ	C13-C12-C11	-2.51	116.77	120.25
2	A	1501	NAG	C1-O5-C5	2.48	115.51	112.19
3	B	1902	5NZ	C31-C33-N8	-2.44	120.22	122.80
3	A	1502	5NZ	C27-C26-N4	2.38	106.93	103.90
3	A	1502	5NZ	C31-C30-N7	-2.36	114.53	121.86
3	A	1502	5NZ	C24-C25-C26	-2.32	109.86	113.63
3	B	1902	5NZ	C10-C4-C5	-2.27	116.95	119.93
3	B	1902	5NZ	C17-C12-C13	2.20	121.62	117.68
3	A	1502	5NZ	C32-C31-C30	-2.18	120.78	124.70
3	A	1502	5NZ	C28-C29-N7	2.13	116.39	110.29
3	B	1902	5NZ	C40-C39-C37	-2.11	118.91	120.90
4	B	1901	P6G	O7-C6-C5	2.11	119.98	110.35
3	A	1502	5NZ	C40-C39-C37	-2.06	118.96	120.90
3	A	1502	5NZ	C10-C4-C5	-2.01	117.29	119.93
3	B	1902	5NZ	C3-C4-C10	2.00	120.20	117.72

There are no chirality outliers.

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1502	5NZ	C29-C28-N4-N5
3	A	1502	5NZ	N4-C28-C29-N7
3	B	1902	5NZ	N4-C28-C29-N7
4	B	1901	P6G	O1-C2-C3-O4
4	B	1901	P6G	O13-C14-C15-O16
4	B	1901	P6G	O16-C17-C18-O19
2	A	1501	NAG	C4-C5-C6-O6
3	B	1902	5NZ	C29-C28-N4-N5
4	B	1901	P6G	C11-C12-O13-C14
4	B	1901	P6G	O7-C8-C9-O10
3	A	1502	5NZ	C31-C30-N7-C29

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	1501	NAG	O5-C5-C6-O6
3	A	1502	5NZ	C20-C21-C22-C24
3	A	1502	5NZ	C39-C30-N7-C29
4	B	1901	P6G	C6-C5-O4-C3
3	B	1902	5NZ	C31-C30-N7-C29
3	B	1902	5NZ	C39-C30-N7-C29

There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1502	5NZ	3	0
4	B	1901	P6G	3	0
3	B	1902	5NZ	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	536/543 (98%)	0.54	34 (6%) 26 20	29, 46, 74, 116	2 (0%)
1	B	531/543 (97%)	1.24	80 (15%) 5 4	32, 52, 81, 106	1 (0%)
All	All	1067/1086 (98%)	0.89	114 (10%) 11 9	29, 48, 79, 116	3 (0%)

All (114) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	540	LEU	8.6
1	B	76	LEU	7.5
1	B	72	TYR	6.7
1	B	497	SER	6.6
1	A	341	TYR	6.2
1	A	258	PRO	5.9
1	B	124	TYR	5.9
1	B	341	TYR	5.5
1	A	72	TYR	5.1
1	B	495	SER	5.0
1	B	337	TYR	4.9
1	A	493	ARG	4.6
1	A	264	GLY	4.6
1	B	336	SER	4.6
1	A	495	SER	4.5
1	A	541	SER	4.4
1	A	540	LEU	4.3
1	B	345	GLY	4.2
1	B	447	HIS	4.2
1	B	492	PRO	4.1
1	B	78	PRO	3.9
1	B	493	ARG	3.8
1	B	340	VAL	3.7
1	B	496	LYS	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	347	SER	3.6
1	A	347	SER	3.5
1	B	344	PRO	3.5
1	B	346	PHE	3.5
1	A	124	TYR	3.4
1	B	258	PRO	3.4
1	B	269	LEU	3.4
1	B	335	GLY	3.3
1	B	339	LEU	3.3
1	B	122	GLY	3.3
1	B	349	ASP	3.3
1	B	23	LYS	3.3
1	B	498	PRO	3.3
1	B	363	VAL	3.2
1	B	460	ASP	3.2
1	A	4	GLU	3.1
1	B	342	GLY	3.1
1	A	447	HIS	3.1
1	B	74	ASP	3.0
1	A	323	ASP	3.0
1	B	77	TYR	2.9
1	B	457	LEU	2.9
1	A	336	SER	2.9
1	B	265	ASN	2.9
1	A	144	ALA	2.9
1	B	367	VAL	2.9
1	B	446	PRO	2.8
1	A	462	SER	2.8
1	B	165	ARG	2.8
1	B	458	PRO	2.8
1	A	76	LEU	2.8
1	B	463	LEU	2.7
1	B	280	ASP	2.7
1	A	23	LYS	2.7
1	B	395	ARG	2.7
1	A	2	GLY	2.7
1	A	3	ARG	2.6
1	A	345	GLY	2.6
1	B	286	TRP	2.6
1	B	107	ARG	2.6
1	B	449	TYR	2.6
1	B	539	LEU	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	265	ASN	2.6
1	A	492	PRO	2.6
1	A	539	LEU	2.5
1	A	496	LYS	2.5
1	A	342	GLY	2.5
1	B	75	THR	2.5
1	B	462	SER	2.5
1	B	424	ARG	2.5
1	A	7	GLN	2.5
1	B	68	VAL	2.4
1	A	216	LEU	2.4
1	B	22	LEU	2.4
1	B	146	LEU	2.4
1	B	451	ILE	2.4
1	B	454	ILE	2.4
1	B	104	PRO	2.4
1	B	285	GLU	2.3
1	B	323	ASP	2.3
1	A	522	ARG	2.3
1	B	73	VAL	2.3
1	A	386	LEU	2.3
1	B	271	ALA	2.3
1	B	358	GLN	2.3
1	B	401	VAL	2.3
1	A	109	ALA	2.3
1	B	238	THR	2.3
1	B	10	VAL	2.3
1	B	92	LEU	2.3
1	B	313	GLU	2.2
1	B	389	GLU	2.2
1	B	465	TYR	2.2
1	B	34	GLY	2.2
1	B	79	GLY	2.2
1	B	360	LEU	2.2
1	B	190	PHE	2.2
1	B	109	ALA	2.2
1	B	253	ARG	2.2
1	B	43	GLY	2.1
1	B	293	SER	2.1
1	B	70	TYR	2.1
1	B	456	GLY	2.1
1	B	81	GLU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	351	GLU	2.1
1	B	284	HIS	2.1
1	A	337	TYR	2.1
1	B	455	PHE	2.1
1	A	219[A]	ARG	2.0
1	A	463	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	5NZ	B	1902	49/49	0.61	0.28	35,52,89,91	0
2	NAG	A	1501	14/15	0.71	0.17	77,85,87,88	0
3	5NZ	A	1502	49/49	0.73	0.24	28,42,79,81	0
4	P6G	B	1901	19/19	0.83	0.18	52,62,69,69	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 5NZ B 1902:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 5NZ A 1502:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around P6G B 1901:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.