



Full wwPDB NMR Structure Validation Report ⓘ

Mar 10, 2026 – 07:58 AM UTC

PDB ID : 1FKC / pdb_00001fkc
Title : HUMAN PRION PROTEIN (MUTANT E200K) FRAGMENT 90-231
Authors : Zhang, Y.; Swietnicki, W.; Zagorski, M.G.; Surewicz, W.K.; Soennichsen, F.D.
Deposited on : 2000-08-09

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	142	

2 Ensemble composition and analysis ⓘ

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1731 atoms, of which 837 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called PRION PROTEIN.

Mol	Chain	Residues	Atoms						Trace
1	A	107	Total	C	H	N	O	S	0
			1731	553	837	157	175	9	


There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	200	LYS	GLU	engineered mutation	UNP P04156

4 Residue-property plots [i](#)

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: PRION PROTEIN

Chain A: 



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing molecular dynamics*.

Of the 60 calculated structures, 1 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	structure solution	0.9
ARIA	structure solution	0.9
CNS	refinement	0.9

No chemical shift data was provided.

6 Model quality

6.1 Standard geometry

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	894	837	833	14
All	All	894	837	833	14

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:A:176:VAL:O	1:A:180:VAL:HG23	0.58	1.99
1:A:129:MET:HA	1:A:129:MET:HE2	0.53	1.80
1:A:212:GLN:HA	1:A:215:ILE:HD12	0.52	1.80
1:A:150:TYR:HB2	1:A:205:MET:SD	0.51	2.45
1:A:130:LEU:HD23	1:A:131:GLY:N	0.49	2.23
1:A:186:GLN:O	1:A:190:THR:HG22	0.46	2.10
1:A:173:ASN:O	1:A:177:HIS:HB2	0.45	2.12
1:A:216:THR:O	1:A:220:ARG:HG3	0.44	2.12
1:A:129:MET:O	1:A:162:TYR:HA	0.44	2.12
1:A:125:LEU:HG	1:A:126:GLY:N	0.43	2.28
1:A:161:VAL:HG12	1:A:183:THR:HG21	0.42	1.91
1:A:139:ILE:HG13	1:A:209:VAL:HG22	0.41	1.92
1:A:173:ASN:CG	1:A:174:ASN:N	0.40	2.79
1:A:215:ILE:O	1:A:218:TYR:HB3	0.40	2.16

6.3 Torsion angles

6.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	105/142 (74%)	92 (88%)	12 (11%)	1 (1%)	15	65
All	All	105/142 (74%)	92 (88%)	12 (11%)	1 (1%)	15	65

All 1 Ramachandran outliers are listed below.

Mol	Chain	Res	Type
1	A	195	GLY

6.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	99/120 (82%)	71 (72%)	28 (28%)	1	19
All	All	99/120 (82%)	71 (72%)	28 (28%)	1	19

All 28 residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	125	LEU
1	A	129	MET
1	A	134	MET
1	A	136	ARG
1	A	139	ILE
1	A	151	ARG
1	A	159	ASN
1	A	164	ARG
1	A	166	MET
1	A	168	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	172	GLN
1	A	177	HIS
1	A	182	ILE
1	A	184	ILE
1	A	185	LYS
1	A	186	GLN
1	A	194	LYS
1	A	196	GLU
1	A	199	THR
1	A	200	LYS
1	A	202	ASP
1	A	204	LYS
1	A	205	MET
1	A	211	GLU
1	A	212	GLN
1	A	221	GLU
1	A	228	ARG
1	A	231	SER

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided