



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:39 AM UTC

PDB ID : 1FL1 / pdb_00001fl1
Title : KSHV PROTEASE
Authors : Reiling, K.K.; Pray, T.R.; Craik, C.S.; Stroud, R.M.
Deposited on : 2000-08-11
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	230	10% 72% 11% 17%
1	B	230	41% 74% 12% 10%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3157 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	192	Total	C	N	O	S	0	0	0
			1437	926	246	261	4			
1	B	207	Total	C	N	O	S	0	0	0
			1550	1001	259	286	4			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	204	GLY	SER	engineered mutation	UNP O36607
B	204	GLY	SER	engineered mutation	UNP O36607

- Molecule 2 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	K	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	83	Total	O	0	0
			83	83		
3	B	86	Total	O	0	0
			86	86		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: PROTEASE



• Molecule 1: PROTEASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	53.58Å 53.58Å 323.06Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	28.00 – 2.20 28.00 – 2.20	Depositor EDS
% Data completeness (in resolution range)	(Not available) (28.00-2.20) 97.5 (28.00-2.20)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	9.85 (at 2.20Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.225 , 0.260 0.283 , 0.302	Depositor DCC
R_{free} test set	1642 reflections (5.87%)	wwPDB-VP
Wilson B-factor (Å ²)	46.2	Xtriage
Anisotropy	0.005	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 57.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.056 for -h,-k,l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	3157	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.44	0/1470	0.88	2/2005 (0.1%)
1	B	0.48	0/1587	0.93	4/2168 (0.2%)
All	All	0.46	0/3057	0.91	6/4173 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	11	VAL	N-CA-C	-7.94	104.78	113.43
1	B	11	VAL	N-CA-C	-7.48	104.02	113.22
1	B	118	ILE	CA-C-N	-6.37	102.16	121.98
1	B	118	ILE	C-N-CA	-6.37	102.16	121.98
1	A	137	LEU	N-CA-C	-5.82	99.81	109.46
1	B	137	LEU	N-CA-C	-5.64	100.10	109.46

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1437	0	1427	15	0
1	B	1550	0	1536	23	0
2	B	1	0	0	0	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	83	0	0	0	1
3	B	86	0	0	5	1
All	All	3157	0	2963	38	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:226:VAL:HB	1:B:228:LEU:HD23	1.77	0.65
1:B:177:SER:HB3	3:B:1076:HOH:O	1.99	0.63
1:B:89:VAL:O	1:B:99:LYS:HE3	1.99	0.62
1:B:107:HIS:CE1	1:B:228:LEU:HD22	2.39	0.58
1:B:29:GLN:HG3	1:B:31:THR:H	1.70	0.56
1:B:79:LEU:HD22	1:B:83:LEU:HD22	1.88	0.55
1:B:102:LEU:O	1:B:105:ILE:HG22	2.07	0.54
1:B:209:ARG:HD3	3:B:1040:HOH:O	2.08	0.54
1:A:181:LEU:C	1:A:183:ASP:H	2.17	0.52
1:B:97:LEU:HD21	1:B:225:PRO:HG2	1.91	0.51
1:A:73:SER:HB3	1:A:76:PHE:HB3	1.92	0.51
1:B:144:ARG:HH12	1:B:212:LEU:HB3	1.78	0.49
1:B:96:ASN:ND2	3:B:1018:HOH:O	2.26	0.49
1:B:144:ARG:NE	1:B:144:ARG:HA	2.28	0.48
1:A:2:ALA:HA	1:A:152:HIS:HB3	1.95	0.48
1:A:100:GLU:O	1:A:104:GLU:HG3	2.14	0.48
1:A:5:LEU:HD23	1:A:71:ILE:HB	1.96	0.47
1:A:84:ALA:O	1:A:90:ALA:HB2	2.14	0.47
1:A:113:LEU:HG	1:A:148:ALA:HB2	1.98	0.46
1:A:209:ARG:HD3	3:B:1042:HOH:O	2.16	0.46
1:B:147:VAL:HG22	3:B:1013:HOH:O	2.16	0.44
1:A:82:ARG:HH11	1:A:82:ARG:HG2	1.82	0.44
1:B:144:ARG:NH1	1:B:212:LEU:HB3	2.32	0.44
1:B:160:ARG:HG3	1:B:226:VAL:CG1	2.49	0.43
1:B:118:ILE:H	1:B:118:ILE:HG12	1.42	0.42
1:A:5:LEU:HB3	1:A:71:ILE:HB	2.00	0.42
1:B:113:LEU:HG	1:B:148:ALA:HB2	2.01	0.42
1:A:116:SER:HB3	1:A:134:HIS:CE1	2.55	0.42
1:B:41:PRO:HD2	1:B:133:GLN:O	2.20	0.41
1:A:181:LEU:C	1:A:183:ASP:N	2.79	0.41
1:B:182:GLU:O	1:B:182:GLU:CG	2.68	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:63:HIS:HD2	1:B:163:SER:O	2.03	0.41
1:B:153:ASP:O	1:B:157:VAL:HG23	2.19	0.41
1:A:6:TYR:CE1	1:A:181:LEU:HD11	2.56	0.41
1:A:82:ARG:HG2	1:A:82:ARG:NH1	2.36	0.41
1:A:186:THR:HA	1:A:187:PRO:HD3	1.87	0.41
1:B:140:LEU:CD1	1:B:213:LEU:HD13	2.51	0.41
1:B:213:LEU:HD12	1:B:213:LEU:HA	1.88	0.41

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:1001:K:K	2:B:1001:K:K[6_765]	0.96	1.24
3:A:290:HOH:O	3:A:290:HOH:O[5_555]	2.04	0.16
3:B:1064:HOH:O	3:B:1064:HOH:O[6_765]	2.08	0.12

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	186/230 (81%)	177 (95%)	8 (4%)	1 (0%)	24	27
1	B	201/230 (87%)	195 (97%)	5 (2%)	1 (0%)	24	27
All	All	387/460 (84%)	372 (96%)	13 (3%)	2 (0%)	24	27

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	32	ASP
1	A	182	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	153/200 (76%)	151 (99%)	2 (1%)	61	76
1	B	168/200 (84%)	157 (94%)	11 (6%)	15	18
All	All	321/400 (80%)	308 (96%)	13 (4%)	28	38

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	79	LEU
1	A	226	VAL
1	B	5	LEU
1	B	29	GLN
1	B	32	ASP
1	B	79	LEU
1	B	83	LEU
1	B	102	LEU
1	B	117	SER
1	B	118	ILE
1	B	160	ARG
1	B	213	LEU
1	B	228	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	188	ASN
1	B	63	HIS
1	B	133	GLN
1	B	188	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	192/230 (83%)	0.99	23 (11%) 9 6	32, 45, 75, 81	0
1	B	207/230 (90%)	1.98	94 (45%) 0 0	34, 48, 72, 92	0
All	All	399/460 (86%)	1.50	117 (29%) 1 1	32, 46, 75, 92	0

All (117) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	15	SER	5.7
1	B	181	LEU	4.8
1	B	117	SER	4.8
1	B	2	ALA	4.4
1	B	185	SER	4.3
1	B	118	ILE	4.1
1	B	59	PHE	4.1
1	B	179	CYS	4.1
1	B	157	VAL	4.1
1	B	130	PRO	3.9
1	A	183	ASP	3.9
1	B	187	PRO	3.9
1	B	119	HIS	3.8
1	B	33	TYR	3.8
1	B	69	GLY	3.8
1	B	120	PRO	3.8
1	A	181	LEU	3.8
1	B	54	TRP	3.7
1	B	57	GLY	3.6
1	A	184	LEU	3.6
1	B	158	VAL	3.6
1	B	213	LEU	3.4
1	B	35	PRO	3.4
1	B	184	LEU	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	173	LEU	3.3
1	B	31	THR	3.2
1	B	36	VAL	3.2
1	B	80	ALA	3.2
1	B	177	SER	3.1
1	B	8	GLY	3.1
1	B	14	VAL	3.1
1	B	34	LEU	3.1
1	B	178	SER	3.0
1	B	132	PHE	3.0
1	B	94	VAL	3.0
1	B	116	SER	3.0
1	A	147	VAL	3.0
1	B	172	ILE	3.0
1	A	13	VAL	3.0
1	B	61	VAL	3.0
1	B	149	VAL	3.0
1	B	93	PRO	3.0
1	B	193	LEU	2.9
1	B	180	ARG	2.9
1	B	230	ALA	2.9
1	A	39	PRO	2.9
1	B	12	ASP	2.9
1	B	154	ALA	2.9
1	B	168	GLU	2.8
1	B	192	PRO	2.8
1	B	176	VAL	2.8
1	B	160	ARG	2.8
1	B	29	GLN	2.8
1	B	102	LEU	2.7
1	B	58	LEU	2.7
1	A	175	HIS	2.6
1	A	208	ASP	2.6
1	A	47	LEU	2.6
1	B	9	GLY	2.6
1	B	65	ILE	2.6
1	B	161	PHE	2.6
1	B	223	LEU	2.5
1	B	97	LEU	2.5
1	B	68	THR	2.5
1	B	175	HIS	2.5
1	B	91	ARG	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	113	LEU	2.5
1	B	13	VAL	2.5
1	A	179	CYS	2.4
1	B	43	THR	2.4
1	B	156	TRP	2.4
1	A	40	LEU	2.4
1	B	106	LEU	2.4
1	B	92	ALA	2.4
1	B	77	LEU	2.4
1	B	174	GLN	2.4
1	B	165	SER	2.3
1	A	132	PHE	2.3
1	A	138	CYS	2.3
1	B	131	VAL	2.3
1	B	55	THR	2.3
1	B	224	SER	2.3
1	A	14	VAL	2.3
1	A	38	GLU	2.3
1	B	42	ILE	2.3
1	B	41	PRO	2.3
1	B	163	SER	2.3
1	A	3	GLN	2.3
1	B	133	GLN	2.3
1	B	153	ASP	2.2
1	B	90	ALA	2.2
1	A	193	LEU	2.2
1	B	56	LEU	2.2
1	B	66	PHE	2.2
1	B	225	PRO	2.2
1	B	67	CYS	2.2
1	B	221	SER	2.2
1	A	188	ASN	2.2
1	B	30	VAL	2.1
1	B	89	VAL	2.1
1	B	10	PHE	2.1
1	B	169	ARG	2.1
1	B	96	ASN	2.1
1	B	208	ASP	2.1
1	B	164	VAL	2.1
1	B	112	GLY	2.1
1	B	73	SER	2.1
1	B	170	ALA	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	171	HIS	2.1
1	A	94	VAL	2.1
1	A	2	ALA	2.1
1	B	46	HIS	2.1
1	A	92	ALA	2.0
1	B	84	ALA	2.0
1	B	134	HIS	2.0
1	B	182	GLU	2.0
1	A	211	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	K	B	1001	1/1	0.94	0.08	47,47,47,47	0

6.5 Other polymers [i](#)

There are no such residues in this entry.