



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 04:44 PM UTC

PDB ID : 3GA6 / pdb_00003ga6
Title : Mth0212 in complex with two DNA helices
Authors : Lakomek, K.; Dickmanns, A.; Ficner, R.
Deposited on : 2009-02-16
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	265	
1	B	265	
2	F	12	
2	H	12	
3	D	12	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	G	12	 67% 33%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	7002	-	-	X	-
4	GOL	A	7031	-	-	X	-
4	GOL	B	7022	-	-	X	-
4	GOL	B	7033	-	-	X	-

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 5629 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Exodeoxyribonuclease.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	254	Total	C	N	O	S	0	0	0
			2122	1362	372	380	8			
1	B	255	Total	C	N	O	S	0	0	0
			2127	1365	373	381	8			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1002	ALA	THR	engineered mutation	UNP O26314
A	1151	ASN	ASP	engineered mutation	UNP O26314
A	1258	LEU	-	expression tag	UNP O26314
A	1259	GLU	-	expression tag	UNP O26314
A	1260	HIS	-	expression tag	UNP O26314
A	1261	HIS	-	expression tag	UNP O26314
A	1262	HIS	-	expression tag	UNP O26314
A	1263	HIS	-	expression tag	UNP O26314
A	1264	HIS	-	expression tag	UNP O26314
A	1265	HIS	-	expression tag	UNP O26314
B	2002	ALA	THR	engineered mutation	UNP O26314
B	2151	ASN	ASP	engineered mutation	UNP O26314
B	2258	LEU	-	expression tag	UNP O26314
B	2259	GLU	-	expression tag	UNP O26314
B	2260	HIS	-	expression tag	UNP O26314
B	2261	HIS	-	expression tag	UNP O26314
B	2262	HIS	-	expression tag	UNP O26314
B	2263	HIS	-	expression tag	UNP O26314
B	2264	HIS	-	expression tag	UNP O26314
B	2265	HIS	-	expression tag	UNP O26314


- Molecule 2 is a DNA chain called 5'-D(*GP*CP*TP*GP*CP*GP*CP*AP*GP*GP*GP*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	F	11	Total	C	N	O	P	0	0	0
			227	107	46	64	10			
2	H	11	Total	C	N	O	P	0	0	0
			227	106	44	66	11			

- Molecule 3 is a DNA chain called 5'-D(*GP*CP*CP*CP*TP*GP*UP*GP*CP*AP*GP*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	G	12	Total	C	N	O	P	0	0	1
			207	95	36	65	11			
3	D	12	Total	C	N	O	P	0	0	0
			240	114	44	71	11			

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Na	0	0
			1	1		

- Molecule 6 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	O	P	0	0
			5	4	1		
6	B	1	Total	O	P	0	0
			5	4	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	157	Total	O	0	0
			157	157		
7	B	155	Total	O	0	0
			155	155		
7	F	6	Total	O	0	0
			6	6		
7	G	13	Total	O	0	0
			13	13		
7	H	16	Total	O	0	0
			16	16		
7	D	7	Total	O	0	0
			7	7		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Exodeoxyribonuclease



• Molecule 1: Exodeoxyribonuclease



• Molecule 2: 5'-D(*GP*CP*TP*GP*CP*GP*CP*AP*GP*GP*GP*C)-3'



• Molecule 2: 5'-D(*GP*CP*TP*GP*CP*GP*CP*AP*GP*GP*GP*C)-3'






- Molecule 3: 5'-D(*GP*CP*CP*CP*TP*GP*UP*GP*CP*AP*GP*C)-3'



- Molecule 3: 5'-D(*GP*CP*CP*CP*TP*GP*UP*GP*CP*AP*GP*C)-3'



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	54.83Å 126.66Å 54.83Å 90.00° 93.14° 90.00°	Depositor
Resolution (Å)	33.71 – 1.90 33.71 – 1.90	Depositor EDS
% Data completeness (in resolution range)	89.4 (33.71-1.90) 95.5 (33.71-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.09 (at 1.89Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.155 , 0.203 0.159 , 0.199	Depositor DCC
R_{free} test set	3397 reflections (6.04%)	wwPDB-VP
Wilson B-factor (Å ²)	27.7	Xtriage
Anisotropy	0.334	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 42.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.488 for l,-k,h	Xtriage
Reported twinning fraction	0.505 for l,-k,h	Depositor
Outliers	1 of 56227 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	5629	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.90% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, NA, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.48	0/2180	0.87	5/2939 (0.2%)
1	B	0.47	0/2185	0.82	4/2946 (0.1%)
2	F	0.25	0/255	0.80	0/393
2	H	0.33	0/254	1.04	1/390 (0.3%)
3	D	0.46	0/268	1.09	1/411 (0.2%)
3	G	0.92	1/230 (0.4%)	1.30	1/354 (0.3%)
All	All	0.49	1/5372 (0.0%)	0.89	12/7433 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	G	1	DG	O3'-P	-8.27	1.48	1.61

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	G	11	DG	P-O3'-C3'	11.74	137.81	120.20
1	A	1061	THR	CA-C-N	5.87	126.06	120.31
1	A	1061	THR	C-N-CA	5.87	126.06	120.31
1	A	1078	PRO	O-C-N	5.83	123.99	121.31
1	B	2176	VAL	CB-CA-C	-5.67	104.62	111.88

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2122	0	2060	62	0
1	B	2127	0	2065	83	0
2	F	227	0	124	11	0
2	H	227	0	123	6	0
3	D	240	0	134	23	0
3	G	207	0	110	6	0
4	A	66	0	88	24	0
4	B	48	0	64	20	0
5	A	1	0	0	0	0
6	B	10	0	0	0	0
7	A	157	0	0	2	0
7	B	155	0	0	7	0
7	D	7	0	0	0	0
7	F	6	0	0	0	0
7	G	13	0	0	0	0
7	H	16	0	0	0	0
All	All	5629	0	4768	186	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 186 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:G:7:DU:H3	2:H:5:DG:H22	0.99	0.96
1:B:2175:PRO:HG2	4:B:7032:GOL:H12	1.46	0.94
1:B:2193:PHE:HB2	4:B:7033:GOL:H11	1.52	0.90
2:F:6:DG:H2''	2:F:7:DC:H5'	1.55	0.88
1:A:1131:ALA:HB1	4:A:7002:GOL:H31	1.56	0.87

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	252/265 (95%)	243 (96%)	8 (3%)	1 (0%)	30	22
1	B	253/265 (96%)	243 (96%)	10 (4%)	0	100	100
All	All	505/530 (95%)	486 (96%)	18 (4%)	1 (0%)	43	36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1216	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	225/235 (96%)	209 (93%)	16 (7%)	13	7
1	B	225/235 (96%)	208 (92%)	17 (8%)	12	5
All	All	450/470 (96%)	417 (93%)	33 (7%)	13	6

5 of 33 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	2214	GLU
1	B	2220	ARG
1	B	2255	ILE
1	A	1204	THR
1	A	1166	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1012	ASN
1	A	1145	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	2037	GLN
1	B	2137	ASN
1	B	2202	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 22 ligands modelled in this entry, 1 is monoatomic - leaving 21 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	A	7012	-	5,5,5	0.47	0	5,5,5	0.21	0
4	GOL	B	7032	-	5,5,5	0.40	0	5,5,5	0.24	0
4	GOL	A	7031	-	5,5,5	0.65	0	5,5,5	0.40	0
4	GOL	A	7020	-	5,5,5	0.43	0	5,5,5	0.44	0
4	GOL	B	7011	-	5,5,5	0.37	0	5,5,5	0.25	0
6	PO4	B	7205	-	4,4,4	0.94	0	6,6,6	0.48	0
4	GOL	A	7024	-	5,5,5	0.37	0	5,5,5	0.14	0
4	GOL	A	7030	-	5,5,5	0.44	0	5,5,5	0.41	0
4	GOL	B	7003	-	5,5,5	0.33	0	5,5,5	0.23	0
4	GOL	A	7004	-	5,5,5	0.44	0	5,5,5	0.43	0
4	GOL	A	7002	-	5,5,5	0.55	0	5,5,5	0.44	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	B	7021	-	5,5,5	0.45	0	5,5,5	0.27	0
4	GOL	A	7010	-	5,5,5	0.41	0	5,5,5	0.58	0
4	GOL	A	7013	-	5,5,5	0.32	0	5,5,5	0.47	0
4	GOL	A	7015	-	5,5,5	0.48	0	5,5,5	0.73	0
4	GOL	B	7022	-	5,5,5	0.49	0	5,5,5	0.60	0
6	PO4	B	7204	-	4,4,4	0.90	0	6,6,6	0.61	0
4	GOL	B	7027	-	5,5,5	0.34	0	5,5,5	0.33	0
4	GOL	A	7006	5	5,5,5	0.35	0	5,5,5	0.45	0
4	GOL	B	7009	-	5,5,5	0.36	0	5,5,5	0.27	0
4	GOL	B	7033	-	5,5,5	0.43	0	5,5,5	0.34	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	7012	-	-	2/4/4/4	-
4	GOL	B	7032	-	-	2/4/4/4	-
4	GOL	A	7031	-	-	1/4/4/4	-
4	GOL	A	7020	-	-	2/4/4/4	-
4	GOL	B	7011	-	-	2/4/4/4	-
4	GOL	A	7024	-	-	4/4/4/4	-
4	GOL	A	7030	-	-	0/4/4/4	-
4	GOL	B	7003	-	-	0/4/4/4	-
4	GOL	A	7004	-	-	4/4/4/4	-
4	GOL	A	7002	-	-	4/4/4/4	-
4	GOL	B	7021	-	-	2/4/4/4	-
4	GOL	A	7010	-	-	3/4/4/4	-
4	GOL	A	7013	-	-	2/4/4/4	-
4	GOL	A	7015	-	-	1/4/4/4	-
4	GOL	B	7022	-	-	2/4/4/4	-
4	GOL	B	7027	-	-	0/4/4/4	-
4	GOL	A	7006	5	-	4/4/4/4	-
4	GOL	B	7009	-	-	2/4/4/4	-
4	GOL	B	7033	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 37 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	7002	GOL	O1-C1-C2-C3
4	A	7004	GOL	O1-C1-C2-C3
4	A	7006	GOL	O1-C1-C2-C3
4	A	7020	GOL	O1-C1-C2-C3
4	A	7024	GOL	O1-C1-C2-C3

There are no ring outliers.

14 monomers are involved in 44 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	7012	GOL	2	0
4	B	7032	GOL	2	0
4	A	7031	GOL	5	0
4	A	7020	GOL	1	0
4	B	7011	GOL	2	0
4	A	7030	GOL	3	0
4	B	7003	GOL	2	0
4	A	7004	GOL	2	0
4	A	7002	GOL	7	0
4	B	7021	GOL	2	0
4	A	7015	GOL	3	0
4	B	7022	GOL	4	0
4	A	7006	GOL	1	0
4	B	7033	GOL	8	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	254/265 (95%)	-1.56	0 100 100	16, 29, 48, 75	2 (0%)
1	B	255/265 (96%)	-1.56	0 100 100	18, 30, 49, 72	3 (1%)
2	F	11/12 (91%)	-1.36	0 100 100	46, 79, 88, 113	0
2	H	11/12 (91%)	-1.60	0 100 100	30, 38, 43, 99	0
3	D	11/12 (91%)	-1.48	0 100 100	40, 61, 87, 99	0
3	G	11/12 (91%)	-1.50	0 100 100	37, 41, 60, 133	0
All	All	553/578 (95%)	-1.55	0 100 100	16, 31, 56, 133	5 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GOL	B	7009	6/6	0.97	0.04	76,79,81,82	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GOL	A	7004	6/6	0.98	0.05	48,52,54,54	0
4	GOL	B	7032	6/6	0.98	0.05	67,68,71,71	0
4	GOL	A	7010	6/6	0.99	0.03	36,40,45,48	0
4	GOL	A	7012	6/6	0.99	0.04	37,38,41,41	0
4	GOL	A	7013	6/6	0.99	0.03	41,43,46,46	0
4	GOL	A	7015	6/6	0.99	0.04	32,42,45,45	0
4	GOL	A	7020	6/6	0.99	0.03	27,31,37,38	0
4	GOL	A	7024	6/6	0.99	0.03	57,58,59,59	0
4	GOL	A	7030	6/6	0.99	0.04	37,39,41,42	0
4	GOL	B	7003	6/6	0.99	0.05	65,72,74,79	0
4	GOL	A	7002	6/6	0.99	0.04	38,46,50,50	0
4	GOL	B	7011	6/6	0.99	0.04	28,34,38,41	0
4	GOL	B	7021	6/6	0.99	0.04	39,44,44,45	0
4	GOL	B	7022	6/6	0.99	0.04	35,41,44,48	0
4	GOL	B	7027	6/6	0.99	0.03	56,60,61,63	0
4	GOL	A	7006	6/6	0.99	0.03	32,45,46,51	0
4	GOL	B	7033	6/6	0.99	0.05	32,36,37,37	0
6	PO4	B	7204	5/5	0.99	0.04	86,88,90,90	0
6	PO4	B	7205	5/5	0.99	0.04	101,103,103,103	0
4	GOL	A	7031	6/6	1.00	0.06	21,34,35,39	0
5	NA	A	7401	1/1	1.00	0.02	30,30,30,30	0

6.5 Other polymers

There are no such residues in this entry.