



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 09:31 PM UTC

PDB ID : 4GIZ / pdb_00004giz
Title : Crystal structure of full-length human papillomavirus oncoprotein E6 in complex with LXXLL peptide of ubiquitin ligase E6AP at 2.55 Å resolution
Authors : McEwen, A.G.; Zanier, K.; Charbonnier, S.; Poussin, P.; Cura, V.; Vande Pol, S.; Trave, G.; Cavarelli, J.
Deposited on : 2012-08-09
Resolution : 2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION






The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1091 (2.54-2.54)
Clashscore	190562	1120 (2.54-2.54)
Ramachandran outliers	187476	1106 (2.54-2.54)
Sidechain outliers	187428	1106 (2.54-2.54)
RSRZ outliers	180081	1091 (2.54-2.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	382	
1	B	382	
2	C	142	
2	D	142	
3	E	6	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	6	 100%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 9036 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Maltose-binding periplasmic protein, UBIQUITIN LIGASE EA6P: chimeric protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	382	Total	C	N	O	S	0	1	0
			2949	1896	480	567	6			
1	B	382	Total	C	N	O	S	0	1	0
			2951	1896	480	569	6			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	83	ALA	ASP	engineered mutation	UNP P0AEX9
A	84	ALA	LYS	engineered mutation	UNP P0AEX9
A	240	ALA	LYS	engineered mutation	UNP P0AEX9
A	360	ALA	GLU	engineered mutation	UNP P0AEX9
A	363	ALA	LYS	engineered mutation	UNP P0AEX9
A	364	ALA	ASP	engineered mutation	UNP P0AEX9
B	83	ALA	ASP	engineered mutation	UNP P0AEX9
B	84	ALA	LYS	engineered mutation	UNP P0AEX9
B	240	ALA	LYS	engineered mutation	UNP P0AEX9
B	360	ALA	GLU	engineered mutation	UNP P0AEX9
B	363	ALA	LYS	engineered mutation	UNP P0AEX9
B	364	ALA	ASP	engineered mutation	UNP P0AEX9

- Molecule 2 is a protein called Protein E6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	142	Total	C	N	O	S	0	1	0
			1206	755	222	217	12			
2	D	142	Total	C	N	O	S	0	3	0
			1220	763	225	220	12			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	47	ARG	PHE	engineered mutation	UNP P03126
C	80	SER	CYS	engineered mutation	UNP P03126
C	97	SER	CYS	engineered mutation	UNP P03126
C	111	SER	CYS	engineered mutation	UNP P03126
C	140	SER	CYS	engineered mutation	UNP P03126
D	47	ARG	PHE	engineered mutation	UNP P03126
D	80	SER	CYS	engineered mutation	UNP P03126
D	97	SER	CYS	engineered mutation	UNP P03126
D	111	SER	CYS	engineered mutation	UNP P03126
D	140	SER	CYS	engineered mutation	UNP P03126

- Molecule 3 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
3	E	6	Total	C	O	0	0	0
			67	36	31			
3	F	6	Total	C	O	0	0	0
			67	36	31			

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	2	Total	Zn	0	0
			2	2		
4	D	2	Total	Zn	0	0
			2	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	236	Total	O	0	0
			236	236		
5	B	181	Total	O	0	0
			181	181		
5	C	79	Total	O	0	0
			79	79		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	D	76	Total	O	0	0
			76	76		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Maltose-binding periplasmic protein, UBIQUITIN LIGASE EA6P: chimeric protein

Chain A: 





- Molecule 1: Maltose-binding periplasmic protein, UBIQUITIN LIGASE EA6P: chimeric protein

Chain B: 





- Molecule 2: Protein E6

Chain C: 





- Molecule 2: Protein E6

Chain D: 



- Molecule 3: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain E: 



- Molecule 3: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain F:

100%

GLC1
GLC2
GLC3
GLC4
GLC5
GLC6

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	106.28Å 134.94Å 138.79Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.33 – 2.55 37.33 – 2.55	Depositor EDS
% Data completeness (in resolution range)	99.9 (37.33-2.55) 99.8 (37.33-2.55)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.09 (at 2.54Å)	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
R, R_{free}	0.166 , 0.196 0.166 , 0.197	Depositor DCC
R_{free} test set	3325 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	45.9	Xtriage
Anisotropy	0.578	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 44.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.026 for -h,l,k	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9036	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.79	1/3021 (0.0%)	1.27	7/4104 (0.2%)
1	B	0.78	0/3020	1.26	10/4103 (0.2%)
2	C	0.78	0/1232	1.18	2/1655 (0.1%)
2	D	0.78	0/1249	1.20	3/1678 (0.2%)
All	All	0.78	1/8522 (0.0%)	1.24	22/11540 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	133	ILE	CA-C	5.08	1.56	1.52

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	132	GLU	CA-C-N	7.73	125.11	120.24
1	A	132	GLU	C-N-CA	7.73	125.11	120.24
1	B	132	GLU	CA-C-N	6.03	125.15	120.33
1	B	132	GLU	C-N-CA	6.03	125.15	120.33
1	A	16	LYS	N-CA-C	-5.78	106.27	113.38
1	A	292	GLU	CB-CG-CD	5.78	122.43	112.60
1	B	16	LYS	N-CA-C	-5.68	106.39	113.38
1	B	243	TYR	CA-C-N	5.66	125.96	121.61
1	B	243	TYR	C-N-CA	5.66	125.96	121.61
1	A	243	TYR	CA-C-N	5.63	125.94	121.61
1	A	243	TYR	C-N-CA	5.63	125.94	121.61
1	B	178	ASP	CA-CB-CG	5.39	117.99	112.60
1	B	165	ASP	CA-CB-CG	5.22	117.83	112.60
2	C	2	PHE	CA-C-N	5.19	127.69	120.63
2	C	2	PHE	C-N-CA	5.19	127.69	120.63
1	B	297	ASP	CA-CB-CG	5.17	117.77	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	309	GLU	CB-CG-CD	5.17	121.39	112.60
2	D	2	PHE	CA-C-N	5.12	127.60	120.63
2	D	2	PHE	C-N-CA	5.12	127.60	120.63
1	A	31	ASP	CA-CB-CG	5.12	117.72	112.60
1	B	31	ASP	CA-CB-CG	5.07	117.67	112.60
2	D	142	SER	N-CA-C	5.03	118.27	110.17

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2949	0	2917	6	0
1	B	2951	0	2912	5	0
2	C	1206	0	1191	1	0
2	D	1220	0	1204	1	0
3	E	67	0	54	0	0
3	F	67	0	52	0	0
4	C	2	0	0	0	0
4	D	2	0	0	0	0
5	A	236	0	0	1	0
5	B	181	0	0	1	0
5	C	79	0	0	0	0
5	D	76	0	0	0	0
All	All	9036	0	8330	12	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (12) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:274:LYS:O	1:A:278:LYS:HG2	2.03	0.57
1:A:69:GLY:HA3	1:A:333:ASN:O	2.07	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:296:LYS:HE2	5:A:513:HOH:O	2.12	0.49
1:B:80:ILE:HD12	1:B:82:PRO:HD3	1.96	0.47
1:B:278:LYS:NZ	5:B:669:HOH:O	2.48	0.46
1:A:63:TRP:HB3	1:A:68:PHE:HE1	1.81	0.46
1:B:63:TRP:HB3	1:B:68:PHE:HE1	1.81	0.45
1:A:80:ILE:HD12	1:A:82:PRO:HD3	2.00	0.43
1:B:69:GLY:HA3	1:B:333:ASN:O	2.20	0.41
1:A:149:MET:HE2	1:A:213:ILE:HG22	2.02	0.41
1:B:149:MET:HE2	1:B:213:ILE:HG22	2.02	0.41
2:C:35:GLN:HG2	2:D:69:PHE:CE2	2.55	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	381/382 (100%)	369 (97%)	12 (3%)	0	100	100
1	B	381/382 (100%)	372 (98%)	9 (2%)	0	100	100
2	C	141/142 (99%)	136 (96%)	5 (4%)	0	100	100
2	D	143/142 (101%)	135 (94%)	8 (6%)	0	100	100
All	All	1046/1048 (100%)	1012 (97%)	34 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	300/299 (100%)	296 (99%)	4 (1%)	61	76
1	B	300/299 (100%)	296 (99%)	4 (1%)	61	76
2	C	137/136 (101%)	136 (99%)	1 (1%)	76	85
2	D	139/136 (102%)	137 (99%)	2 (1%)	59	75
All	All	876/870 (101%)	865 (99%)	11 (1%)	61	76

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	61	ILE
1	A	80	ILE
1	A	179	ILE
1	A	257	LYS
1	B	61	ILE
1	B	80	ILE
1	B	179	ILE
1	B	309	GLU
2	C	129	ARG
2	D	64	ASP
2	D	143	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	283	ASN
1	A	356	GLN
1	B	73	GLN
1	B	283	ASN
1	B	356	GLN
1	B	368	ASN
2	C	36	GLN
2	C	91	GLN
2	D	36	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

12 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	GLC	E	1	3	12,12,12	1.34	2 (16%)	17,17,17	1.86	6 (35%)
3	GLC	E	2	3	11,11,12	3.17	7 (63%)	15,15,17	1.67	4 (26%)
3	GLC	E	3	3	11,11,12	2.91	6 (54%)	15,15,17	1.44	2 (13%)
3	GLC	E	4	3	11,11,12	3.06	5 (45%)	15,15,17	1.62	2 (13%)
3	GLC	E	5	3	11,11,12	3.41	8 (72%)	15,15,17	2.44	4 (26%)
3	GLC	E	6	3	11,11,12	3.24	6 (54%)	15,15,17	2.64	6 (40%)
3	GLC	F	1	3	12,12,12	1.27	0	17,17,17	1.81	5 (29%)
3	GLC	F	2	3	11,11,12	3.28	6 (54%)	15,15,17	1.61	4 (26%)
3	GLC	F	3	3	11,11,12	2.91	5 (45%)	15,15,17	1.30	3 (20%)
3	GLC	F	4	3	11,11,12	3.10	5 (45%)	15,15,17	1.50	4 (26%)
3	GLC	F	5	3	11,11,12	3.49	9 (81%)	15,15,17	2.06	4 (26%)
3	GLC	F	6	3	11,11,12	3.26	6 (54%)	15,15,17	2.65	5 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GLC	E	1	3	-	0/2/22/22	0/1/1/1
3	GLC	E	2	3	-	0/2/19/22	0/1/1/1
3	GLC	E	3	3	-	0/2/19/22	0/1/1/1
3	GLC	E	4	3	-	0/2/19/22	0/1/1/1
3	GLC	E	5	3	-	2/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GLC	E	6	3	-	2/2/19/22	0/1/1/1
3	GLC	F	1	3	-	0/2/22/22	0/1/1/1
3	GLC	F	2	3	-	1/2/19/22	0/1/1/1
3	GLC	F	3	3	-	0/2/19/22	0/1/1/1
3	GLC	F	4	3	-	0/2/19/22	0/1/1/1
3	GLC	F	5	3	-	1/2/19/22	0/1/1/1
3	GLC	F	6	3	-	2/2/19/22	0/1/1/1

All (65) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	6	GLC	O5-C5	6.62	1.56	1.43
3	F	4	GLC	O5-C5	6.54	1.56	1.43
3	E	6	GLC	O5-C5	6.45	1.56	1.43
3	E	4	GLC	O5-C5	6.26	1.55	1.43
3	F	2	GLC	O5-C5	6.25	1.55	1.43
3	F	5	GLC	O5-C5	6.22	1.55	1.43
3	F	5	GLC	C6-C5	-5.92	1.32	1.51
3	E	5	GLC	C6-C5	-5.88	1.32	1.51
3	E	5	GLC	O5-C5	5.82	1.54	1.43
3	F	2	GLC	C2-C3	-5.75	1.43	1.52
3	E	2	GLC	C2-C3	-5.75	1.43	1.52
3	F	3	GLC	O5-C5	5.37	1.53	1.43
3	E	6	GLC	C2-C3	-5.30	1.44	1.52
3	E	2	GLC	O5-C5	5.27	1.53	1.43
3	F	6	GLC	C2-C3	-5.14	1.44	1.52
3	E	3	GLC	O5-C5	5.11	1.53	1.43
3	E	5	GLC	C2-C3	-5.01	1.44	1.52
3	F	5	GLC	C2-C3	-4.91	1.45	1.52
3	E	3	GLC	C2-C3	-4.75	1.45	1.52
3	F	3	GLC	C6-C5	-4.64	1.36	1.51
3	F	3	GLC	C2-C3	-4.56	1.45	1.52
3	F	2	GLC	C6-C5	-4.55	1.36	1.51
3	F	4	GLC	C2-C3	-4.52	1.45	1.52
3	E	2	GLC	C6-C5	-4.51	1.36	1.51
3	E	3	GLC	C6-C5	-4.50	1.36	1.51
3	E	4	GLC	C2-C3	-4.47	1.45	1.52
3	E	4	GLC	C6-C5	-4.45	1.36	1.51
3	F	4	GLC	C6-C5	-4.28	1.37	1.51
3	F	6	GLC	C6-C5	-4.25	1.37	1.51
3	E	6	GLC	C6-C5	-4.23	1.37	1.51
3	E	2	GLC	C1-C2	-3.26	1.44	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	3	GLC	C1-C2	-3.25	1.44	1.52
3	F	5	GLC	C4-C5	3.03	1.59	1.53
3	E	4	GLC	O2-C2	2.96	1.49	1.43
3	F	4	GLC	C1-C2	-2.79	1.45	1.52
3	F	2	GLC	C1-C2	-2.74	1.45	1.52
3	F	3	GLC	C1-C2	-2.73	1.45	1.52
3	E	6	GLC	O2-C2	2.68	1.48	1.43
3	F	5	GLC	O2-C2	2.67	1.48	1.43
3	F	2	GLC	O3-C3	2.65	1.49	1.43
3	E	5	GLC	O2-C2	2.62	1.48	1.43
3	F	6	GLC	O2-C2	2.58	1.48	1.43
3	F	4	GLC	O2-C2	2.53	1.48	1.43
3	E	5	GLC	C4-C5	2.53	1.58	1.53
3	E	6	GLC	C1-C2	-2.50	1.46	1.52
3	E	4	GLC	C1-C2	-2.50	1.46	1.52
3	F	6	GLC	O5-C1	2.50	1.47	1.43
3	E	5	GLC	C1-C2	-2.46	1.46	1.52
3	E	5	GLC	O3-C3	2.46	1.49	1.43
3	E	2	GLC	O3-C3	2.40	1.48	1.43
3	F	6	GLC	C1-C2	-2.39	1.46	1.52
3	F	3	GLC	O2-C2	2.36	1.48	1.43
3	E	6	GLC	O5-C1	2.35	1.47	1.43
3	E	1	GLC	O1-C1	-2.35	1.32	1.39
3	F	5	GLC	C1-C2	-2.32	1.46	1.52
3	E	3	GLC	C4-C3	-2.32	1.46	1.52
3	F	2	GLC	O2-C2	2.26	1.48	1.43
3	F	5	GLC	O3-C3	2.20	1.48	1.43
3	E	2	GLC	C4-C3	-2.15	1.46	1.52
3	E	1	GLC	O5-C1	2.14	1.48	1.42
3	E	2	GLC	O2-C2	2.13	1.47	1.43
3	F	5	GLC	O5-C1	2.11	1.47	1.43
3	E	5	GLC	C4-C3	-2.10	1.46	1.52
3	F	5	GLC	C4-C3	-2.10	1.46	1.52
3	E	3	GLC	O2-C2	2.07	1.47	1.43

All (49) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	6	GLC	C1-O5-C5	6.94	121.48	112.19
3	E	5	GLC	C1-O5-C5	6.83	121.34	112.19
3	E	6	GLC	C1-O5-C5	6.81	121.32	112.19
3	F	5	GLC	C1-O5-C5	5.82	119.98	112.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	6	GLC	C3-C4-C5	4.18	117.81	110.23
3	E	6	GLC	C3-C4-C5	4.14	117.74	110.23
3	E	6	GLC	C1-C2-C3	3.72	115.06	109.64
3	E	1	GLC	O5-C5-C4	3.71	116.39	109.70
3	E	1	GLC	O5-C1-C2	3.70	116.81	110.30
3	F	6	GLC	C1-C2-C3	3.67	114.99	109.64
3	E	5	GLC	C1-C2-C3	3.59	114.87	109.64
3	E	6	GLC	C2-C3-C4	3.57	117.14	110.86
3	E	4	GLC	C2-C3-C4	3.55	117.10	110.86
3	E	5	GLC	C3-C4-C5	3.51	116.59	110.23
3	E	3	GLC	C1-O5-C5	3.37	116.71	112.19
3	F	1	GLC	O5-C5-C4	3.35	115.73	109.70
3	F	4	GLC	C2-C3-C4	3.21	116.51	110.86
3	E	2	GLC	C1-O5-C5	3.14	116.39	112.19
3	F	1	GLC	C1-O5-C5	3.07	119.59	113.65
3	F	6	GLC	C2-C3-C4	3.04	116.20	110.86
3	F	1	GLC	C4-C3-C2	3.02	116.12	110.83
3	E	4	GLC	C1-O5-C5	2.98	116.18	112.19
3	E	1	GLC	C4-C3-C2	2.89	115.90	110.83
3	F	5	GLC	C1-C2-C3	2.88	113.83	109.64
3	E	5	GLC	C2-C3-C4	2.86	115.89	110.86
3	F	1	GLC	O5-C1-C2	2.85	115.31	110.30
3	E	2	GLC	C2-C3-C4	2.84	115.86	110.86
3	F	2	GLC	C2-C3-C4	2.84	115.85	110.86
3	F	2	GLC	C1-C2-C3	2.79	113.71	109.64
3	E	3	GLC	C2-C3-C4	2.69	115.60	110.86
3	F	4	GLC	C1-O5-C5	2.65	115.73	112.19
3	E	1	GLC	C1-C2-C3	2.60	115.65	110.36
3	E	1	GLC	C1-O5-C5	2.47	118.44	113.65
3	E	1	GLC	O4-C4-C3	-2.45	104.59	110.38
3	F	3	GLC	C1-O5-C5	2.41	115.42	112.19
3	E	2	GLC	C1-C2-C3	2.40	113.14	109.64
3	F	6	GLC	O6-C6-C5	2.37	119.40	111.33
3	F	5	GLC	C3-C4-C5	2.36	114.51	110.23
3	E	6	GLC	O6-C6-C5	2.33	119.28	111.33
3	F	5	GLC	C2-C3-C4	2.31	114.92	110.86
3	E	2	GLC	C3-C4-C5	2.30	114.40	110.23
3	F	2	GLC	C1-O5-C5	2.25	115.20	112.19
3	F	1	GLC	C1-C2-C3	2.21	114.87	110.36
3	E	6	GLC	O5-C1-C2	2.20	116.05	110.79
3	F	3	GLC	C2-C3-C4	2.11	114.58	110.86
3	F	4	GLC	C1-C2-C3	2.10	112.69	109.64

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	4	GLC	O6-C6-C5	2.02	118.22	111.33
3	F	3	GLC	O2-C2-C3	2.01	114.31	110.15
3	F	2	GLC	O5-C5-C6	2.01	111.57	107.66

There are no chirality outliers.


All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	E	6	GLC	O5-C5-C6-O6
3	E	6	GLC	C4-C5-C6-O6
3	F	6	GLC	C4-C5-C6-O6
3	F	6	GLC	O5-C5-C6-O6
3	E	5	GLC	C4-C5-C6-O6
3	F	2	GLC	C4-C5-C6-O6
3	F	5	GLC	C4-C5-C6-O6
3	E	5	GLC	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	382/382 (100%)	-0.44	0 100 100	26, 46, 71, 101	1 (0%)
1	B	382/382 (100%)	-0.29	1 (0%) 90 92	27, 51, 76, 107	1 (0%)
2	C	142/142 (100%)	-0.28	6 (4%) 40 41	27, 47, 89, 117	1 (0%)
2	D	142/142 (100%)	-0.22	7 (4%) 35 34	23, 48, 89, 116	3 (2%)
All	All	1048/1048 (100%)	-0.33	14 (1%) 75 75	23, 48, 80, 117	6 (0%)

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	143	SER	4.0
2	D	143	SER	4.0
2	C	142	SER	3.2
2	C	92	TYR	2.9
2	D	5	PRO	2.7
2	D	141	ARG	2.5
2	C	137[A]	MET	2.4
2	C	3	GLN	2.3
2	D	92	TYR	2.3
2	D	3	GLN	2.3
1	B	176	LYS	2.2
2	D	2	PHE	2.2
2	C	91	GLN	2.2
2	D	129	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GLC	F	6	11/12	0.66	0.19	93,98,99,99	0
3	GLC	E	6	11/12	0.74	0.17	95,99,102,102	0
3	GLC	F	5	11/12	0.93	0.08	69,74,78,85	0
3	GLC	E	5	11/12	0.94	0.09	66,71,75,85	0
3	GLC	E	1	12/12	0.95	0.08	45,48,57,60	0
3	GLC	F	1	12/12	0.97	0.06	47,52,61,63	0
3	GLC	F	4	11/12	0.97	0.08	43,51,54,60	0
3	GLC	E	2	11/12	0.97	0.06	35,37,39,40	0
3	GLC	E	4	11/12	0.97	0.06	44,46,49,57	0
3	GLC	F	3	11/12	0.98	0.05	38,43,46,49	0
3	GLC	F	2	11/12	0.98	0.04	39,43,45,46	0
3	GLC	E	3	11/12	0.99	0.05	36,38,39,43	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain E:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	ZN	C	201	1/1	1.00	0.01	40,40,40,40	0
4	ZN	C	202	1/1	1.00	0.02	36,36,36,36	0
4	ZN	D	201	1/1	1.00	0.01	46,46,46,46	0
4	ZN	D	202	1/1	1.00	0.02	38,38,38,38	0

6.5 Other polymers [i](#)

There are no such residues in this entry.