



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 03:53 AM UTC

PDB ID : 8GLE / pdb_00008gle
Title : Crystal Structure of Human CD1b in Complex with Lysosulfatide
Authors : Shahine, A.
Deposited on : 2023-03-22
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	300	% 83% 10% 7%
2	B	101	4% 85% 12% ..
3	C	7	14% 86%
4	D	3	33% 33% 33%

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 3536 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called T-cell surface glycoprotein CD1b.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	280	Total	C	N	O	S	0	6	0
			2218	1422	377	409	10			

There are 23 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	279	GLY	-	expression tag	UNP P29016
A	280	SER	-	expression tag	UNP P29016
A	281	GLY	-	expression tag	UNP P29016
A	282	LEU	-	expression tag	UNP P29016
A	283	ASN	-	expression tag	UNP P29016
A	284	ASP	-	expression tag	UNP P29016
A	285	ILE	-	expression tag	UNP P29016
A	286	PHE	-	expression tag	UNP P29016
A	287	GLU	-	expression tag	UNP P29016
A	288	ALA	-	expression tag	UNP P29016
A	289	GLN	-	expression tag	UNP P29016
A	290	LYS	-	expression tag	UNP P29016
A	291	ILE	-	expression tag	UNP P29016
A	292	GLU	-	expression tag	UNP P29016
A	293	TRP	-	expression tag	UNP P29016
A	294	HIS	-	expression tag	UNP P29016
A	295	GLU	-	expression tag	UNP P29016
A	296	HIS	-	expression tag	UNP P29016
A	297	HIS	-	expression tag	UNP P29016
A	298	HIS	-	expression tag	UNP P29016
A	299	HIS	-	expression tag	UNP P29016
A	300	HIS	-	expression tag	UNP P29016
A	301	HIS	-	expression tag	UNP P29016


- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	99	Total	C	N	O	S	0	2	0
			823	525	141	155	2			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	1	PRO	-	expression tag	UNP P61769
B	2	LYS	-	expression tag	UNP P61769

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-3)][alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	7	Total	C	N	O	0	0	0
			81	46	2	33			

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	D	3	Total	C	N	O	0	0	0
			38	22	2	14			

- Molecule 5 is (2S,3R,4E)-2-amino-3-hydroxyoctadec-4-en-1-yl 3-O-sulfo-beta-D-galactopyranoside (CCD ID: SGF) (formula: C₂₄H₄₇NO₁₀S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	S	3	0
			36	24	1	10	1		

- Molecule 6 is tetracosyl octadecanoate (CCD ID: CUY) (formula: C₄₂H₈₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			33	31	2		

- Molecule 7 is IODIDE ION (CCD ID: IOD) (formula: I).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total I 1 1	0	0

- Molecule 8 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total C O 4 2 2	0	0
8	A	1	Total C O 4 2 2	0	0
8	A	1	Total C O 4 2 2	0	0
8	B	1	Total C O 4 2 2	0	0

- Molecule 9 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	2	Total Cl 2 2	0	0
9	B	1	Total Cl 1 1	0	0


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	225	Total 225	O 225	0	0
10	B	62	Total 62	O 62	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: T-cell surface glycoprotein CD1b




- Molecule 2: Beta-2-microglobulin



- Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-3)][alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	57.87Å 79.53Å 93.17Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.16 – 1.85 49.16 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.7 (49.16-1.85) 99.6 (49.16-1.85)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.31 (at 1.86Å)	Xtriage
Refinement program	PHENIX 1.14_3260	Depositor
R, R_{free}	0.188 , 0.229 0.191 , 0.231	Depositor DCC
R_{free} test set	1905 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	26.2	Xtriage
Anisotropy	0.375	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 46.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3536	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, CUY, IOD, BMA, SGF, FUC, EDO, CL, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.31	0/2295	0.54	0/3115
2	B	0.28	0/846	0.47	0/1148
All	All	0.30	0/3141	0.53	0/4263

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2218	0	2143	24	0
2	B	823	0	773	11	0
3	C	81	0	70	1	0
4	D	38	0	34	1	0
5	A	36	0	47	4	0
6	A	33	0	0	0	0
7	A	1	0	0	0	0
8	A	12	0	18	2	0
8	B	4	0	6	1	0
9	A	2	0	0	0	0
9	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
10	A	225	0	0	4	0
10	B	62	0	0	2	0
All	All	3536	0	3091	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:43:LYS:O	2:B:46:GLU:HG2	1.59	1.03
1:A:240:ASN:ND2	10:A:501:HOH:O	2.23	0.71
2:B:19:ASN:ND2	10:B:402:HOH:O	2.30	0.65
8:A:405:EDO:H12	2:B:34:PRO:HG3	1.82	0.61
1:A:264:LYS:NZ	1:A:272:ASP:OD2	2.34	0.59
1:A:6:GLY:HA2	8:A:406:EDO:H11	1.89	0.54
1:A:187:PRO:HB3	1:A:211:PHE:HB3	1.90	0.53
1:A:120:GLY:O	2:B:5:ARG:NH2	2.39	0.53
1:A:115[B]:ARG:HG2	1:A:115[B]:ARG:HH11	1.75	0.52
1:A:217:TRP:HB3	1:A:264:LYS:HB2	1.91	0.52
1:A:246:ARG:HD2	10:A:536:HOH:O	2.09	0.51
1:A:63:VAL:HG13	5:A:401:SGF:H1	1.92	0.51
1:A:65:GLU:HG3	10:A:589:HOH:O	2.10	0.51
1:A:171:LEU:HD12	3:C:2:NAG:H82	1.95	0.49
2:B:91:GLN:H	8:B:301:EDO:H11	1.77	0.49
1:A:66:LEU:HD13	5:A:401:SGF:H9	1.96	0.48
1:A:157:THR:HG21	5:A:401:SGF:O3	2.15	0.47
2:B:99:ARG:H	2:B:99:ARG:HG3	1.40	0.47
1:A:31:TRP:CZ3	1:A:36:GLN:HB2	2.51	0.46
2:B:14:ARG:NH2	10:B:405:HOH:O	2.43	0.45
1:A:117:ALA:HB2	2:B:62:TRP:CE2	2.52	0.44
1:A:58:PHE:CZ	1:A:169:TYR:HB2	2.52	0.44
1:A:190:TRP:CZ3	2:B:16:PRO:HD3	2.53	0.44
1:A:178:LYS:NZ	10:A:518:HOH:O	2.42	0.44
1:A:19:THR:HG21	4:D:3:FUC:H5	2.01	0.43
1:A:76:GLY:O	1:A:80:GLU:HG2	2.19	0.42
1:A:263:VAL:HB	1:A:273:ILE:HB	2.00	0.42
1:A:73:TYR:HA	5:A:401:SGF:H38	2.02	0.41
2:B:14:ARG:HB3	2:B:24:PHE:HB2	2.03	0.41
1:A:78:ALA:O	1:A:82:GLN:HG2	2.21	0.41
1:A:219:MET:HE2	1:A:219:MET:HB2	1.94	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:22:SER:HA	2:B:73:THR:HG22	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	284/300 (95%)	278 (98%)	6 (2%)	0	100	100
2	B	99/101 (98%)	98 (99%)	1 (1%)	0	100	100
All	All	383/401 (96%)	376 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	235/247 (95%)	235 (100%)	0	100	100
2	B	91/96 (95%)	90 (99%)	1 (1%)	65	57
All	All	326/343 (95%)	325 (100%)	1 (0%)	86	85

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	99	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	5	GLN
1	A	142	GLN
1	A	150	GLN
1	A	184	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

10 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	C	1	1,3	14,14,15	0.55	0	17,19,21	0.92	1 (5%)
3	NAG	C	2	3	14,14,15	0.38	0	17,19,21	0.64	0
3	BMA	C	3	3	11,11,12	0.56	0	15,15,17	0.83	1 (6%)
3	MAN	C	4	3	11,11,12	0.85	0	15,15,17	0.87	1 (6%)
3	MAN	C	5	3	11,11,12	0.70	0	15,15,17	1.09	1 (6%)
3	FUC	C	6	3	10,10,11	1.18	1 (10%)	14,14,16	0.90	0
3	FUC	C	7	3	10,10,11	1.07	0	14,14,16	0.72	0
4	NAG	D	1	4,1	14,14,15	0.37	0	17,19,21	0.45	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	D	2	4	14,14,15	0.48	0	17,19,21	1.00	1 (5%)
4	FUC	D	3	4	10,10,11	0.82	0	14,14,16	1.05	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	C	2	3	-	0/6/23/26	0/1/1/1
3	BMA	C	3	3	-	0/2/19/22	0/1/1/1
3	MAN	C	4	3	-	0/2/19/22	0/1/1/1
3	MAN	C	5	3	-	0/2/19/22	0/1/1/1
3	FUC	C	6	3	-	-	0/1/1/1
3	FUC	C	7	3	-	-	0/1/1/1
4	NAG	D	1	4,1	-	2/6/23/26	0/1/1/1
4	NAG	D	2	4	-	0/6/23/26	0/1/1/1
4	FUC	D	3	4	-	-	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	6	FUC	C4-C5	2.31	1.57	1.52

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	5	MAN	C1-O5-C5	3.34	116.67	112.19
4	D	2	NAG	C1-O5-C5	2.94	116.13	112.19
3	C	1	NAG	C1-O5-C5	2.55	115.61	112.19
3	C	3	BMA	C1-O5-C5	2.27	115.22	112.19
3	C	4	MAN	O2-C2-C3	-2.21	105.57	110.15
4	D	3	FUC	O5-C5-C4	2.10	113.33	109.55

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	1	NAG	C3-C2-N2-C7

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	D	1	NAG	C4-C5-C6-O6
3	C	1	NAG	C1-C2-N2-C7
4	D	1	NAG	O5-C5-C6-O6


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	2	NAG	1	0
4	D	3	FUC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 4 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
8	EDO	A	404	-	3,3,3	0.37	0	2,2,2	0.49	0
8	EDO	A	405	-	3,3,3	0.25	0	2,2,2	0.29	0
5	SGF	A	401	-	35,36,36	1.24	4 (11%)	40,46,46	1.87	9 (22%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	EDO	B	301	-	3,3,3	0.48	0	2,2,2	0.45	0
6	CUY	A	402	-	32,32,43	0.65	1 (3%)	32,32,43	1.06	1 (3%)
8	EDO	A	406	-	3,3,3	0.59	0	2,2,2	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	EDO	A	404	-	-	1/1/1/1	-
8	EDO	A	405	-	-	1/1/1/1	-
5	SGF	A	401	-	-	21/31/51/51	0/1/1/1
8	EDO	B	301	-	-	1/1/1/1	-
6	CUY	A	402	-	-	14/31/31/42	-
8	EDO	A	406	-	-	1/1/1/1	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	401	SGF	O4-S	3.03	1.66	1.57
5	A	401	SGF	O6-C43	2.64	1.48	1.41
6	A	402	CUY	OAQ-CAP	2.51	1.40	1.33
5	A	401	SGF	C44-C45	-2.39	1.45	1.52
5	A	401	SGF	O1-C2	-2.32	1.39	1.43

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	401	SGF	C43-C44-C45	-6.87	98.16	110.11
5	A	401	SGF	O-C43-C44	3.13	113.03	108.27
5	A	401	SGF	O6-C47-C46	3.12	115.33	109.70
5	A	401	SGF	C46-C45-C44	-3.01	106.68	110.86
5	A	401	SGF	O8-S-O4	2.85	112.93	106.37
5	A	401	SGF	O9-S-O10	-2.76	101.61	112.24
5	A	401	SGF	C2-C3-C4	-2.68	119.14	124.69
6	A	402	CUY	OAQ-CAP-CAO	2.60	119.76	111.83
5	A	401	SGF	O4-C45-C46	2.24	118.02	107.12
5	A	401	SGF	C45-O4-S	2.14	124.17	119.04

There are no chirality outliers.

All (39) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	A	401	SGF	C-C1-C2-C3
5	A	401	SGF	O-C-C1-C2
5	A	401	SGF	O-C-C1-N
5	A	401	SGF	C44-C45-O4-S
5	A	401	SGF	C46-C45-O4-S
5	A	401	SGF	C45-O4-S-O9
6	A	402	CUY	CAO-CAP-OAQ-CAR
6	A	402	CUY	OBO-CAP-OAQ-CAR
5	A	401	SGF	O6-C43-O-C
5	A	401	SGF	C44-C43-O-C
5	A	401	SGF	C10-C11-C12-C13
5	A	401	SGF	C7-C8-C9-C10
6	A	402	CUY	CAI-CAJ-CAK-CAL
5	A	401	SGF	C9-C10-C11-C12
5	A	401	SGF	C4-C5-C6-C7
8	A	404	EDO	O1-C1-C2-O2
8	A	406	EDO	O1-C1-C2-O2
5	A	401	SGF	C45-O4-S-O8
6	A	402	CUY	CBI-CBJ-CBK-CBL
6	A	402	CUY	CAS-CAT-CAU-CAV
5	A	401	SGF	C14-C15-C16-C17
6	A	402	CUY	CAR-CAS-CAT-CAU
6	A	402	CUY	CAV-CAW-CBM-CBL
6	A	402	CUY	CAF-CAG-CAH-CAI
6	A	402	CUY	CAM-CAN-CAO-CAP
8	A	405	EDO	O1-C1-C2-O2
5	A	401	SGF	C11-C12-C13-C14
5	A	401	SGF	N-C1-C2-C3
5	A	401	SGF	C-C1-C2-O1
5	A	401	SGF	N-C1-C2-O1
6	A	402	CUY	CAG-CAH-CAI-CAJ
5	A	401	SGF	C45-O4-S-O10
6	A	402	CUY	CBG-CBH-CBI-CBJ
5	A	401	SGF	C11-C10-C9-C8
6	A	402	CUY	CAU-CAV-CAW-CBM
5	A	401	SGF	C13-C14-C15-C16
6	A	402	CUY	CAL-CAM-CAN-CAO
8	B	301	EDO	O1-C1-C2-O2
6	A	402	CUY	CBJ-CBK-CBL-CBM

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	A	405	EDO	1	0
5	A	401	SGF	4	0
8	B	301	EDO	1	0
8	A	406	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	280/300 (93%)	-0.16	4 (1%) 73 77	13, 29, 53, 84	6 (2%)
2	B	99/101 (98%)	0.28	4 (4%) 42 46	10, 39, 73, 106	2 (2%)
All	All	379/401 (94%)	-0.04	8 (2%) 63 67	10, 31, 61, 106	8 (2%)

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	279	GLY	3.9
2	B	77	LYS	2.6
1	A	195	PRO	2.6
1	A	283	ASN	2.5
2	B	45	GLY	2.3
2	B	100	ASP	2.3
1	A	270	GLY	2.2
2	B	76	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	D	2	14/15	0.58	0.15	61,64,71,72	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	D	1	14/15	0.77	0.13	54,66,69,74	0
4	FUC	D	3	10/11	0.78	0.13	51,59,66,69	0
3	NAG	C	2	14/15	0.89	0.10	27,33,39,41	0
3	FUC	C	7	10/11	0.90	0.12	30,41,44,49	0
3	MAN	C	5	11/12	0.92	0.12	44,47,53,57	0
3	MAN	C	4	11/12	0.93	0.08	36,41,49,58	0
3	FUC	C	6	10/11	0.93	0.09	33,37,47,50	0
3	BMA	C	3	11/12	0.95	0.07	31,36,40,41	0
3	NAG	C	1	14/15	0.96	0.06	20,25,31,32	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	SGF	A	401	36/36	0.84	0.16	38,57,77,82	3
8	EDO	A	404	4/4	0.88	0.18	60,63,64,64	0
6	CUY	A	402	33/44	0.90	0.12	27,43,57,61	0
8	EDO	B	301	4/4	0.90	0.14	28,33,40,46	0
9	CL	B	302	1/1	0.91	0.23	34,34,34,34	0
8	EDO	A	405	4/4	0.93	0.10	23,31,33,33	0
8	EDO	A	406	4/4	0.94	0.21	22,26,39,47	0
9	CL	A	407	1/1	0.96	0.06	37,37,37,37	0

Continued on next page...




Continued from previous page...


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
9	CL	A	408	1/1	0.99	0.03	24,24,24,24	0
7	IOD	A	403	1/1	0.99	0.02	21,21,21,21	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around SGF A 401:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.