



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 10, 2026 – 01:43 AM UTC

PDB ID : 8HLE / pdb_00008hle
Title : Structure of DddY-DMSOP complex
Authors : Peng, M.; Li, C.Y.; Zhang, Y.Z.
Deposited on : 2022-11-30
Resolution : 1.91 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.91 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1188 (1.92-1.92)
Clashscore	190562	1209 (1.92-1.92)
Ramachandran outliers	187476	1195 (1.92-1.92)
Sidechain outliers	187428	1195 (1.92-1.92)
RSRZ outliers	180081	1188 (1.92-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	407	2% 86% 6% • 7%
1	B	407	2% 88% 5% 7%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7308 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DMSP lyase DddY.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	380	Total	C	N	O	S	0	1	0
			3069	1943	524	585	17			
1	B	380	Total	C	N	O	S	0	1	0
			3069	1943	524	585	17			

There are 14 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	260	ALA	TYR	engineered mutation	UNP N8X9V6
A	402	HIS	-	expression tag	UNP N8X9V6
A	403	HIS	-	expression tag	UNP N8X9V6
A	404	HIS	-	expression tag	UNP N8X9V6
A	405	HIS	-	expression tag	UNP N8X9V6
A	406	HIS	-	expression tag	UNP N8X9V6
A	407	HIS	-	expression tag	UNP N8X9V6
B	260	ALA	TYR	engineered mutation	UNP N8X9V6
B	402	HIS	-	expression tag	UNP N8X9V6
B	403	HIS	-	expression tag	UNP N8X9V6
B	404	HIS	-	expression tag	UNP N8X9V6
B	405	HIS	-	expression tag	UNP N8X9V6
B	406	HIS	-	expression tag	UNP N8X9V6
B	407	HIS	-	expression tag	UNP N8X9V6

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		

- Molecule 3 is 3-[dimethyl(oxidanyl)-\$l^{\{4\}}\$-sulfanyl]propanoic acid (CCD ID: LNI) (formula:

C₅H₁₁O₃S).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	S	0	0
			9	5	3	1		
3	B	1	Total	C	O	S	0	0
			9	5	3	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	561	Total	O	0	0
			561	561		
4	B	589	Total	O	0	0
			589	589		

- Molecule 1: DMSP lyase DddY



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	65.80Å 73.39Å 87.37Å 90.00° 91.14° 90.00°	Depositor
Resolution (Å)	36.72 – 1.91 36.72 – 1.91	Depositor EDS
% Data completeness (in resolution range)	97.2 (36.72-1.91) 97.3 (36.72-1.91)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.63 (at 1.91Å)	Xtriage
Refinement program	PHENIX (1.16_3549: ???)	Depositor
R, R_{free}	0.160 , 0.201 (Not available) , 0.212	Depositor DCC
R_{free} test set	3197 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	16.6	Xtriage
Anisotropy	0.450	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 52.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.039 for h,-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7308	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, LNI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	0/3153	0.58	5/4280 (0.1%)
1	B	0.33	0/3153	0.56	3/4280 (0.1%)
All	All	0.35	0/6306	0.57	8/8560 (0.1%)

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	260	ALA	CA-C-N	-6.29	114.18	120.21
1	A	260	ALA	C-N-CA	-6.29	114.18	120.21
1	B	36	THR	N-CA-C	-5.94	103.10	110.41
1	B	31	LYS	CA-C-N	-5.87	113.89	119.76
1	B	31	LYS	C-N-CA	-5.87	113.89	119.76

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3069	0	2958	28	0
1	B	3069	0	2958	21	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	9	0	0	2	0
3	B	9	0	0	0	0
4	A	561	0	0	3	0
4	B	589	0	0	2	0
All	All	7308	0	5916	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 49 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:350:ILE:C	1:A:351:LYS:HD3	2.05	0.81
1:A:374:CYS:O	1:A:386:MET:HE1	1.82	0.80
1:B:376:PRO:HB3	1:B:381:LYS:CG	2.12	0.80
1:B:376:PRO:HB3	1:B:381:LYS:HG2	1.65	0.77
1:B:381:LYS:NZ	1:B:390:ASP:OD1	2.20	0.75

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	379/407 (93%)	370 (98%)	8 (2%)	1 (0%)	36	26
1	B	379/407 (93%)	370 (98%)	9 (2%)	0	100	100
All	All	758/814 (93%)	740 (98%)	17 (2%)	1 (0%)	48	40

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	380	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	333/354 (94%)	328 (98%)	5 (2%)	57	47
1	B	333/354 (94%)	331 (99%)	2 (1%)	78	75
All	All	666/708 (94%)	659 (99%)	7 (1%)	65	60

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	383	ILE
1	A	386	MET
1	B	382	ASP
1	B	31	LYS
1	A	381	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	213	ASN
1	B	322	GLN
1	A	285	HIS
1	A	322	GLN
1	B	40	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	LNI	A	502	2	7,8,8	1.38	0	8,11,11	1.67	2 (25%)
3	LNI	B	502	2	7,8,8	1.54	1 (14%)	8,11,11	1.45	1 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	LNI	A	502	2	-	3/4/6/6	-
3	LNI	B	502	2	-	4/4/6/6	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	502	LNI	O03-S01	2.25	1.47	1.44

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	LNI	C05-S01-C04	3.07	111.11	106.32
3	A	502	LNI	O03-S01-C05	-2.08	108.97	111.94
3	B	502	LNI	C05-S01-C04	2.03	109.48	106.32

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	502	LNI	C02-C03-S01-O03
3	B	502	LNI	C01-C02-C03-S01
3	A	502	LNI	O01-C01-C02-C03
3	A	502	LNI	O02-C01-C02-C03
3	B	502	LNI	O02-C01-C02-C03

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	502	LNI	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	380/407 (93%)	-0.18	7 (1%) 67 74	9, 17, 34, 64	1 (0%)
1	B	380/407 (93%)	-0.30	8 (2%) 63 70	9, 15, 32, 59	1 (0%)
All	All	760/814 (93%)	-0.24	15 (1%) 65 71	9, 16, 33, 64	2 (0%)

The worst 5 of 15 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	386	MET	4.3
1	A	380	ALA	4.2
1	B	380	ALA	4.2
1	A	302	ASN	4.0
1	B	381	LYS	3.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	LNI	A	502	9/9	0.86	0.12	16,23,36,46	0
3	LNI	B	502	9/9	0.90	0.15	12,23,30,49	0
2	ZN	A	501	1/1	1.00	0.02	14,14,14,14	0
2	ZN	B	501	1/1	1.00	0.02	12,12,12,12	0

6.5 Other polymers [i](#)

There are no such residues in this entry.