



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 12:46 PM UTC

PDB ID : 6IYZ / pdb_00006iyz
Title : Structural basis for activity of TRIC counter-ion channels in calcium release
Authors : Wang, X.H.; Zeng, Y.; Qin, L.; Gao, F.; Su, M.; Chen, Y.H.
Deposited on : 2018-12-18
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	326	13% 65% . 31%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1821 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Trimeric intracellular cation channel type A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	225	1806	1206	292	292	16	0	1	0

There are 31 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	245	SER	CYS	engineered mutation	UNP Q5ZK43
A	297	ALA	-	expression tag	UNP Q5ZK43
A	298	ALA	-	expression tag	UNP Q5ZK43
A	299	ALA	-	expression tag	UNP Q5ZK43
A	300	GLU	-	expression tag	UNP Q5ZK43
A	301	ASN	-	expression tag	UNP Q5ZK43
A	302	LEU	-	expression tag	UNP Q5ZK43
A	303	TYR	-	expression tag	UNP Q5ZK43
A	304	PHE	-	expression tag	UNP Q5ZK43
A	305	GLN	-	expression tag	UNP Q5ZK43
A	306	GLY	-	expression tag	UNP Q5ZK43
A	307	LEU	-	expression tag	UNP Q5ZK43
A	308	GLU	-	expression tag	UNP Q5ZK43
A	309	ASP	-	expression tag	UNP Q5ZK43
A	310	TYR	-	expression tag	UNP Q5ZK43
A	311	LYS	-	expression tag	UNP Q5ZK43
A	312	ASP	-	expression tag	UNP Q5ZK43
A	313	ASP	-	expression tag	UNP Q5ZK43
A	314	ASP	-	expression tag	UNP Q5ZK43
A	315	ASP	-	expression tag	UNP Q5ZK43
A	316	LYS	-	expression tag	UNP Q5ZK43
A	317	HIS	-	expression tag	UNP Q5ZK43
A	318	HIS	-	expression tag	UNP Q5ZK43
A	319	HIS	-	expression tag	UNP Q5ZK43
A	320	HIS	-	expression tag	UNP Q5ZK43
A	321	HIS	-	expression tag	UNP Q5ZK43
A	322	HIS	-	expression tag	UNP Q5ZK43

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	323	HIS	-	expression tag	UNP Q5ZK43
A	324	HIS	-	expression tag	UNP Q5ZK43
A	325	HIS	-	expression tag	UNP Q5ZK43
A	326	HIS	-	expression tag	UNP Q5ZK43

- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total Cl 2 2	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	13	Total O 13 13	0	0

- Molecule 1: Trimeric intracellular cation channel type A



4 Data and refinement statistics

Property	Value	Source
Space group	F 41 3 2	Depositor
Cell constants a, b, c, α , β , γ	268.21 Å 268.21 Å 268.21 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.41 – 2.20 47.41 – 2.20	Depositor EDS
% Data completeness (in resolution range)	100.0 (47.41-2.20) 100.0 (47.41-2.20)	Depositor EDS
R_{merge}	0.39	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.11 (at 2.20 Å)	Xtriage
Refinement program	PHENIX 1.14rc2_3191	Depositor
R, R_{free}	0.251 , 0.274 0.255 , 0.275	Depositor DCC
R_{free} test set	2199 reflections (5.19%)	wwPDB-VP
Wilson B-factor (Å ²)	54.7	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 34.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	1821	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.08	0/1870	0.22	0/2545

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1806	0	1791	7	0
2	A	2	0	0	0	0
3	A	13	0	0	0	0
All	All	1821	0	1791	7	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (7) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:129:LYS:HB3	1:A:155:LYS:HE2	1.93	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:68:LEU:HD13	1:A:162:MET:HE3	1.94	0.49
1:A:145:TRP:CZ3	1:A:202:THR:HG22	2.48	0.48
1:A:29:TYR:CD2	1:A:122:LYS:HG3	2.48	0.48
1:A:65:SER:HB2	1:A:155:LYS:HA	1.99	0.45
1:A:198:THR:O	1:A:202:THR:HG23	2.18	0.43
1:A:104:LEU:HG	1:A:108:CYS:SG	2.59	0.42

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	224/326 (69%)	222 (99%)	2 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	192/278 (69%)	190 (99%)	2 (1%)	68	81

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	86	SER
1	A	142	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	225/326 (69%)	1.28	41 (18%) 3 2	26, 56, 88, 132	1 (0%)

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	234	ALA	13.0
1	A	204	TRP	6.7
1	A	233	HIS	6.6
1	A	206	PRO	6.2
1	A	10	GLY	6.1
1	A	11	GLU	5.8
1	A	39	TYR	5.6
1	A	232	SER	5.1
1	A	135[A]	HIS	5.0
1	A	12	LEU	5.0
1	A	230	THR	4.3
1	A	207	VAL	4.2
1	A	102	MET	4.2
1	A	231	HIS	3.9
1	A	36	TYR	3.9
1	A	211	ASN	3.3
1	A	142	HIS	3.3
1	A	208	SER	3.2
1	A	108	CYS	2.9
1	A	70	ASP	2.9
1	A	68	LEU	2.8
1	A	215	PHE	2.8
1	A	192	TYR	2.8
1	A	45	GLU	2.8
1	A	65	SER	2.7
1	A	212	LEU	2.5
1	A	48	ARG	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	16	PHE	2.4
1	A	226	PHE	2.4
1	A	143	HIS	2.4
1	A	17	ALA	2.3
1	A	165	PHE	2.3
1	A	19	VAL	2.3
1	A	24	LEU	2.2
1	A	14	ALA	2.2
1	A	210	ALA	2.1
1	A	203	HIS	2.1
1	A	202	THR	2.1
1	A	71	LEU	2.0
1	A	96	LEU	2.0
1	A	104	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CL	A	402	1/1	0.87	0.64	104,104,104,104	0
2	CL	A	401	1/1	0.94	0.37	65,65,65,65	1

6.5 Other polymers [i](#)

There are no such residues in this entry.