



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 04:17 PM UTC

PDB ID : 5JC1 / pdb_00005jc1
Title : Structure of Plasmodium falciparum DXR in complex with a beta-substituted fosmidomycin analogue, LC55 and manganese
Authors : Sooriyaarachchi, S.; Bergfors, T.; Jones, T.A.; Mowbray, S.L.
Deposited on : 2016-04-14
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.66-1.66)
Clashscore	190562	2662 (1.66-1.66)
Ramachandran outliers	187476	2621 (1.66-1.66)
Sidechain outliers	187428	2621 (1.66-1.66)
RSRZ outliers	180081	2564 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	422	7% 94%
1	B	422	5% 96%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7236 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 1-deoxy-D-xylulose 5-phosphate reductoisomerase, apicoplastic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	410	Total	C	N	O	S	0	6	0
			3317	2126	543	626	22			
1	B	410	Total	C	N	O	S	0	5	0
			3305	2120	541	624	20			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	67	MET	-	initiating methionine	UNP Q8IKG4
A	68	ALA	-	expression tag	UNP Q8IKG4
A	69	HIS	-	expression tag	UNP Q8IKG4
A	70	HIS	-	expression tag	UNP Q8IKG4
A	71	HIS	-	expression tag	UNP Q8IKG4
A	72	HIS	-	expression tag	UNP Q8IKG4
A	73	HIS	-	expression tag	UNP Q8IKG4
A	74	HIS	-	expression tag	UNP Q8IKG4
B	67	MET	-	initiating methionine	UNP Q8IKG4
B	68	ALA	-	expression tag	UNP Q8IKG4
B	69	HIS	-	expression tag	UNP Q8IKG4
B	70	HIS	-	expression tag	UNP Q8IKG4
B	71	HIS	-	expression tag	UNP Q8IKG4
B	72	HIS	-	expression tag	UNP Q8IKG4
B	73	HIS	-	expression tag	UNP Q8IKG4
B	74	HIS	-	expression tag	UNP Q8IKG4

- Molecule 2 is [(2R)-2-{2-[hydroxy(methyl)amino]-2-oxoethyl}-5-(3-methoxyphenyl)pentyl]p
hosphonic acid (CCD ID: 6JB) (formula: C₁₅H₂₄NO₆P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			23	15	1	6	1		
2	B	1	Total	C	N	O	P	0	0
			23	15	1	6	1		

- Molecule 3 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mn	0	0
			1	1		
3	B	1	Total	Mn	0	0
			1	1		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	241	Total O 241 241	0	0
6	B	241	Total O 241 241	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	51.58Å 56.46Å 85.71Å 103.84° 103.25° 100.15°	Depositor
Resolution (Å)	30.00 – 1.65 30.00 – 1.65	Depositor EDS
% Data completeness (in resolution range)	97.4 (30.00-1.65) 97.4 (30.00-1.65)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.68 (at 1.65Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.179 , 0.204 0.189 , 0.210	Depositor DCC
R_{free} test set	5194 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å ²)	16.7	Xtriage
Anisotropy	0.483	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 35.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.012 for -h,-k,h+k+l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7236	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 6JB, MN, GOL, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.57	0/3380	0.78	2/4564 (0.0%)
1	B	0.57	1/3374 (0.0%)	0.80	0/4557
All	All	0.57	1/6754 (0.0%)	0.79	2/9121 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	362	ILE	CA-CB	5.22	1.57	1.54

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	118	VAL	N-CA-C	5.37	116.09	110.62
1	A	275	GLN	N-CA-C	5.05	117.51	111.71

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3317	0	3353	3	0
1	B	3305	0	3348	0	0
2	A	23	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	23	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	6	0	8	0	0
4	B	6	0	8	0	0
5	A	28	0	42	0	0
5	B	44	0	66	0	0
6	A	241	0	0	1	0
6	B	241	0	0	0	0
All	All	7236	0	6825	3	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All (3) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:150:ASN:ND2	6:A:604:HOH:O	2.49	0.42
1:A:217:LYS:HE2	1:A:458:GLU:O	2.20	0.41
1:A:340:ILE:HD12	1:A:391:PHE:HZ	1.85	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	414/422 (98%)	402 (97%)	12 (3%)	0	100	100
1	B	413/422 (98%)	402 (97%)	11 (3%)	0	100	100
All	All	827/844 (98%)	804 (97%)	23 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	383/388 (99%)	377 (98%)	6 (2%)	55	34
1	B	382/388 (98%)	378 (99%)	4 (1%)	68	52
All	All	765/776 (99%)	755 (99%)	10 (1%)	61	42

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	116	LYS
1	A	118	VAL
1	A	149	LYS
1	A	252	LEU
1	A	292	LYS
1	A	297	LYS
1	B	142	GLU
1	B	155	LYS
1	B	297	LYS
1	B	349	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	150	ASN
1	A	220	ASN
1	A	244	ASN
1	A	255	ASN
1	A	446	GLN
1	B	92	ASN
1	B	124	GLN
1	B	289	ASN
1	B	428	ASN
1	B	446	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 24 ligands modelled in this entry, 2 are monoatomic - leaving 22 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	EDO	B	501	-	3,3,3	0.36	0	2,2,2	0.93	0
5	EDO	A	506	-	3,3,3	0.38	0	2,2,2	0.36	0
4	GOL	A	503	-	5,5,5	0.39	0	5,5,5	0.56	0
5	EDO	B	512	-	3,3,3	0.54	0	2,2,2	0.49	0
5	EDO	B	510	-	3,3,3	0.42	0	2,2,2	0.15	0
5	EDO	A	508	-	3,3,3	0.45	0	2,2,2	0.30	0
5	EDO	B	513	-	3,3,3	0.43	0	2,2,2	0.40	0
5	EDO	A	510	-	3,3,3	0.47	0	2,2,2	0.35	0
5	EDO	A	504	-	3,3,3	0.42	0	2,2,2	0.38	0
5	EDO	A	509	-	3,3,3	0.41	0	2,2,2	0.34	0
5	EDO	A	507	-	3,3,3	0.46	0	2,2,2	0.31	0
5	EDO	A	505	-	3,3,3	0.47	0	2,2,2	0.43	0
5	EDO	B	507	-	3,3,3	0.36	0	2,2,2	0.53	0
4	GOL	B	504	-	5,5,5	0.35	0	5,5,5	0.66	0
5	EDO	B	508	-	3,3,3	0.44	0	2,2,2	0.34	0
5	EDO	B	506	-	3,3,3	0.46	0	2,2,2	0.18	0
2	6JB	B	502	3	20,23,23	1.11	2 (10%)	24,31,31	1.03	2 (8%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	EDO	B	509	-	3,3,3	0.44	0	2,2,2	0.34	0
5	EDO	B	505	-	3,3,3	0.34	0	2,2,2	0.61	0
5	EDO	B	511	-	3,3,3	0.41	0	2,2,2	0.29	0
2	6JB	A	501	3	20,23,23	1.12	1 (5%)	24,31,31	1.13	2 (8%)
5	EDO	B	514	-	3,3,3	0.38	0	2,2,2	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	B	501	-	-	1/1/1/1	-
5	EDO	A	506	-	-	1/1/1/1	-
4	GOL	A	503	-	-	0/4/4/4	-
5	EDO	B	512	-	-	0/1/1/1	-
5	EDO	B	510	-	-	0/1/1/1	-
5	EDO	A	508	-	-	0/1/1/1	-
5	EDO	B	513	-	-	1/1/1/1	-
5	EDO	A	510	-	-	0/1/1/1	-
5	EDO	A	504	-	-	1/1/1/1	-
5	EDO	A	509	-	-	1/1/1/1	-
5	EDO	A	507	-	-	1/1/1/1	-
5	EDO	A	505	-	-	0/1/1/1	-
5	EDO	B	507	-	-	0/1/1/1	-
4	GOL	B	504	-	-	0/4/4/4	-
5	EDO	B	508	-	-	0/1/1/1	-
5	EDO	B	506	-	-	1/1/1/1	-
2	6JB	B	502	3	-	7/21/21/21	0/1/1/1
5	EDO	B	509	-	-	1/1/1/1	-
5	EDO	B	505	-	-	1/1/1/1	-
5	EDO	B	511	-	-	1/1/1/1	-
2	6JB	A	501	3	-	6/21/21/21	0/1/1/1
5	EDO	B	514	-	-	0/1/1/1	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	6JB	P5-C1	3.74	1.82	1.79
2	B	502	6JB	P5-C1	3.34	1.82	1.79
2	B	502	6JB	P5-O7	-2.29	1.49	1.55

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	502	6JB	O12-C4-C3	-3.36	115.79	122.07
2	A	501	6JB	O12-C4-C3	-3.14	116.21	122.07
2	A	501	6JB	C23-O22-C19	3.12	124.19	117.50
2	B	502	6JB	C23-O22-C19	2.64	123.15	117.50

There are no chirality outliers.

All (23) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	B	505	EDO	O1-C1-C2-O2
5	A	509	EDO	O1-C1-C2-O2
5	B	511	EDO	O1-C1-C2-O2
5	A	504	EDO	O1-C1-C2-O2
5	B	501	EDO	O1-C1-C2-O2
5	B	509	EDO	O1-C1-C2-O2
2	A	501	6JB	C2-C3-C4-N9
2	B	502	6JB	C2-C3-C4-N9
2	A	501	6JB	C2-C3-C4-O12
2	B	502	6JB	C2-C3-C4-O12
2	A	501	6JB	P5-C1-C2-C3
2	B	502	6JB	P5-C1-C2-C3
2	B	502	6JB	C2-C1-P5-O8
2	B	502	6JB	C1-C2-C3-C4
5	A	506	EDO	O1-C1-C2-O2
5	B	513	EDO	O1-C1-C2-O2
5	A	507	EDO	O1-C1-C2-O2
5	B	506	EDO	O1-C1-C2-O2
2	A	501	6JB	C1-C2-C3-C4
2	A	501	6JB	O12-C4-N9-O10
2	B	502	6JB	O12-C4-N9-O10
2	A	501	6JB	P5-C1-C2-C13
2	B	502	6JB	P5-C1-C2-C13

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	410/422 (97%)	0.18	29 (7%)	22 25	6, 18, 46, 73	6 (1%)
1	B	410/422 (97%)	0.12	23 (5%)	30 34	6, 17, 44, 70	5 (1%)
All	All	820/844 (97%)	0.15	52 (6%)	26 29	6, 17, 45, 73	11 (1%)

All (52) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	296	TRP	5.9
1	A	184	PHE	5.5
1	B	184	PHE	4.8
1	B	118	VAL	4.3
1	B	296	TRP	4.3
1	B	141	TYR	4.0
1	A	295	LYS	3.6
1	B	188	TYR	3.5
1	B	294	PRO	3.4
1	A	294	PRO	3.4
1	A	293	HIS	3.4
1	A	140	VAL	3.4
1	B	297	LYS	3.3
1	B	485	HIS	3.3
1	B	293	HIS	3.2
1	A	141	TYR	3.1
1	A	485	HIS	3.1
1	A	77	PRO	3.0
1	B	486	ASN	2.9
1	A	192	TYR	2.8
1	B	140	VAL	2.8
1	A	188	TYR	2.8
1	B	295	LYS	2.8
1	A	486	ASN	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	153	ASP	2.7
1	A	160[A]	CYS	2.6
1	B	144	LEU	2.6
1	A	297	LYS	2.6
1	A	163	GLU	2.6
1	B	182	ASP	2.5
1	B	150	ASN	2.4
1	B	117	SER	2.4
1	A	183	SER	2.3
1	A	186	GLY	2.3
1	A	159	LEU	2.3
1	B	148	VAL	2.3
1	A	136	HIS	2.3
1	A	185	GLN	2.3
1	A	144	LEU	2.3
1	B	77	PRO	2.2
1	A	155	LYS	2.2
1	A	147	LEU	2.2
1	A	164	GLY	2.2
1	B	122	TYR	2.2
1	A	173	SER	2.2
1	B	160	CYS	2.1
1	A	181	ILE	2.1
1	A	116	LYS	2.1
1	A	118	VAL	2.1
1	B	147	LEU	2.1
1	B	138	LYS	2.1
1	A	139	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	EDO	B	510	4/4	0.76	0.16	44,45,45,46	0
4	GOL	A	503	6/6	0.77	0.14	29,31,32,33	0
5	EDO	B	505	4/4	0.81	0.14	25,26,27,27	0
5	EDO	A	509	4/4	0.81	0.17	48,48,49,50	0
5	EDO	A	504	4/4	0.82	0.15	28,28,29,29	0
5	EDO	B	506	4/4	0.83	0.17	38,38,38,39	0
4	GOL	B	504	6/6	0.83	0.12	28,29,29,29	0
5	EDO	B	511	4/4	0.83	0.14	38,39,39,40	0
5	EDO	A	505	4/4	0.84	0.13	24,27,27,29	0
5	EDO	B	513	4/4	0.84	0.14	32,35,36,38	0
5	EDO	A	507	4/4	0.85	0.14	38,39,40,42	0
5	EDO	A	510	4/4	0.85	0.15	40,40,41,43	0
5	EDO	B	509	4/4	0.85	0.15	33,35,37,38	0
5	EDO	B	501	4/4	0.86	0.16	22,25,25,27	0
5	EDO	B	507	4/4	0.86	0.14	21,25,27,30	0
5	EDO	B	514	4/4	0.86	0.13	30,31,32,33	0
5	EDO	B	508	4/4	0.87	0.14	26,30,31,35	0
5	EDO	A	506	4/4	0.87	0.13	40,40,41,43	0
5	EDO	A	508	4/4	0.92	0.10	44,44,45,45	0
5	EDO	B	512	4/4	0.96	0.06	15,15,15,16	0
2	6JB	A	501	23/23	0.97	0.06	14,16,19,21	0
2	6JB	B	502	23/23	0.98	0.06	13,15,18,20	0
3	MN	A	502	1/1	1.00	0.02	12,12,12,12	0
3	MN	B	503	1/1	1.00	0.03	12,12,12,12	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 6JB A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 6JB B 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.