



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 7, 2026 – 07:23 AM JST

PDB ID : 8JC3 / pdb_00008jc3
Title : Crystal structure of the P450 BM3 heme domain mutant F87A-T268V in complex with Pyd-N-C4-Phe and hydroxylamine
Authors : Jiang, Y.; Dong, S.; Feng, Y.; Cong, Z.
Deposited on : 2023-05-10
Resolution : 1.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.015 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50

i

X-RAY DIFFRACTION

A.

 R_{free}

1	A	465
---	---	-----

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 8281 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Bifunctional cytochrome P450/NADPH-P450 reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	451	Total	C	N	O	S	0	1	0
			3567	2282	606	662	17			
1	A	450	Total	C	N	O	S	0	1	0
			3562	2279	605	661	17			

There are 22 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	-1	GLY	-	expression tag	UNP P14779
B	87	ALA	PHE	engineered mutation	UNP P14779
B	268	VAL	THR	engineered mutation	UNP P14779
B	456	LEU	-	expression tag	UNP P14779
B	457	GLU	-	expression tag	UNP P14779
B	458	HIS	-	expression tag	UNP P14779
B	459	HIS	-	expression tag	UNP P14779
B	460	HIS	-	expression tag	UNP P14779
B	461	HIS	-	expression tag	UNP P14779
B	462	HIS	-	expression tag	UNP P14779
B	463	HIS	-	expression tag	UNP P14779
A	-1	GLY	-	expression tag	UNP P14779
A	87	ALA	PHE	engineered mutation	UNP P14779
A	268	VAL	THR	engineered mutation	UNP P14779
A	456	LEU	-	expression tag	UNP P14779
A	457	GLU	-	expression tag	UNP P14779
A	458	HIS	-	expression tag	UNP P14779
A	459	HIS	-	expression tag	UNP P14779
A	460	HIS	-	expression tag	UNP P14779
A	461	HIS	-	expression tag	UNP P14779
A	462	HIS	-	expression tag	UNP P14779
A	463	HIS	-	expression tag	UNP P14779

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is HYDROXYAMINE (CCD ID: HOA) (formula: H_3NO) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	N	O	0	0
			2	1	1		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	N	O	0	0
			2	1	1		

- Molecule 4 is 4-(pyridin-4-ylamino)butanoic acid (CCD ID: LXO) (formula: C₉H₁₂N₂O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	1
			12	9	2	1		
4	B	1	Total	C	N	O	0	1
			12	9	2	1		
4	A	1	Total	C	N	O	0	1
			12	9	2	1		
4	A	1	Total	C	N	O	0	1
			12	9	2	1		

- Molecule 5 is PHENYLALANINE (CCD ID: PHE) (formula: C₉H₁₁NO₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	N	O	0	1
			12	9	1	2		
5	B	1	Total	C	N	O	0	1
			12	9	1	2		
5	A	1	Total	C	N	O	0	1
			12	9	1	2		
5	A	1	Total	C	N	O	0	1
			12	9	1	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	491	Total	O	0	0
			491	491		
6	A	475	Total	O	0	0
			475	475		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Bifunctional cytochrome P450/NADPH-P450 reductase



- Molecule 1: Bifunctional cytochrome P450/NADPH-P450 reductase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	58.56Å 147.58Å 64.60Å 90.00° 100.21° 90.00°	Depositor
Resolution (Å)	45.42 – 1.82 45.42 – 1.82	Depositor EDS
% Data completeness (in resolution range)	97.4 (45.42-1.82) 97.4 (45.42-1.82)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.04 (at 1.82Å)	Xtriage
Refinement program	PHENIX (1.18.2_3874: ???)	Depositor
R, R_{free}	0.190 , 0.212 0.191 , 0.211	Depositor DCC
R_{free} test set	5000 reflections (5.20%)	wwPDB-VP
Wilson B-factor (Å ²)	22.0	Xtriage
Anisotropy	0.454	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 42.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8281	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HOA, LXO, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.23	2/3646 (0.1%)	0.40	0/4936
1	B	0.37	5/3651 (0.1%)	0.44	0/4943
All	All	0.31	7/7297 (0.1%)	0.42	0/9879

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	436	THR	C-O	-6.27	1.16	1.24
1	A	268	VAL	C-O	-6.23	1.16	1.24
1	B	149	THR	C-O	-5.90	1.17	1.24
1	B	288	GLN	C-O	-5.68	1.17	1.24
1	A	264	ALA	C-O	-5.30	1.17	1.24
1	B	285	HIS	C-O	-5.24	1.17	1.24
1	B	150	LEU	C-O	-5.21	1.18	1.24

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3562	0	3485	25	0
1	B	3567	0	3487	30	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	43	0	30	2	0
2	B	43	0	30	3	0
3	A	2	0	0	1	0
3	B	2	0	0	1	0
4	A	24	0	0	0	0
4	B	24	0	0	0	0
5	A	24	0	16	0	0
5	B	24	0	16	0	0
6	A	475	0	0	4	0
6	B	491	0	0	4	0
All	All	8281	0	7064	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:383:SER:HB2	1:A:104:LEU:HD21	1.51	0.89
1:A:264:ALA:HB1	3:A:502:HOA:N	1.90	0.85
1:A:118:MET:SD	6:A:780:HOH:O	2.44	0.74
1:B:388:HIS:HA	1:B:391:LYS:HD3	1.70	0.74
1:B:264:ALA:HB1	3:B:502:HOA:N	2.06	0.70
1:B:118:MET:SD	6:B:635:HOH:O	2.49	0.70
1:A:194:ASP:N	6:A:606:HOH:O	2.29	0.66
1:A:158:PHE:HD2	1:A:219:ILE:HG21	1.61	0.66
1:B:43:GLU:HG2	1:B:48:VAL:HG22	1.80	0.64
1:A:43:GLU:HG2	1:A:48:VAL:HG22	1.82	0.61
1:B:158:PHE:HD2	1:B:219:ILE:HG21	1.69	0.58
1:B:153:ILE:HG23	1:B:262:LEU:HD23	1.87	0.56
1:A:148:LEU:HD21	1:A:413:VAL:HG21	1.88	0.55
1:B:387:GLN:HG2	1:B:388:HIS:CD2	2.41	0.55
1:A:171:HIS:HB3	1:A:174:ILE:HD12	1.88	0.55
1:B:120:VAL:HG11	1:B:302:VAL:HG13	1.90	0.54
1:B:268:VAL:HG21	2:B:501:HEM:C4B	2.43	0.53
1:B:426:HIS:CD2	1:B:447:LYS:HE2	2.45	0.52
1:B:434:LYS:HB2	1:B:442:GLU:HB2	1.92	0.52
1:A:367:TRP:HB2	1:A:371:VAL:HG12	1.94	0.50
1:B:126:LEU:HD21	1:B:145:MET:HE1	1.94	0.49
1:A:368:GLY:O	1:A:371:VAL:HG13	2.13	0.48
1:B:366:ILE:HG21	1:B:389:ALA:HB1	1.95	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:15:LYS:HD2	1:B:43:GLU:HB3	1.97	0.47
1:B:97:LYS:HE3	6:B:604:HOH:O	2.15	0.47
2:A:501:HEM:HBC2	2:A:501:HEM:HMC1	1.97	0.46
1:A:81:PHE:HB3	1:A:209:ILE:HG12	1.98	0.46
1:A:426:HIS:CD2	1:A:447:LYS:HE2	2.51	0.46
1:A:388:HIS:HA	1:A:391:LYS:HD3	1.97	0.45
1:B:41:LYS:NZ	6:B:631:HOH:O	2.48	0.45
1:B:181:LEU:HD21	1:B:263:ILE:HG23	1.97	0.45
2:B:501:HEM:HMC1	2:B:501:HEM:HBC2	1.99	0.45
1:A:120:VAL:HG11	1:A:302:VAL:HG13	1.98	0.45
1:B:272:LEU:HD13	1:B:322:LEU:HG	1.99	0.45
1:B:59:LYS:NZ	6:B:621:HOH:O	2.45	0.44
1:A:15:LYS:HD2	1:A:43:GLU:HB3	2.00	0.43
1:A:366:ILE:HG21	1:A:389:ALA:HB1	1.99	0.43
1:B:149:THR:O	1:B:153:ILE:HG22	2.18	0.43
2:A:501:HEM:HBB2	2:A:501:HEM:HMB2	2.00	0.42
1:B:298:LEU:HD22	1:B:303:PRO:HB3	2.00	0.42
1:B:173:PHE:CE1	1:B:262:LEU:HD13	2.55	0.42
1:A:153:ILE:HD11	1:A:261:PHE:HB3	2.02	0.42
2:B:501:HEM:HMB2	2:B:501:HEM:HBB2	2.02	0.42
1:B:367:TRP:HB2	1:B:371:VAL:HG12	2.02	0.42
1:A:312:LYS:NZ	6:A:612:HOH:O	2.32	0.42
1:A:422:ASP:OD2	6:A:601:HOH:O	2.22	0.42
1:B:370:ASP:O	1:B:378:ARG:NH2	2.45	0.41
1:B:148:LEU:HD21	1:B:413:VAL:HG21	2.01	0.41
1:A:24:LYS:NZ	1:A:432:ASP:OD1	2.45	0.41
1:A:298:LEU:HD22	1:A:303:PRO:HB3	2.02	0.41
1:B:253:ASN:O	1:B:257:GLN:HG2	2.21	0.41
1:A:253:ASN:O	1:A:257:GLN:HG2	2.21	0.41
1:B:153:ILE:HD11	1:B:261:PHE:HB3	2.01	0.41
1:B:171:HIS:HB3	1:B:174:ILE:HD12	2.02	0.41
1:A:434:LYS:HB2	1:A:442:GLU:HB2	2.03	0.41
1:A:266:HIS:CG	1:A:267:GLU:N	2.89	0.41
1:B:44:ALA:HB3	1:B:47:ARG:HB3	2.02	0.40
1:A:272:LEU:HD13	1:A:322:LEU:HG	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	447/465 (96%)	435 (97%)	12 (3%)	0	100	100
1	B	448/465 (96%)	436 (97%)	12 (3%)	0	100	100
All	All	895/930 (96%)	871 (97%)	24 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	377/407 (93%)	376 (100%)	1 (0%)	86	84
1	B	377/407 (93%)	375 (100%)	2 (0%)	81	77
All	All	754/814 (93%)	751 (100%)	3 (0%)	84	81

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	148	LEU
1	B	409	GLU
1	A	148	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	70	ASN
1	B	204	GLN
1	B	310	GLN
1	A	70	ASN
1	A	204	GLN
1	A	359	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	PHE	A	506[B]	4	11,12,12	0.70	0	14,15,15	1.03	1 (7%)
5	PHE	B	504[A]	4	11,12,12	0.66	0	14,15,15	0.77	0
4	LXO	B	505[B]	5	12,12,13	0.86	1 (8%)	13,13,15	0.83	0
3	HOA	B	502	2	0,1,1	-	-	-	-	-
3	HOA	A	502	2	0,1,1	-	-	-	-	-
4	LXO	B	503[A]	5	12,12,13	0.69	1 (8%)	13,13,15	1.41	2 (15%)
5	PHE	A	504[A]	4	11,12,12	0.68	0	14,15,15	0.95	1 (7%)
4	LXO	A	503[A]	5	12,12,13	0.67	0	13,13,15	1.47	3 (23%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEM	B	501	1,3	50,50,50	1.38	8 (16%)	66,82,82	1.19	5 (7%)
5	PHE	B	506[B]	4	11,12,12	0.72	0	14,15,15	0.68	0
4	LXO	A	505[B]	5	12,12,13	0.97	1 (8%)	13,13,15	1.53	3 (23%)
2	HEM	A	501	1,3	50,50,50	1.38	9 (18%)	66,82,82	1.27	7 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	PHE	A	506[B]	4	-	0/8/8/8	0/1/1/1
5	PHE	B	504[A]	4	-	0/8/8/8	0/1/1/1
4	LXO	B	505[B]	5	-	2/5/6/7	0/1/1/1
4	LXO	B	503[A]	5	-	1/5/6/7	0/1/1/1
5	PHE	A	504[A]	4	-	0/8/8/8	0/1/1/1
4	LXO	A	503[A]	5	-	1/5/6/7	0/1/1/1
2	HEM	B	501	1,3	-	1/14/54/54	-
5	PHE	B	506[B]	4	-	0/8/8/8	0/1/1/1
4	LXO	A	505[B]	5	-	3/5/6/7	0/1/1/1
2	HEM	A	501	1,3	-	0/14/54/54	-

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	HEM	FE-ND	3.33	2.05	1.94
2	A	501	HEM	FE-NA	3.31	2.06	1.95
2	B	501	HEM	FE-NC	3.29	2.06	1.95
2	B	501	HEM	CAC-C3C	3.01	1.55	1.47
2	A	501	HEM	FE-NB	3.00	2.04	1.94
2	B	501	HEM	CAB-C3B	3.00	1.55	1.47
2	A	501	HEM	CAC-C3C	2.95	1.55	1.47
2	A	501	HEM	CAB-C3B	2.95	1.55	1.47
2	B	501	HEM	FE-NA	2.86	2.04	1.95
2	A	501	HEM	FE-NC	2.85	2.04	1.95
2	A	501	HEM	FE-ND	2.80	2.03	1.94
4	A	505[B]	LXO	C29-N28	2.79	1.46	1.38
4	B	505[B]	LXO	C29-N28	2.65	1.46	1.38
2	B	501	HEM	FE-NB	2.41	2.02	1.94
2	B	501	HEM	CMB-C2B	2.21	1.55	1.50
2	B	501	HEM	CMC-C2C	2.12	1.55	1.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	HEM	CMC-C2C	2.03	1.55	1.50
4	B	503[A]	LXO	C29-N28	2.02	1.44	1.38
2	A	501	HEM	CMA-C3A	2.02	1.55	1.50
2	A	501	HEM	CMD-C2D	2.01	1.55	1.50

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	HEM	C4D-ND-C1D	3.24	108.42	105.07
2	B	501	HEM	C4D-ND-C1D	3.02	108.19	105.07
2	A	501	HEM	C1B-NB-C4B	3.02	108.19	105.07
2	A	501	HEM	C4C-NC-C1C	2.96	108.25	105.35
4	A	505[B]	LXO	C27-N28-C29	2.86	129.25	122.94
4	A	503[A]	LXO	C26-C27-N28	-2.85	103.83	111.49
2	B	501	HEM	C4A-NA-C1A	2.85	108.14	105.35
2	B	501	HEM	C1B-NB-C4B	2.83	107.99	105.07
5	A	506[B]	PHE	OXT-C-CA	2.68	122.51	113.38
2	B	501	HEM	C4C-NC-C1C	2.68	107.97	105.35
2	A	501	HEM	C4A-NA-C1A	2.64	107.94	105.35
2	A	501	HEM	CBA-CAA-C2A	-2.54	105.56	112.63
4	B	503[A]	LXO	C33-N32-C31	2.44	122.58	116.85
5	A	504[A]	PHE	OXT-C-CA	2.36	121.43	113.38
4	A	505[B]	LXO	C30-C29-N28	2.31	125.77	120.97
2	A	501	HEM	C3D-C4D-ND	-2.23	107.69	110.17
4	B	503[A]	LXO	C26-C25-C24	-2.17	104.60	114.26
4	A	503[A]	LXO	C26-C25-C24	-2.14	104.72	114.26
4	A	505[B]	LXO	C34-C29-N28	-2.13	116.56	120.97
2	A	501	HEM	C3B-C2B-C1B	2.11	108.05	106.49
4	A	503[A]	LXO	C33-N32-C31	2.09	121.77	116.85
2	B	501	HEM	C3B-C2B-C1B	2.00	107.97	106.49

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	505[B]	LXO	C24-C25-C26-C27
4	A	505[B]	LXO	C24-C25-C26-C27
4	B	503[A]	LXO	C25-C26-C27-N28
4	A	503[A]	LXO	C25-C26-C27-N28
4	A	505[B]	LXO	C25-C26-C27-N28
4	B	505[B]	LXO	C25-C26-C27-N28
4	A	505[B]	LXO	C26-C27-N28-C29

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	B	501	HEM	CAA-CBA-CGA-O2A




There are no ring outliers.


4 monomers are involved in 7 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	502	HOA	1	0
3	A	502	HOA	1	0
2	B	501	HEM	3	0
2	A	501	HEM	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	450/465 (96%)	0.19	17 (3%)	44 50	12, 23, 42, 66	1 (0%)
1	B	451/465 (96%)	0.15	19 (4%)	40 46	12, 23, 41, 60	1 (0%)
All	All	901/930 (96%)	0.17	36 (3%)	42 48	12, 23, 41, 66	2 (0%)

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	384	ALA	4.6
1	B	136	ASP	4.4
1	A	136	ASP	4.4
1	A	197	ALA	4.2
1	A	194	ASP	4.2
1	A	198	TYR	3.4
1	A	227	GLY	3.3
1	B	197	ALA	3.3
1	B	194	ASP	3.2
1	A	195	ASP	3.2
1	A	264	ALA	3.1
1	A	196	PRO	3.1
1	B	193	PRO	3.0
1	B	263	ILE	2.9
1	B	454	PRO	2.8
1	A	226	SER	2.8
1	B	196	PRO	2.8
1	A	199	ASP	2.8
1	A	175	THR	2.6
1	A	200	GLU	2.6
1	B	227	GLY	2.5
1	B	453	ILE	2.5
1	B	385	ILE	2.4
1	B	153	ILE	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	189	GLN	2.3
1	B	13	GLU	2.3
1	A	174	ILE	2.2
1	B	195	ASP	2.2
1	B	369	ASP	2.2
1	B	3	LYS	2.1
1	A	225	ALA	2.1
1	A	3	LYS	2.1
1	A	215	LEU	2.1
1	A	263	ILE	2.1
1	B	198	TYR	2.0
1	B	225	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	HOA	A	502	2/2	0.78	0.23	23,23,23,27	0
4	LXO	A	503[A]	12/13	0.84	0.15	14,18,20,21	12
4	LXO	A	505[B]	12/13	0.87	0.14	15,20,22,23	12
3	HOA	B	502	2/2	0.90	0.16	20,20,20,27	0
4	LXO	B	503[A]	12/13	0.90	0.12	13,15,18,18	12
5	PHE	A	504[A]	12/12	0.90	0.11	15,17,20,21	12
5	PHE	B	506[B]	12/12	0.92	0.10	14,17,21,21	12
4	LXO	B	505[B]	12/13	0.93	0.10	14,17,19,19	12
5	PHE	B	504[A]	12/12	0.94	0.09	12,16,18,20	12
5	PHE	A	506[B]	12/12	0.96	0.08	13,17,21,23	12
2	HEM	B	501	43/43	0.98	0.05	11,14,16,21	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	HEM	A	501	43/43	0.98	0.05	11,15,18,23	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around HOA A 502:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around LXO A 503 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around LXO A 505 (B):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HOA B 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around LXO B 503 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around LXO B 505 (B):**


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.