



Full wwPDB NMR Structure Validation Report ⓘ

Mar 8, 2026 – 05:38 PM UTC

PDB ID : 2JT6 / pdb_00002jt6
BMRB ID : 15396
Title : Solution structure of matrix metalloproteinase 3 (MMP-3) in the presence of 3
-4'-cyanobiphenyl-4-yloxy)-n-hydroxypropionamide (MMP-3 inhibitor VII)
Authors : Alcaraz, L.A.; Banci, L.; Bertini, I.; Cantini, F.; Donaire, A.; Gonnelli, L.
Deposited on : 2007-07-23

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 14%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	161	

2 Ensemble composition and analysis ⓘ

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.

3 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2529 atoms, of which 1226 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Stromelysin-1.

Mol	Chain	Residues	Atoms						Trace
1	A	161	Total	C	H	N	O	S	0
			2491	818	1213	213	245	2	


- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	
2	A	2	Total	Zn
			2	2

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	
3	A	2	Total	Ca
			2	2

- Molecule 4 is 3-[(4'-cyanobiphenyl-4-yl)oxy]-N-hydroxypropanamide (CCD ID: JT6) (formula: C₁₆H₁₄N₂O₃).




Mol	Chain	Residues	Atoms				
4	A	1	Total	C	H	N	O
			34	16	13	2	3

4 Residue-property plots

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: Stromelysin-1



5 Refinement protocol and experimental data overview

The models were refined using the following method: *molecular dynamics*.

Of the 100 calculated structures, 1 were deposited, based on the following criterion: *structures with the least restraint violations*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR NIH	refinement	
AutoDock	refinement	3.0

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	298
Number of shifts mapped to atoms	298
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	14%

6 Model quality ⓘ

6.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CA, JT6

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1278	1213	1207	25
All	All	1303	1226	1220	25

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:A:149:ARG:HA	1:A:157:PHE:HB3	0.69	1.62
1:A:120:ALA:HB1	1:A:200:ALA:HA	0.67	1.66
1:A:135:LEU:HD13	1:A:137:GLU:HG2	0.65	1.68
1:A:200:ALA:HB1	1:A:218:LEU:HD21	0.59	1.73
1:A:169:ALA:HB3	1:A:170:PRO:HD3	0.55	1.79
1:A:148:VAL:HG22	1:A:149:ARG:H	0.51	1.64
1:A:135:LEU:HB2	1:A:137:GLU:OE2	0.51	2.05
1:A:176:GLY:HA2	1:A:206:SER:CB	0.51	2.35
1:A:135:LEU:HD11	1:A:140:ALA:HB2	0.51	1.82
1:A:209:LEU:HD23	1:A:219:MET:HG3	0.49	1.82
1:A:99:TYR:CZ	1:A:134:ARG:HB2	0.47	2.44
1:A:142:ILE:HA	1:A:176:GLY:O	0.47	2.08
1:A:176:GLY:HA2	1:A:206:SER:HB2	0.46	1.87

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:A:108:LEU:HB3	1:A:109:PRO:HD2	0.46	1.88
1:A:164:LEU:HD22	1:A:193:THR:HG23	0.45	1.88
1:A:137:GLU:H	1:A:137:GLU:CD	0.45	2.19
1:A:187:THR:HG21	1:A:191:THR:HB	0.45	1.89
1:A:187:THR:O	1:A:194:ASN:HA	0.44	2.12
1:A:220:TYR:CD1	1:A:221:PRO:HD2	0.44	2.47
1:A:122:LYS:HA	1:A:125:GLU:HB3	0.42	1.92
1:A:89:ILE:HG13	1:A:93:ARG:NH1	0.42	2.30
1:A:212:SER:HB3	1:A:237:ASP:OD2	0.42	2.15
1:A:119:LYS:O	1:A:123:VAL:HB	0.41	2.16
1:A:149:ARG:CA	1:A:157:PHE:HB3	0.41	2.40
1:A:212:SER:OG	1:A:219:MET:HB2	0.41	2.16

6.3 Torsion angles

6.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	159/161 (99%)	125 (79%)	29 (18%)	5 (3%)	5	37
All	All	159/161 (99%)	125 (79%)	29 (18%)	5 (3%)	5	37

All 5 Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	137	GLU
1	A	138	GLY
1	A	154	PHE
1	A	169	ALA
1	A	174	ILE

6.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation

was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	135/135 (100%)	102 (76%)	33 (24%)	2	26
All	All	135/135 (100%)	102 (76%)	33 (24%)	2	26

All 33 residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	98	THR
1	A	102	VAL
1	A	110	LYS
1	A	111	ASP
1	A	113	VAL
1	A	119	LYS
1	A	122	LYS
1	A	123	VAL
1	A	125	GLU
1	A	127	VAL
1	A	128	THR
1	A	131	THR
1	A	133	SER
1	A	134	ARG
1	A	135	LEU
1	A	136	TYR
1	A	137	GLU
1	A	143	MET
1	A	145	SER
1	A	157	PHE
1	A	162	ASN
1	A	163	VAL
1	A	187	THR
1	A	190	THR
1	A	191	THR
1	A	195	LEU
1	A	206	SER
1	A	215	THR
1	A	224	HIS
1	A	225	SER
1	A	227	THR
1	A	230	THR
1	A	236	GLN

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.6 Ligand geometry ⓘ

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
4	JT6	A	1	2	22,22,22	1.97	5 (22%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
4	JT6	A	1	2	26,28,28	2.75	8 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means

no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	JT6	A	1	2	-	0,14,14,14	0,2,2,2

All bond outliers are listed below. They are sorted according to the Z-score.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1	JT6	O5-N2	4.82	1.28	1.40
4	A	1	JT6	C13-C8	3.22	1.44	1.38
4	A	1	JT6	O7-C8	2.76	1.31	1.37
4	A	1	JT6	C9-C8	2.61	1.43	1.38
4	A	1	JT6	C19-C14	2.42	1.44	1.39

All angle outliers are listed below. They are sorted according to the Z-score.


Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1	JT6	O5-N2-C1	9.76	134.22	119.79
4	A	1	JT6	C6-O7-C8	6.40	134.56	117.93
4	A	1	JT6	O4-C1-C3	2.72	117.09	122.02
4	A	1	JT6	O4-C1-N2	2.59	126.45	123.27
4	A	1	JT6	C9-C8-C13	2.51	116.50	120.16
4	A	1	JT6	C18-C17-C16	2.14	115.29	118.96
4	A	1	JT6	C9-C10-C11	2.05	123.75	121.12
4	A	1	JT6	C13-C12-C11	2.03	123.72	121.12

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 14% for the well-defined parts and 14% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	298
Number of shifts mapped to atoms	298
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	1

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	0	—	None (insufficient data)
$^{13}\text{C}_\beta$	0	—	None (insufficient data)
$^{13}\text{C}'$	0	—	None (insufficient data)
^{15}N	149	0.16 ± 0.54	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 14%, i.e. 298 atoms were assigned a chemical shift out of a possible 2152. 0 out of 24 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	298/798 (37%)	149/325 (46%)	0/322 (0%)	149/151 (99%)
Sidechain	0/1108 (0%)	0/723 (0%)	0/352 (0%)	0/33 (0%)

Continued on next page...

Continued from previous page...

	Total	¹H	¹³C	¹⁵N
Aromatic	0/246 (0%)	0/122 (0%)	0/111 (0%)	0/13 (0%)
Overall	298/2152 (14%)	149/1170 (13%)	0/785 (0%)	149/197 (76%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 14%, i.e. 298 atoms were assigned a chemical shift out of a possible 2152. 0 out of 24 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹H	¹³C	¹⁵N
Backbone	298/798 (37%)	149/325 (46%)	0/322 (0%)	149/151 (99%)
Sidechain	0/1108 (0%)	0/723 (0%)	0/352 (0%)	0/33 (0%)
Aromatic	0/246 (0%)	0/122 (0%)	0/111 (0%)	0/13 (0%)
Overall	298/2152 (14%)	149/1170 (13%)	0/785 (0%)	149/197 (76%)

7.1.4 Statistically unusual chemical shifts [i](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	218	LEU	H	11.51	5.09 – 11.34	5.3

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:

