



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 02:41 AM UTC

PDB ID : 5KDW / pdb_00005kdw
Title : IMPa metallopeptidase from Pseudomonas aeruginosa
Authors : Noach, I.; Boraston, A.B.
Deposited on : 2016-06-08
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	885	 97%
1	B	885	 97%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	PO4	A	1024	-	-	X	-
4	PO4	A	1025	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 15932 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Metallopeptidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	883	Total	C	N	O	S	0	9	0
			6843	4278	1225	1324	16			
1	B	883	Total	C	N	O	S	0	8	0
			6824	4268	1220	1320	16			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	41	MET	-	initiating methionine	UNP Q9I5W4
A	924	LEU	-	expression tag	UNP Q9I5W4
A	925	GLU	-	expression tag	UNP Q9I5W4
B	41	MET	-	initiating methionine	UNP Q9I5W4
B	924	LEU	-	expression tag	UNP Q9I5W4
B	925	GLU	-	expression tag	UNP Q9I5W4

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		
3	B	1	Total	Zn	0	0
			1	1		

- Molecule 4 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	P	0	0
			5	4	1		
4	A	1	Total	O	P	0	0
			5	4	1		
4	A	1	Total	O	P	0	0
			5	4	1		
4	A	1	Total	O	P	0	0
			5	4	1		
4	B	1	Total	O	P	0	0
			5	4	1		
4	B	1	Total	O	P	0	0
			5	4	1		
4	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1027	Total	O	0	0
			1027	1027		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1033	Total 1033	O 1033	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Metallopeptidase



- Molecule 1: Metallopeptidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	98.92Å 132.73Å 103.07Å 90.00° 103.81° 90.00°	Depositor
Resolution (Å)	100.09 – 1.85 100.09 – 1.85	Depositor EDS
% Data completeness (in resolution range)	100.0 (100.09-1.85) 100.0 (100.09-1.85)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.15 (at 1.86Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.153 , 0.186 0.165 , 0.195	Depositor DCC
R_{free} test set	10928 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	15.6	Xtriage
Anisotropy	0.327	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 45.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.065 for l,-k,h	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	15932	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.12% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, ZN, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.99	0/6991	0.96	4/9514 (0.0%)
1	B	1.01	0/6972	0.96	3/9490 (0.0%)
All	All	1.00	0/13963	0.96	7/19004 (0.0%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	602	ASP	N-CA-C	7.39	120.27	111.33
1	B	602	ASP	CB-CA-C	-7.01	98.76	110.68
1	A	607	VAL	CB-CA-C	-6.12	103.87	112.14
1	A	69	ARG	CG-CD-NE	-5.33	100.29	112.00
1	B	681	TYR	CA-CB-CG	5.26	123.37	113.90
1	A	681	TYR	CA-CB-CG	5.13	123.14	113.90
1	A	837	ARG	NE-CZ-NH2	5.03	123.73	119.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6843	0	6626	19	0
1	B	6824	0	6604	16	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	84	0	126	2	0
2	B	84	0	126	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	20	0	0	4	0
4	B	15	0	0	0	0
5	A	1027	0	0	9	0
5	B	1033	0	0	3	0
All	All	15932	0	13482	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:642[A]:GLU:OE1	5:A:1102:HOH:O	1.96	0.84
1:A:771[A]:GLU:OE1	5:A:1101:HOH:O	1.96	0.83
1:A:555:GLN:HA	1:A:555:GLN:OE1	1.87	0.74
1:B:103:MET:HE2	1:B:271:GLU:HG2	1.70	0.72
1:A:633:GLN:NE2	5:A:1104:HOH:O	2.18	0.71
1:A:646:ARG:NH1	5:A:1109:HOH:O	2.25	0.68
1:A:103:MET:HE2	1:A:271:GLU:HG2	1.79	0.65
1:A:145:THR:OG1	5:A:1103:HOH:O	2.15	0.64
1:A:758:GLU:OE2	1:A:768[A]:ARG:NH1	2.26	0.62
1:A:638:ALA:O	1:A:642[B]:GLU:HG2	2.01	0.60
1:A:385:ASN:HB2	5:A:1679:HOH:O	2.02	0.58
4:A:1024:PO4:O3	4:A:1025:PO4:O1	2.24	0.56
1:A:642[B]:GLU:O	1:A:646:ARG:HG3	2.07	0.54
1:A:469:ILE:HD13	1:A:477:LEU:HD12	1.89	0.54
1:B:579:LEU:C	1:B:579:LEU:HD12	2.35	0.52
1:B:385:ASN:OD1	5:B:1101:HOH:O	2.19	0.51
1:A:361:GLN:HE22	2:A:1007:EDO:HO1	1.60	0.47
1:B:469:ILE:O	1:B:469:ILE:HD12	2.15	0.47
1:B:574:ASP:HA	5:B:1165:HOH:O	2.15	0.47
1:A:579:LEU:C	1:A:579:LEU:HD12	2.40	0.46
4:A:1024:PO4:O4	4:A:1025:PO4:O4	2.33	0.45
1:B:469:ILE:HD12	1:B:469:ILE:C	2.42	0.44
1:B:254:TRP:CE3	1:B:278:TYR:HD1	2.34	0.44
1:B:575:LYS:HA	1:B:575:LYS:HD2	1.70	0.44
1:B:598:SER:O	1:B:602:ASP:HB2	2.18	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:111:ASP:OD2	4:A:1026:PO4:O2	2.36	0.43
1:A:725:HIS:HD2	1:A:804:TRP:CG	2.37	0.43
4:A:1024:PO4:O2	5:A:1105:HOH:O	2.21	0.43
2:A:1016:EDO:H21	5:A:1919:HOH:O	2.18	0.42
1:A:330:ARG:NH1	5:A:1125:HOH:O	2.46	0.42
1:A:784:ALA:O	1:A:788:GLN:HG2	2.20	0.41
1:A:642[A]:GLU:O	1:A:646:ARG:HG3	2.21	0.41
1:B:465:LEU:HD12	1:B:465:LEU:C	2.46	0.41
1:B:103:MET:HE3	1:B:103:MET:HB3	1.85	0.41
1:B:725:HIS:HD2	1:B:804:TRP:CG	2.39	0.41
1:B:761:PRO:HD2	5:B:1463:HOH:O	2.21	0.41
1:B:372:LEU:HD23	1:B:372:LEU:HA	1.95	0.41
1:B:696:HIS:CE1	1:B:719:ASN:HD21	2.39	0.41
1:B:784:ALA:O	1:B:788:GLN:HG2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	890/885 (101%)	877 (98%)	13 (2%)	0	100	100
1	B	889/885 (100%)	875 (98%)	14 (2%)	0	100	100
All	All	1779/1770 (100%)	1752 (98%)	27 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	707/701 (101%)	704 (100%)	3 (0%)	84	82
1	B	704/701 (100%)	700 (99%)	4 (1%)	78	73
All	All	1411/1402 (101%)	1404 (100%)	7 (0%)	84	77

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	771[A]	GLU
1	A	771[B]	GLU
1	A	924	LEU
1	B	45	GLU
1	B	575	LYS
1	B	646	ARG
1	B	739	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	83	GLN
1	A	361	GLN
1	A	903	ASN
1	B	163	GLN
1	B	255	ASN
1	B	262	GLN
1	B	270	GLN
1	B	802	GLN
1	B	903	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 51 ligands modelled in this entry, 2 are monoatomic - leaving 49 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	A	1011	-	3,3,3	0.45	0	2,2,2	0.59	0
2	EDO	B	1010	-	3,3,3	0.30	0	2,2,2	0.57	0
2	EDO	A	1021	-	3,3,3	0.55	0	2,2,2	0.24	0
4	PO4	B	1023	-	4,4,4	0.68	0	6,6,6	1.05	0
4	PO4	A	1024	-	4,4,4	0.72	0	6,6,6	0.68	0
4	PO4	B	1024	-	4,4,4	1.07	0	6,6,6	1.41	2 (33%)
2	EDO	A	1014	-	3,3,3	0.43	0	2,2,2	0.62	0
2	EDO	A	1020	-	3,3,3	0.63	0	2,2,2	0.16	0
2	EDO	B	1011	-	3,3,3	0.30	0	2,2,2	0.48	0
2	EDO	B	1017	-	3,3,3	0.31	0	2,2,2	0.76	0
2	EDO	A	1007	-	3,3,3	0.45	0	2,2,2	0.07	0
2	EDO	B	1007	-	3,3,3	0.22	0	2,2,2	0.06	0
2	EDO	A	1006	-	3,3,3	0.21	0	2,2,2	0.69	0
2	EDO	A	1018	-	3,3,3	0.57	0	2,2,2	0.20	0
2	EDO	A	1013	-	3,3,3	0.24	0	2,2,2	1.17	0
2	EDO	B	1002	-	3,3,3	0.51	0	2,2,2	0.39	0
4	PO4	A	1026	-	4,4,4	1.13	1 (25%)	6,6,6	1.16	0
2	EDO	A	1012	-	3,3,3	0.29	0	2,2,2	0.28	0
2	EDO	B	1014	-	3,3,3	0.17	0	2,2,2	1.40	0
4	PO4	B	1025	-	4,4,4	0.96	0	6,6,6	1.29	1 (16%)
2	EDO	A	1009	-	3,3,3	0.44	0	2,2,2	0.24	0
2	EDO	B	1020	-	3,3,3	0.24	0	2,2,2	1.04	0
2	EDO	A	1019	-	3,3,3	0.42	0	2,2,2	0.42	0
2	EDO	A	1001	-	3,3,3	0.38	0	2,2,2	0.55	0
2	EDO	B	1016	-	3,3,3	0.59	0	2,2,2	0.38	0
2	EDO	B	1013	-	3,3,3	0.35	0	2,2,2	0.51	0
2	EDO	A	1017	-	3,3,3	0.35	0	2,2,2	1.47	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	EDO	B	1021	-	3,3,3	0.65	0	2,2,2	0.17	0
2	EDO	B	1019	-	3,3,3	0.29	0	2,2,2	0.94	0
2	EDO	B	1018	-	3,3,3	0.31	0	2,2,2	0.58	0
2	EDO	B	1001	-	3,3,3	0.27	0	2,2,2	0.63	0
2	EDO	A	1002	-	3,3,3	0.53	0	2,2,2	0.26	0
4	PO4	A	1023	-	4,4,4	0.89	0	6,6,6	1.61	1 (16%)
2	EDO	A	1003	-	3,3,3	0.18	0	2,2,2	1.16	0
2	EDO	A	1008	-	3,3,3	0.79	0	2,2,2	0.52	0
2	EDO	A	1005	-	3,3,3	0.53	0	2,2,2	0.65	0
2	EDO	A	1015	-	3,3,3	0.42	0	2,2,2	0.44	0
2	EDO	B	1012	-	3,3,3	0.26	0	2,2,2	0.66	0
2	EDO	A	1016	-	3,3,3	0.29	0	2,2,2	0.65	0
2	EDO	A	1004	-	3,3,3	0.42	0	2,2,2	0.50	0
2	EDO	B	1003	-	3,3,3	0.64	0	2,2,2	0.11	0
2	EDO	B	1004	-	3,3,3	0.30	0	2,2,2	0.31	0
2	EDO	B	1008	-	3,3,3	0.30	0	2,2,2	0.44	0
2	EDO	B	1005	-	3,3,3	0.53	0	2,2,2	0.76	0
2	EDO	B	1009	-	3,3,3	0.38	0	2,2,2	0.47	0
2	EDO	B	1015	-	3,3,3	0.64	0	2,2,2	0.11	0
2	EDO	B	1006	-	3,3,3	0.58	0	2,2,2	0.47	0
2	EDO	A	1010	-	3,3,3	0.46	0	2,2,2	0.28	0
4	PO4	A	1025	-	4,4,4	0.74	0	6,6,6	1.04	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	1011	-	-	0/1/1/1	-
2	EDO	B	1010	-	-	1/1/1/1	-
2	EDO	A	1021	-	-	0/1/1/1	-
2	EDO	A	1014	-	-	1/1/1/1	-
2	EDO	A	1020	-	-	1/1/1/1	-
2	EDO	B	1011	-	-	0/1/1/1	-
2	EDO	B	1017	-	-	1/1/1/1	-
2	EDO	A	1007	-	-	1/1/1/1	-
2	EDO	B	1007	-	-	1/1/1/1	-
2	EDO	A	1006	-	-	0/1/1/1	-
2	EDO	A	1018	-	-	1/1/1/1	-
2	EDO	A	1013	-	-	0/1/1/1	-
2	EDO	B	1002	-	-	0/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	1012	-	-	0/1/1/1	-
2	EDO	B	1014	-	-	0/1/1/1	-
2	EDO	A	1009	-	-	1/1/1/1	-
2	EDO	B	1020	-	-	1/1/1/1	-
2	EDO	A	1019	-	-	0/1/1/1	-
2	EDO	A	1001	-	-	1/1/1/1	-
2	EDO	B	1016	-	-	1/1/1/1	-
2	EDO	B	1013	-	-	1/1/1/1	-
2	EDO	A	1017	-	-	0/1/1/1	-
2	EDO	B	1021	-	-	0/1/1/1	-
2	EDO	B	1019	-	-	1/1/1/1	-
2	EDO	B	1018	-	-	1/1/1/1	-
2	EDO	B	1001	-	-	0/1/1/1	-
2	EDO	A	1002	-	-	0/1/1/1	-
2	EDO	A	1003	-	-	0/1/1/1	-
2	EDO	A	1008	-	-	0/1/1/1	-
2	EDO	A	1005	-	-	0/1/1/1	-
2	EDO	A	1015	-	-	0/1/1/1	-
2	EDO	B	1012	-	-	0/1/1/1	-
2	EDO	A	1016	-	-	1/1/1/1	-
2	EDO	A	1004	-	-	0/1/1/1	-
2	EDO	B	1003	-	-	1/1/1/1	-
2	EDO	B	1004	-	-	1/1/1/1	-
2	EDO	B	1008	-	-	0/1/1/1	-
2	EDO	B	1005	-	-	0/1/1/1	-
2	EDO	B	1009	-	-	1/1/1/1	-
2	EDO	B	1015	-	-	0/1/1/1	-
2	EDO	B	1006	-	-	0/1/1/1	-
2	EDO	A	1010	-	-	1/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1026	PO4	P-O1	2.21	1.55	1.50

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1023	PO4	O4-P-O1	-3.10	99.98	110.95
4	B	1024	PO4	O2-P-O1	-2.64	101.63	110.95
4	B	1025	PO4	O2-P-O1	-2.17	103.29	110.95
4	B	1024	PO4	O4-P-O3	2.03	114.24	107.91

There are no chirality outliers.

All (19) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1010	EDO	O1-C1-C2-O2
2	B	1003	EDO	O1-C1-C2-O2
2	B	1004	EDO	O1-C1-C2-O2
2	B	1016	EDO	O1-C1-C2-O2
2	B	1020	EDO	O1-C1-C2-O2
2	A	1018	EDO	O1-C1-C2-O2
2	A	1020	EDO	O1-C1-C2-O2
2	A	1016	EDO	O1-C1-C2-O2
2	B	1013	EDO	O1-C1-C2-O2
2	A	1007	EDO	O1-C1-C2-O2
2	B	1009	EDO	O1-C1-C2-O2
2	B	1010	EDO	O1-C1-C2-O2
2	B	1018	EDO	O1-C1-C2-O2
2	B	1007	EDO	O1-C1-C2-O2
2	B	1017	EDO	O1-C1-C2-O2
2	B	1019	EDO	O1-C1-C2-O2
2	A	1001	EDO	O1-C1-C2-O2
2	A	1009	EDO	O1-C1-C2-O2
2	A	1014	EDO	O1-C1-C2-O2

There are no ring outliers.

5 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1024	PO4	3	0
2	A	1007	EDO	1	0
4	A	1026	PO4	1	0
2	A	1016	EDO	1	0
4	A	1025	PO4	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	883/885 (99%)	-0.31	7 (0%) 82 86	8, 20, 34, 62	9 (1%)
1	B	883/885 (99%)	-0.35	11 (1%) 76 80	8, 19, 34, 55	8 (0%)
All	All	1766/1770 (99%)	-0.33	18 (1%) 79 83	8, 20, 34, 62	17 (0%)

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	556	PRO	4.4
1	B	557	GLY	3.9
1	B	42	ALA	3.1
1	A	557	GLY	3.1
1	B	602	ASP	2.7
1	B	759	ALA	2.6
1	A	558	GLU	2.5
1	B	109	ASN	2.4
1	A	759	ALA	2.3
1	B	649	ASP	2.3
1	B	558	GLU	2.3
1	B	601	LYS	2.2
1	A	42	ALA	2.2
1	A	206	CYS	2.2
1	A	672[A]	CYS	2.2
1	B	646	ARG	2.2
1	B	600	ASP	2.1
1	B	672[A]	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	EDO	B	1020	4/4	0.75	0.19	44,46,48,48	0
2	EDO	B	1016	4/4	0.84	0.16	43,48,51,56	0
2	EDO	A	1016	4/4	0.84	0.16	46,47,49,55	0
2	EDO	A	1020	4/4	0.85	0.17	50,52,53,55	0
2	EDO	A	1014	4/4	0.86	0.17	40,45,46,52	0
2	EDO	A	1008	4/4	0.86	0.16	34,37,41,42	0
2	EDO	A	1018	4/4	0.86	0.17	33,35,36,42	0
2	EDO	A	1017	4/4	0.88	0.18	26,34,40,55	0
2	EDO	B	1003	4/4	0.88	0.15	41,43,45,49	0
2	EDO	B	1021	4/4	0.88	0.15	31,33,34,38	0
2	EDO	A	1015	4/4	0.89	0.15	52,52,54,54	0
2	EDO	A	1005	4/4	0.89	0.12	30,32,33,33	0
2	EDO	B	1013	4/4	0.89	0.15	35,40,43,44	0
4	PO4	A	1025	5/5	0.89	0.09	41,45,50,51	0
2	EDO	A	1010	4/4	0.90	0.15	46,47,49,53	0
2	EDO	A	1019	4/4	0.90	0.14	44,44,50,54	0
2	EDO	B	1017	4/4	0.90	0.15	43,44,44,46	0
2	EDO	A	1007	4/4	0.91	0.13	36,38,40,43	0
2	EDO	B	1004	4/4	0.91	0.12	34,37,40,43	0
2	EDO	B	1018	4/4	0.91	0.13	31,37,40,48	0
2	EDO	B	1006	4/4	0.91	0.10	26,28,29,29	0
2	EDO	B	1011	4/4	0.91	0.15	46,46,46,47	0
2	EDO	A	1003	4/4	0.91	0.12	35,36,38,42	0
4	PO4	A	1026	5/5	0.91	0.15	30,42,43,44	0
2	EDO	A	1013	4/4	0.92	0.13	30,36,38,38	0
2	EDO	A	1011	4/4	0.92	0.14	30,41,42,46	0
2	EDO	B	1014	4/4	0.92	0.14	30,35,38,38	0
2	EDO	A	1021	4/4	0.92	0.12	27,33,33,39	0
2	EDO	B	1009	4/4	0.92	0.11	30,32,33,39	0
2	EDO	B	1012	4/4	0.93	0.10	40,42,44,46	0
2	EDO	B	1008	4/4	0.93	0.11	37,38,41,42	0
2	EDO	B	1005	4/4	0.93	0.09	33,33,34,35	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	EDO	B	1010	4/4	0.93	0.11	38,38,39,43	0
2	EDO	A	1012	4/4	0.93	0.11	34,36,37,41	0
2	EDO	B	1007	4/4	0.94	0.09	35,36,37,43	0
2	EDO	A	1006	4/4	0.94	0.09	31,33,34,39	0
2	EDO	A	1009	4/4	0.94	0.09	32,32,32,34	0
4	PO4	B	1023	5/5	0.94	0.14	28,35,38,39	0
4	PO4	B	1025	5/5	0.94	0.13	30,30,40,41	0
2	EDO	B	1015	4/4	0.95	0.10	26,31,31,32	0
2	EDO	B	1001	4/4	0.95	0.09	23,29,29,31	0
2	EDO	B	1019	4/4	0.95	0.10	25,30,36,44	0
2	EDO	A	1001	4/4	0.96	0.10	27,31,33,33	0
4	PO4	A	1024	5/5	0.96	0.07	37,43,46,47	0
2	EDO	B	1002	4/4	0.96	0.07	19,22,24,24	0
2	EDO	A	1002	4/4	0.97	0.07	17,20,21,22	0
2	EDO	A	1004	4/4	0.97	0.07	27,27,29,31	0
3	ZN	A	1022	1/1	0.97	0.07	29,29,29,29	1
3	ZN	B	1022	1/1	0.97	0.09	27,27,27,27	1
4	PO4	B	1024	5/5	0.97	0.12	23,23,27,28	0
4	PO4	A	1023	5/5	0.97	0.15	25,26,32,34	0

6.5 Other polymers

There are no such residues in this entry.