



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:40 PM UTC

PDB ID : 7KWU / pdb_00007kwu
Title : Crystal Structure of HIV-1 RT in Complex with 16c (K07-15)
Authors : Ruiz, F.X.; Arnold, E.
Deposited on : 2020-12-02
Resolution : 2.02 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.02 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	13299 (2.04-2.00)
Clashscore	190562	1022 (2.02-2.02)
Ramachandran outliers	187476	1014 (2.02-2.02)
Sidechain outliers	187428	1014 (2.02-2.02)
RSRZ outliers	180081	13314 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	557	14% 86% 13%
2	B	429	13% 87% 9% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SO4	A	603	-	-	X	-

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 8826 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Reverse transcriptase p66.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	556	Total	C	N	O	S	0	5	0
			4557	2947	760	842	8			

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	initiating methionine	UNP P03366
A	0	VAL	-	expression tag	UNP P03366
A	172	ALA	LYS	engineered mutation	UNP P03366
A	173	ALA	LYS	engineered mutation	UNP P03366
A	280	SER	CYS	engineered mutation	UNP P03366

- Molecule 2 is a protein called Reverse transcriptase p51.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	414	Total	C	N	O	S	0	7	0
			3482	2271	574	630	7			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	GLY	-	expression tag	UNP P03366
B	280	SER	CYS	engineered mutation	UNP P03366

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Mg	0	0
			1	1		

- Molecule 6 is 4-[(4-{[4-(4-cyano-2,6-dimethylphenoxy)-5-(pyridin-4-yl)pyrimidin-2-yl]amino}piperidin-1-yl)methyl]benzamide (CCD ID: K7F) (formula: C₃₁H₃₁N₇O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	1
			40	31	7	2		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	387	Total O 387 387	0	1
7	B	309	Total O 309 309	0	3


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Reverse transcriptase p66



• Molecule 2: Reverse transcriptase p51



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	163.38Å 72.62Å 109.72Å 90.00° 100.20° 90.00°	Depositor
Resolution (Å)	29.54 – 2.02 29.54 – 2.02	Depositor EDS
% Data completeness (in resolution range)	100.0 (29.54-2.02) 99.9 (29.54-2.02)	Depositor EDS
R_{merge}	0.50	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.42 (at 2.03Å)	Xtriage
Refinement program	PHENIX 1.18.2_3874	Depositor
R, R_{free}	0.211 , 0.259 0.212 , 0.260	Depositor DCC
R_{free} test set	4102 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	31.7	Xtriage
Anisotropy	0.045	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 53.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8826	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: K7F, SO4, MG, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.19	0/4676	0.42	5/6357 (0.1%)
2	B	0.17	0/3586	0.36	2/4875 (0.0%)
All	All	0.18	0/8262	0.40	7/11232 (0.1%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	66	LYS	N-CA-C	6.94	121.80	111.04
2	B	90	VAL	N-CA-C	6.26	116.93	110.36
1	A	90	VAL	N-CA-C	-6.21	106.45	112.29
1	A	222	GLN	N-CA-C	-5.90	98.24	110.80
1	A	270	ILE	N-CA-CB	-5.82	104.27	112.12
2	B	90	VAL	CB-CA-C	-5.53	104.80	111.88
1	A	66	LYS	CB-CA-C	-5.29	102.97	111.39

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4557	0	4602	56	0
2	B	3482	0	3488	26	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	20	0	0	2	0
3	B	10	0	0	1	0
4	A	8	0	12	1	0
4	B	12	0	18	0	0
5	A	1	0	0	0	0
6	A	40	0	0	1	0
7	A	387	0	0	4	0
7	B	309	0	0	2	0
All	All	8826	0	8120	80	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (80) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:424:LYS:HE2	7:A:780:HOH:O	1.51	1.07
1:A:296:THR:HG23	1:A:299:ALA:H	1.45	0.82
2:B:99:GLY:HA2	2:B:102:LYS:HE2	1.75	0.69
1:A:195:ILE:HD12	1:A:195:ILE:H	1.59	0.67
1:A:241:VAL:HG11	1:A:244:ILE:HD11	1.77	0.66
2:B:5:ILE:HG23	2:B:6:GLU:HG3	1.76	0.66
1:A:13:LYS:HB2	1:A:16:MET:HG3	1.79	0.64
1:A:16:MET:HE3	1:A:83:ARG:HA	1.80	0.64
1:A:277:ARG:NH1	3:A:603:SO4:O3	2.31	0.63
1:A:244:ILE:HD13	1:A:267:ALA:HB2	1.81	0.62
1:A:544:GLY:HA2	2:B:286[B]:THR:HG22	1.83	0.59
1:A:39:THR:O	1:A:43:LYS:HG3	2.02	0.59
2:B:108:VAL:HB	2:B:232:TYR:HB3	1.86	0.57
2:B:424:LYS:HD2	7:B:775:HOH:O	2.05	0.56
1:A:28:GLU:HG2	1:A:32:LYS:HE3	1.88	0.56
2:B:195:ILE:O	2:B:195:ILE:HG13	2.05	0.55
1:A:543:GLY:N	2:B:283:LEU:O	2.29	0.55
2:B:68:SER:HB2	2:B:230:MET:HE1	1.89	0.55
1:A:253:THR:HG22	1:A:292:VAL:HG22	1.88	0.55
2:B:424:LYS:NZ	3:B:502:SO4:O2	2.39	0.54
1:A:270:ILE:HG13	1:A:314:VAL:HG13	1.90	0.53
2:B:175:ASN:OD1	2:B:201:LYS:NZ	2.40	0.53
1:A:536:VAL:HB	1:A:542:ILE:HD12	1.91	0.52
1:A:209:LEU:HD13	1:A:216:THR:HG21	1.91	0.52
1:A:296:THR:HG23	1:A:299:ALA:N	2.21	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:549:ASP:OD2	1:A:550:LYS:HE3	2.10	0.51
1:A:70:LYS:HD2	1:A:72:ARG:HE	1.75	0.51
1:A:332:GLN:OE1	1:A:338[B]:THR:OG1	2.29	0.51
1:A:194:GLU:OE2	1:A:194:GLU:N	2.25	0.51
1:A:335:GLY:HA3	1:A:356[A]:ARG:HD3	1.93	0.51
1:A:220:LYS:O	1:A:220:LYS:HD2	2.11	0.50
1:A:220:LYS:HD2	1:A:220:LYS:C	2.37	0.50
1:A:332:GLN:CD	1:A:338[B]:THR:OG1	2.55	0.49
2:B:236:PRO:HA	2:B:239:TRP:CD2	2.47	0.49
1:A:332:GLN:CD	1:A:338[B]:THR:HG1	2.20	0.49
1:A:513:SER:O	1:A:519:ASN:ND2	2.34	0.48
1:A:94:ILE:HG21	1:A:230:MET:HE2	1.95	0.48
1:A:221:HIS:O	1:A:222:GLN:C	2.56	0.48
2:B:65:LYS:HD2	2:B:72:ARG:HD2	1.96	0.48
1:A:21:VAL:HB	1:A:59:PRO:HD3	1.96	0.48
1:A:177:ASP:OD2	1:A:178:ILE:HG23	2.15	0.47
1:A:241:VAL:HG12	1:A:242:GLN:O	2.15	0.47
1:A:178:ILE:HG22	1:A:191:SER:HB3	1.98	0.46
2:B:173:LYS:O	2:B:173:LYS:HD2	2.15	0.46
1:A:376:THR:O	1:A:380[B]:ILE:HG12	2.16	0.46
2:B:235:HIS:HB3	2:B:238:LYS:HE3	1.98	0.46
1:A:338[B]:THR:HG23	7:A:753:HOH:O	2.16	0.45
1:A:356[B]:ARG:HD3	1:A:367:GLN:NE2	2.32	0.45
1:A:103:LYS:NZ	7:A:707:HOH:O	2.40	0.45
1:A:28:GLU:HG3	1:A:135:ILE:HD12	1.98	0.45
2:B:328:GLU:HG2	2:B:390:LYS:HD3	1.98	0.44
1:A:65:LYS:HD3	1:A:70:LYS:HE2	1.99	0.44
2:B:191:SER:OG	2:B:198:HIS:ND1	2.48	0.44
2:B:103:LYS:HD3	2:B:103:LYS:HA	1.66	0.44
2:B:229:TRP:CE2	2:B:230:MET:HG2	2.52	0.44
1:A:369:THR:O	1:A:373:GLN:HG2	2.18	0.44
1:A:236:PRO:HA	6:A:608[A]:K7F:C6	2.48	0.44
1:A:166:LYS:HD2	4:A:606:EDO:H11	1.98	0.44
1:A:261:VAL:HG22	1:A:265:ASN:ND2	2.33	0.43
1:A:542:ILE:O	1:A:546:GLU:HG2	2.19	0.43
1:A:424:LYS:CE	7:A:780:HOH:O	2.33	0.43
1:A:122:GLU:H	1:A:122:GLU:CD	2.27	0.42
1:A:357:MET:HE3	1:A:357:MET:HB2	1.82	0.42
2:B:121:ASP:O	2:B:125:ARG:HG3	2.20	0.42
2:B:195:ILE:O	2:B:195:ILE:CG1	2.66	0.42
2:B:197:GLN:HG3	2:B:200:THR:CG2	2.49	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:442:VAL:HB	1:A:481:ALA:HB1	2.02	0.42
2:B:60[B]:VAL:HG12	2:B:75:VAL:HG22	2.01	0.42
1:A:244:ILE:CG2	1:A:263:LYS:HD3	2.50	0.42
1:A:513:SER:N	1:A:519:ASN:OD1	2.52	0.41
1:A:199:ARG:O	1:A:202:ILE:HB	2.20	0.41
2:B:200:THR:O	2:B:204:GLU:N	2.52	0.41
1:A:363:ASN:HA	1:A:511:ASP:OD1	2.21	0.41
2:B:68:SER:HB2	2:B:230:MET:CE	2.50	0.41
2:B:21:VAL:HB	2:B:59:PRO:HD3	2.03	0.41
1:A:334:GLN:HB3	3:A:603:SO4:O3	2.21	0.41
1:A:405:TYR:CE2	1:A:407:GLN:HB2	2.57	0.40
2:B:394[A]:GLN:NE2	7:B:639:HOH:O	2.54	0.40
1:A:253:THR:O	1:A:257:ILE:HG13	2.21	0.40
1:A:476:LYS:HE3	1:A:476:LYS:HB3	1.59	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	559/557 (100%)	545 (98%)	14 (2%)	0	100	100
2	B	417/429 (97%)	405 (97%)	11 (3%)	1 (0%)	43	40
All	All	976/986 (99%)	950 (97%)	25 (3%)	1 (0%)	48	45

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	89	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	499/495 (101%)	499 (100%)	0	100	100
2	B	381/390 (98%)	380 (100%)	1 (0%)	86	86
All	All	880/885 (99%)	879 (100%)	1 (0%)	88	89

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	90	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	394	GLN
1	A	509	GLN
2	B	147	ASN
2	B	151	GLN
2	B	208	HIS
2	B	269	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 13 ligands modelled in this entry, 1 is monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	A	601	-	4,4,4	0.23	0	6,6,6	0.08	0
4	EDO	B	504	-	3,3,3	0.42	0	2,2,2	0.36	0
3	SO4	A	604	-	4,4,4	0.25	0	6,6,6	0.12	0
3	SO4	B	502	-	4,4,4	0.24	0	6,6,6	0.07	0
4	EDO	A	606	-	3,3,3	0.43	0	2,2,2	0.38	0
4	EDO	B	505	-	3,3,3	0.42	0	2,2,2	0.36	0
3	SO4	A	602	-	4,4,4	0.24	0	6,6,6	0.09	0
3	SO4	A	603	-	4,4,4	0.24	0	6,6,6	0.12	0
4	EDO	B	503	-	3,3,3	0.41	0	2,2,2	0.37	0
4	EDO	A	605	-	3,3,3	0.42	0	2,2,2	0.37	0
3	SO4	B	501	-	4,4,4	0.23	0	6,6,6	0.08	0
6	K7F	A	608[A]	-	44,44,44	0.15	0	58,61,61	1.02	2 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	B	504	-	-	0/1/1/1	-
4	EDO	A	606	-	-	0/1/1/1	-
4	EDO	B	505	-	-	0/1/1/1	-
4	EDO	B	503	-	-	0/1/1/1	-
4	EDO	A	605	-	-	0/1/1/1	-
6	K7F	A	608[A]	-	-	4/22/32/32	0/5/5/5

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	608[A]	K7F	C26-C24-C14	5.37	128.74	122.76
6	A	608[A]	K7F	C13-N2-C12	-4.79	116.88	124.32

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	A	608[A]	K7F	C25-C24-C26-C27
6	A	608[A]	K7F	C25-C24-C26-C30
6	A	608[A]	K7F	C14-C24-C26-C27
6	A	608[A]	K7F	C14-C24-C26-C30

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	502	SO4	1	0
4	A	606	EDO	1	0
3	A	603	SO4	2	0
6	A	608[A]	K7F	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	556/557 (99%)	0.73	78 (14%) 6 5	12, 47, 101, 160	5 (0%)
2	B	414/429 (96%)	0.59	54 (13%) 7 6	13, 43, 99, 162	7 (1%)
All	All	970/986 (98%)	0.67	132 (13%) 7 6	12, 45, 100, 162	12 (1%)

All (132) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	92	LEU	6.3
1	A	544	GLY	6.0
1	A	543	GLY	4.9
1	A	552	VAL	4.8
2	B	90	VAL	4.6
2	B	4	PRO	4.5
2	B	95	PRO	4.3
1	A	260	LEU	4.2
1	A	114	ALA	4.1
1	A	283	LEU	3.9
1	A	315[A]	HIS	3.9
2	B	195	ILE	3.9
1	A	244	ILE	3.8
2	B	315	HIS	3.6
2	B	5	ILE	3.6
2	B	91	GLN	3.6
2	B	230	MET	3.6
1	A	257	ILE	3.5
1	A	92	LEU	3.5
2	B	228	LEU	3.5
1	A	227	PHE	3.5
1	A	195	ILE	3.4
2	B	87	PHE	3.4
2	B	196	GLY	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	549	ASP	3.4
2	B	209	LEU	3.3
1	A	109	LEU	3.2
2	B	88	TRP	3.2
2	B	202	ILE	3.1
1	A	541	GLY	3.1
1	A	546	GLU	3.1
2	B	94	ILE	2.9
1	A	290	THR	2.9
1	A	116	PHE	2.9
2	B	98	ALA	2.9
1	A	256	ASP	2.9
2	B	212	TRP	2.9
2	B	361	HIS	2.9
1	A	314	VAL	2.8
1	A	71	TRP	2.8
1	A	254	VAL	2.8
1	A	261	VAL	2.8
2	B	67	ASP	2.8
1	A	183	TYR	2.8
1	A	295	LEU	2.8
1	A	292	VAL	2.8
1	A	542	ILE	2.8
1	A	547	GLN	2.8
2	B	316	GLY	2.8
2	B	229	TRP	2.8
1	A	251	SER	2.7
2	B	198	HIS	2.7
2	B	225	PRO	2.7
2	B	66	LYS	2.7
2	B	193	LEU	2.7
1	A	-1	MET	2.7
1	A	112	GLY	2.7
2	B	7	THR	2.7
1	A	38	CYS	2.7
1	A	286	THR	2.6
2	B	227	PHE	2.6
1	A	24	TRP	2.6
1	A	288	ALA	2.6
1	A	90	VAL	2.6
1	A	271	TYR	2.6
1	A	203	GLU	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	223	LYS	2.6
1	A	550	LYS	2.6
2	B	171	PHE	2.5
1	A	122	GLU	2.5
1	A	245	VAL	2.5
2	B	241	VAL	2.5
2	B	93	GLY	2.5
2	B	240	THR	2.5
2	B	360	ALA	2.5
1	A	289	LEU	2.5
2	B	283	LEU	2.5
2	B	168	LEU	2.5
1	A	548	VAL	2.5
2	B	232	TYR	2.4
2	B	226	PRO	2.4
2	B	284	ARG	2.4
2	B	358	ARG	2.4
1	A	40	GLU	2.4
1	A	200	THR	2.4
1	A	294	PRO	2.4
1	A	554	ALA	2.4
1	A	187	LEU	2.4
2	B	359	GLY	2.4
2	B	176	PRO	2.4
2	B	210	LEU	2.3
2	B	234	LEU	2.3
1	A	70	LYS	2.3
2	B	69	THR	2.3
1	A	310	LEU	2.3
1	A	293	ILE	2.3
1	A	217	PRO	2.3
1	A	243	PRO	2.3
1	A	231	GLY	2.3
1	A	234	LEU	2.3
2	B	237	ASP	2.3
1	A	553	SER	2.3
1	A	202	ILE	2.3
2	B	286[A]	THR	2.3
1	A	218	ASP	2.3
1	A	246	LEU	2.3
1	A	197	GLN	2.2
1	A	115	TYR	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	65	LYS	2.2
2	B	178	ILE	2.2
1	A	299	ALA	2.2
2	B	197	GLN	2.2
2	B	355	ALA	2.2
1	A	551	LEU	2.2
1	A	252	TRP	2.2
1	A	113	ASP	2.2
1	A	106	VAL	2.1
1	A	221	HIS	2.1
2	B	68	SER	2.1
2	B	205	LEU	2.1
2	B	318	TYR	2.1
1	A	25	PRO	2.1
1	A	226	PRO	2.1
1	A	37	ILE	2.1
1	A	241	VAL	2.1
1	A	199	ARG	2.0
1	A	216	THR	2.0
1	A	240	THR	2.0
1	A	279	LEU	2.0
2	B	192	ASP	2.0
1	A	356[A]	ARG	2.0
2	B	213	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SO4	A	603	5/5	0.42	0.14	113,115,123,146	0
3	SO4	A	601	5/5	0.74	0.15	64,69,91,95	0
3	SO4	A	604	5/5	0.75	0.12	70,89,108,110	0
3	SO4	A	602	5/5	0.78	0.10	74,87,109,114	0
6	K7F	A	608[A]	40/40	0.79	0.18	37,58,80,81	0
3	SO4	B	501	5/5	0.83	0.09	63,79,96,99	0
4	EDO	B	505	4/4	0.84	0.18	63,64,66,92	0
3	SO4	B	502	5/5	0.86	0.10	55,67,80,100	0
4	EDO	A	605	4/4	0.89	0.12	34,37,37,42	0
4	EDO	B	503	4/4	0.89	0.12	35,39,43,56	0
4	EDO	A	606	4/4	0.90	0.17	55,57,63,64	0
4	EDO	B	504	4/4	0.92	0.18	36,44,59,71	0
5	MG	A	607	1/1	0.97	0.06	64,64,64,64	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.