



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 5, 2026 – 09:10 PM EDT

PDB ID : 3KX3 / pdb_00003kx3
Title : Crystal structure of Bacillus megaterium BM3 heme domain mutant L86E
Authors : Girvan, H.M.; Levy, C.W.; Leys, D.; Munro, A.W.
Deposited on : 2009-12-02
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.015 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	470	
1	B	470	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 8768 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Bifunctional P-450/NADPH-P450 reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	454	Total	C	N	O	S	0	4	0
			3669	2344	624	684	17			
1	B	456	Total	C	N	O	S	0	9	0
			3738	2380	638	703	17			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	86	GLU	LEU	engineered mutation	UNP P14779
B	86	GLU	LEU	engineered mutation	UNP P14779

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is N-PALMITOYLGLYCINE (CCD ID: 140) (formula: $C_{18}H_{35}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			22	18	1	3		
3	B	1	Total	C	N	O	0	0
			22	18	1	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	613	Total	O	0	0
			613	613		
4	B	618	Total	O	0	0
			618	618		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Bifunctional P-450/NADPH-P450 reductase



- Molecule 1: Bifunctional P-450/NADPH-P450 reductase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	58.96Å 148.25Å 64.21Å 90.00° 98.55° 90.00°	Depositor
Resolution (Å)	39.41 – 1.80 39.41 – 1.80	Depositor EDS
% Data completeness (in resolution range)	96.2 (39.41-1.80) 96.2 (39.41-1.80)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.43 (at 1.81Å)	Xtriage
Refinement program	PHENIX (phenix.refine)	Depositor
R, R_{free}	0.144 , 0.195 0.141 , 0.191	Depositor DCC
R_{free} test set	4812 reflections (4.81%)	wwPDB-VP
Wilson B-factor (Å ²)	17.6	Xtriage
Anisotropy	0.315	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 52.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8768	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, 140

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.36	7/3756 (0.2%)	1.25	0/5077
1	B	1.33	5/3826 (0.1%)	1.23	10/5171 (0.2%)
All	All	1.35	12/7582 (0.2%)	1.24	10/10248 (0.1%)

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	286	VAL	CA-CB	8.03	1.64	1.54
1	A	297	VAL	CA-CB	6.84	1.63	1.54
1	A	412	LEU	C-O	6.21	1.31	1.24
1	A	438	THR	N-CA	6.12	1.54	1.45
1	B	354	MET	SD-CE	-5.88	1.64	1.79
1	A	330	ALA	CA-CB	5.70	1.63	1.53
1	B	269	THR	CA-CB	5.70	1.62	1.53
1	A	42	PHE	C-O	5.52	1.30	1.24
1	B	69	LYS	C-O	5.43	1.30	1.23
1	A	268	THR	C-O	5.39	1.30	1.24
1	B	114	GLY	C-O	-5.34	1.17	1.24
1	A	366	ILE	CA-C	-5.03	1.45	1.52

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	192	ASN	CA-C-N	7.00	126.70	119.56
1	B	192	ASN	C-N-CA	7.00	126.70	119.56
1	B	378	ARG	CB-CA-C	-5.45	99.89	110.46
1	B	96	TRP	CA-CB-CG	-5.37	103.40	113.60
1	B	96	TRP	N-CA-CB	5.33	117.68	109.91
1	B	297	VAL	N-CA-C	5.26	116.74	111.00
1	B	190	ARG	N-CA-C	5.22	116.76	108.67

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	453	ILE	CA-C-O	5.21	122.22	119.15
1	B	5	MET	CA-C-N	5.19	125.47	119.92
1	B	5	MET	C-N-CA	5.19	125.47	119.92

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3669	0	3624	22	0
1	B	3738	0	3680	32	0
2	A	43	0	30	0	0
2	B	43	0	30	2	0
3	A	22	0	34	3	0
3	B	22	0	34	1	0
4	A	613	0	0	9	0
4	B	618	0	0	16	0
All	All	8768	0	7432	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:1470:140:O2	4:B:566:HOH:O	2.01	0.78
1:A:77:PHE:CE1	1:A:187:LYS:HE2	2.19	0.76
3:A:1470:140:O2	4:A:660:HOH:O	2.04	0.76
1:B:236:HIS:CD2	4:B:1252:HOH:O	2.44	0.70
1:A:422:ASP:HB2	1:A:449:LYS:HG3	1.73	0.70
1:B:189:GLN:HB3	4:B:1191:HOH:O	1.90	0.70
1:A:189:GLN:CG	4:A:1066:HOH:O	2.46	0.62
1:B:41:LYS:HE2	1:B:48:VAL:HG21	1.84	0.60
1:A:188:LEU:HD13	3:A:1470:140:HN	1.68	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:31[B]:LYS:HG3	4:A:604:HOH:O	2.04	0.58
1:B:375:ARG:HD3	4:B:761:HOH:O	2.04	0.57
1:B:97:LYS:HD2	4:B:1249:HOH:O	2.05	0.56
1:B:375:ARG:NE	4:B:761:HOH:O	2.36	0.56
1:B:167:ARG:NH1	4:B:737:HOH:O	2.28	0.55
1:A:189:GLN:HG2	4:A:1066:HOH:O	2.07	0.54
1:A:189:GLN:HG3	4:A:1066:HOH:O	2.06	0.54
1:B:382:PRO:HG3	4:B:894:HOH:O	2.08	0.53
1:B:388:HIS:HA	1:B:391:LYS:HD3	1.90	0.53
1:A:104:LEU:HB3	1:A:105:PRO:HD3	1.91	0.52
1:B:51:TYR:CE2	1:B:354:MET:HG2	2.45	0.52
1:B:367:TRP:HB2	1:B:371:VAL:HG12	1.91	0.51
1:B:81:PHE:HB3	1:B:209:ILE:HG12	1.92	0.51
1:B:186[B]:ASN:O	1:B:189:GLN:HG2	2.12	0.50
1:B:243:PRO:HB2	4:B:1093:HOH:O	2.12	0.49
1:A:77:PHE:CZ	1:A:187:LYS:HE2	2.47	0.49
1:B:375:ARG:CD	4:B:761:HOH:O	2.59	0.49
1:A:458:ILE:HG23	4:A:1186:HOH:O	2.13	0.49
1:B:186[A]:ASN:O	1:B:189:GLN:HG2	2.14	0.48
1:A:306:LYS:HG2	4:A:479:HOH:O	2.14	0.47
1:B:236:HIS:CG	4:B:1252:HOH:O	2.64	0.47
1:A:110:GLN:NE2	4:A:640:HOH:O	2.47	0.46
1:B:151:ASP:OD1	1:B:162:PHE:HB2	2.15	0.46
1:B:148:LEU:HD21	1:B:413:VAL:HG21	1.97	0.45
1:A:17:LEU:HB3	1:A:18:PRO:HD3	1.97	0.45
2:B:471:HEM:HBC2	2:B:471:HEM:HMC2	1.99	0.45
1:B:300:ASP:OD2	4:B:755:HOH:O	2.21	0.44
1:B:458:ILE:HG23	4:B:798:HOH:O	2.17	0.44
1:B:272:LEU:HD13	1:B:322:LEU:HG	1.99	0.44
1:A:212:MET:O	1:A:216:VAL:HG23	2.18	0.44
1:A:178:VAL:O	1:A:182:ASP:HB2	2.18	0.43
1:B:5:MET:HE3	1:B:5:MET:HB2	1.77	0.43
1:A:47:ARG:CZ	1:A:73:GLN:HG3	2.48	0.43
2:B:471:HEM:HBC2	2:B:471:HEM:CMC	2.48	0.43
1:B:300:ASP:HB3	1:B:301:PRO:CD	2.49	0.43
1:A:216:VAL:HG21	1:A:259:ILE:HG13	2.01	0.43
1:A:141:VAL:HB	1:A:142:PRO:HD3	2.01	0.42
1:A:453:ILE:HA	1:A:454:PRO:HD3	1.89	0.42
1:A:73:GLN:HB2	3:A:1470:140:O2	2.20	0.42
1:B:205:PHE:CE2	1:B:209:ILE:HD11	2.55	0.42
1:B:169:GLN:NE2	4:B:1045:HOH:O	2.53	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:426:HIS:CD2	1:B:447:LYS:HE3	2.54	0.42
1:B:97:LYS:HB2	4:B:703:HOH:O	2.19	0.42
1:B:104:LEU:N	1:B:105:PRO:CD	2.83	0.42
1:A:229:GLN:N	4:A:911:HOH:O	2.53	0.41
1:B:312:LYS:NZ	4:B:644:HOH:O	2.54	0.41
1:B:280:LEU:HA	1:B:280:LEU:HD23	1.84	0.40
1:B:233:LEU:HG	1:B:237:MET:HE3	2.03	0.40
1:A:5:MET:SD	1:A:50:ARG:HG2	2.62	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	454/470 (97%)	445 (98%)	9 (2%)	0	100	100
1	B	463/470 (98%)	454 (98%)	9 (2%)	0	100	100
All	All	917/940 (98%)	899 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	396/411 (96%)	394 (100%)	2 (0%)	81	80

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	405/411 (98%)	399 (98%)	6 (2%)	57	49
All	All	801/822 (97%)	793 (99%)	8 (1%)	68	64

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	148	LEU
1	A	440	LYS
1	B	97	LYS
1	B	104	LEU
1	B	148	LEU
1	B	200	GLU
1	B	248	PRO
1	B	440	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	70	ASN
1	A	201	ASN
1	A	359	GLN
1	A	403	GLN
1	B	27	GLN
1	B	70	ASN
1	B	73	GLN
1	B	201	ASN
1	B	239	ASN
1	B	359	GLN
1	B	403	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	HEM	A	471	1,4	50,50,50	2.01	17 (34%)	67,82,82	2.59	19 (28%)
2	HEM	B	471	1,4	50,50,50	2.02	15 (30%)	67,82,82	2.72	20 (29%)
3	140	A	1470	-	21,21,21	0.55	0	22,22,22	1.19	1 (4%)
3	140	B	1470	-	21,21,21	0.52	0	22,22,22	1.22	3 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	471	1,4	-	4/14/54/54	-
2	HEM	B	471	1,4	-	2/14/54/54	-
3	140	A	1470	-	-	5/20/20/20	-
3	140	B	1470	-	-	8/20/20/20	-

All (32) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	471	HEM	FE-ND	5.34	2.11	1.94
2	B	471	HEM	FE-NB	5.03	2.10	1.94
2	B	471	HEM	C3D-C2D	4.26	1.45	1.36
2	A	471	HEM	FE-ND	4.14	2.07	1.94
2	A	471	HEM	FE-NB	3.98	2.07	1.94
2	A	471	HEM	C1B-NB	-3.82	1.33	1.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	471	HEM	CMD-C2D	3.73	1.58	1.50
2	B	471	HEM	CMC-C2C	3.68	1.58	1.50
2	A	471	HEM	C3D-C2D	3.61	1.44	1.36
2	B	471	HEM	CAB-C3B	3.61	1.57	1.47
2	A	471	HEM	FE-NC	3.56	2.06	1.95
2	B	471	HEM	CBD-CGD	3.46	1.58	1.50
2	A	471	HEM	C4D-C3D	3.09	1.50	1.45
2	A	471	HEM	CMA-C3A	3.08	1.57	1.50
2	B	471	HEM	CMD-C2D	3.02	1.57	1.50
2	A	471	HEM	C4D-ND	-2.90	1.35	1.40
2	B	471	HEM	C4D-ND	-2.67	1.35	1.40
2	A	471	HEM	CAB-C3B	2.65	1.54	1.47
2	B	471	HEM	CHA-C4D	-2.56	1.33	1.38
2	A	471	HEM	CMC-C2C	2.53	1.56	1.50
2	A	471	HEM	CHA-C4D	-2.46	1.33	1.38
2	A	471	HEM	CBD-CGD	2.42	1.56	1.50
2	A	471	HEM	FE-NA	2.40	2.03	1.95
2	B	471	HEM	CAC-C3C	2.34	1.53	1.47
2	B	471	HEM	FE-NC	2.34	2.02	1.95
2	B	471	HEM	C3B-C4B	2.32	1.49	1.44
2	A	471	HEM	CAD-C3D	2.27	1.57	1.51
2	B	471	HEM	CAD-C3D	2.25	1.57	1.51
2	B	471	HEM	CMA-C3A	2.22	1.55	1.50
2	A	471	HEM	CMB-C2B	2.22	1.55	1.50
2	A	471	HEM	CBA-CGA	2.08	1.55	1.50
2	B	471	HEM	C1B-C2B	2.05	1.48	1.44

All (43) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	471	HEM	C4D-ND-C1D	8.99	115.86	105.21
2	A	471	HEM	C4D-ND-C1D	7.61	114.21	105.21
2	B	471	HEM	C1B-NB-C4B	7.48	114.07	105.21
2	A	471	HEM	CHD-C1D-ND	7.29	132.27	124.42
2	A	471	HEM	CHC-C4B-NB	6.91	131.86	124.42
2	A	471	HEM	C3B-C4B-NB	-6.63	104.71	109.47
2	A	471	HEM	C1B-NB-C4B	6.54	112.95	105.21
2	B	471	HEM	CHB-C1B-NB	6.45	132.34	124.37
2	B	471	HEM	CHA-C4D-ND	6.23	132.07	124.37
2	B	471	HEM	CHD-C1D-ND	6.21	131.10	124.42
2	B	471	HEM	C3B-C4B-NB	-5.56	105.48	109.47
2	A	471	HEM	CHA-C4D-ND	5.45	131.11	124.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	471	HEM	C3D-C4D-ND	-5.45	104.19	110.17
2	B	471	HEM	C4C-C3C-C2C	4.77	110.95	106.81
2	B	471	HEM	C2B-C1B-NB	-4.52	104.65	109.84
2	B	471	HEM	CBD-CAD-C3D	-4.12	101.14	112.53
2	A	471	HEM	C4C-C3C-C2C	3.87	110.17	106.81
2	A	471	HEM	CBD-CAD-C3D	-3.83	101.94	112.53
2	B	471	HEM	C2D-C1D-ND	-3.77	105.55	109.90
2	A	471	HEM	C3B-C2B-C1B	3.66	109.16	106.41
2	A	471	HEM	C2D-C1D-ND	-3.65	105.68	109.90
2	A	471	HEM	C2B-C1B-NB	-3.52	105.79	109.84
2	A	471	HEM	C3D-C4D-ND	-3.40	106.44	110.17
2	A	471	HEM	CHB-C1B-NB	3.21	128.34	124.37
3	A	1470	140	O2-C1-C2	3.03	124.35	112.81
2	B	471	HEM	CHC-C4B-NB	2.93	127.58	124.42
2	B	471	HEM	CMD-C2D-C1D	2.81	129.43	125.03
2	B	471	HEM	CHD-C4C-NC	2.75	127.44	124.45
2	B	471	HEM	C4C-CHD-C1D	-2.64	120.41	126.02
2	A	471	HEM	C4C-CHD-C1D	-2.54	120.62	126.02
2	A	471	HEM	CBA-CAA-C2A	-2.47	105.70	112.53
2	A	471	HEM	CHD-C4C-NC	2.42	127.09	124.45
2	B	471	HEM	C4A-CHB-C1B	-2.33	120.77	126.25
2	B	471	HEM	C3B-C2B-C1B	2.25	108.10	106.41
2	A	471	HEM	CAD-C3D-C4D	2.24	128.61	124.70
2	A	471	HEM	CHC-C1C-NC	2.24	126.89	124.45
3	B	1470	140	C5-C4-C3	-2.20	107.09	113.19
2	B	471	HEM	CBA-CAA-C2A	-2.20	106.46	112.53
2	A	471	HEM	CHB-C4A-NA	2.18	127.81	123.86
3	B	1470	140	O3-C3-N	-2.17	118.78	123.03
2	B	471	HEM	C3C-C2C-C1C	-2.13	105.03	107.05
2	B	471	HEM	CMC-C2C-C1C	2.11	128.44	124.73
3	B	1470	140	C14-C13-C12	-2.01	104.22	114.37

There are no chirality outliers.

All (19) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	1470	140	C7-C8-C9-C10
3	A	1470	140	C9-C10-C11-C12
3	A	1470	140	C7-C8-C9-C10
3	B	1470	140	C9-C10-C11-C12
3	A	1470	140	C10-C11-C12-C13
3	A	1470	140	C11-C12-C13-C14

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	B	1470	140	N-C3-C4-C5
3	B	1470	140	O3-C3-C4-C5
3	B	1470	140	C11-C12-C13-C14
2	B	471	HEM	CAD-CBD-CGD-O1D
2	A	471	HEM	CAD-CBD-CGD-O2D
2	B	471	HEM	CAD-CBD-CGD-O2D
3	B	1470	140	O2-C1-C2-N
2	A	471	HEM	CAD-CBD-CGD-O1D
2	A	471	HEM	CAA-CBA-CGA-O1A
3	B	1470	140	C6-C7-C8-C9
3	A	1470	140	O3-C3-C4-C5
2	A	471	HEM	CAA-CBA-CGA-O2A
3	B	1470	140	C5-C6-C7-C8




There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	471	HEM	2	0
3	A	1470	140	3	0
3	B	1470	140	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	454/470 (96%)	-0.43	4 (0%) 81 81	7, 19, 39, 54	4 (0%)
1	B	456/470 (97%)	-0.50	0 100 100	8, 19, 39, 64	9 (1%)
All	All	910/940 (96%)	-0.46	4 (0%) 88 89	7, 19, 39, 64	13 (1%)

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	192	ASN	2.8
1	A	457	GLY	2.3
1	A	197	ALA	2.3
1	A	381	ASN	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	140	B	1470	22/22	0.85	0.12	21,35,51,58	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	140	A	1470	22/22	0.87	0.12	21,34,51,56	0
2	HEM	A	471	43/43	0.99	0.04	7,9,14,16	0
2	HEM	B	471	43/43	0.99	0.03	8,10,14,16	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 140 B 1470:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 140 Å 1470:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM A 471:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.