



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 08:31 PM UTC

PDB ID : 1M4G / pdb_00001m4g
Title : Aminoglycoside 2'-N-acetyltransferase from Mycobacterium tuberculosis-Complex with Coenzyme A and Ribostamycin
Authors : Vetting, M.W.; Hegde, S.S.; Javid-Majd, F.; Blanchard, J.S.; Roderick, S.L.
Deposited on : 2002-07-02
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	181	 2% 88% 12%
1	B	181	 4% 89% 6% . .

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 3314 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aminoglycoside 2'-N-acetyltransferase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	181	Total	C	N	O	S	0	0	0
			1412	884	265	256	7			
1	B	176	Total	C	N	O	S	0	0	0
			1371	859	257	249	6			

- Molecule 2 is COENZYME A (CCD ID: COA) (formula: $C_{21}H_{36}N_7O_{16}P_3S$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P S	0	0
			48	21	7	16	3 1		
2	B	1	Total	C	N	O	P S	0	0
			48	21	7	16	3 1		

- Molecule 3 is RIBOSTAMYCIN (CCD ID: RIO) (formula: $C_{17}H_{34}N_4O_{10}$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total 31	C 17	N 4	O 10	0	0
3	B	1	Total 31	C 17	N 4	O 10	0	0

- Molecule 4 is 3'-PHOSPHATE-ADENOSINE-5'-DIPHOSPHATE (CCD ID: PAP) (formula: $\text{C}_{10}\text{H}_{16}\text{N}_5\text{O}_{13}\text{P}_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	N	O	P	0	0
			31	10	5	13	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	N	O	P	0	0
			31	10	5	13	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	159	Total	O	0	0
			159	159		
5	B	152	Total	O	0	0
			152	152		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Aminoglycoside 2'-N-acetyltransferase



- Molecule 1: Aminoglycoside 2'-N-acetyltransferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	48.70Å 86.40Å 98.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.80 50.00 – 1.80	Depositor EDS
% Data completeness (in resolution range)	98.0 (50.00-1.80) 98.0 (50.00-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.60 (at 1.80Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.165 , 0.202 0.165 , 0.202	Depositor DCC
R_{free} test set	1922 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	16.0	Xtriage
Anisotropy	0.085	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 45.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3314	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: RIO, COA, PAP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.19	4/1447 (0.3%)	1.23	9/1971 (0.5%)
1	B	1.13	1/1405 (0.1%)	1.21	7/1914 (0.4%)
All	All	1.16	5/2852 (0.2%)	1.22	16/3885 (0.4%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	107	MET	CB-CG	8.32	1.77	1.52
1	A	172	MET	SD-CE	-5.55	1.65	1.79
1	A	149	THR	CA-CB	5.39	1.58	1.53
1	A	75	ALA	CA-CB	5.09	1.59	1.52
1	A	114	GLY	CA-C	5.02	1.57	1.52

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	107	MET	CB-CG-SD	-11.51	78.17	112.70
1	B	7	THR	N-CA-C	-9.53	94.45	109.50
1	A	88	ALA	N-CA-C	7.57	120.41	111.71
1	A	118	SER	N-CA-C	6.23	120.05	110.20
1	B	118	SER	N-CA-C	5.95	119.96	109.96
1	A	86	VAL	N-CA-C	-5.93	99.81	108.11
1	B	119	SER	N-CA-C	-5.46	102.81	110.50
1	A	119	SER	N-CA-C	-5.44	103.35	110.53
1	A	3	THR	N-CA-C	5.40	118.97	112.38
1	A	163	ILE	N-CA-C	-5.33	100.18	108.86
1	A	7	THR	N-CA-C	-5.28	101.86	110.20
1	A	69	LEU	N-CA-C	-5.13	100.54	108.90
1	B	88	ALA	N-CA-C	5.13	117.53	111.33
1	A	79	GLY	N-CA-C	-5.13	101.03	113.18

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	107	MET	CG-SD-CE	-5.09	89.70	100.90
1	B	166	ASP	N-CA-C	-5.01	99.18	107.99

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1412	0	1374	9	0
1	B	1371	0	1331	25	0
2	A	48	0	32	1	1
2	B	48	0	32	0	0
3	A	31	0	34	0	0
3	B	31	0	34	0	0
4	B	62	0	22	0	0
5	A	159	0	0	2	0
5	B	152	0	0	8	1
All	All	3314	0	2859	34	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:107:MET:CG	1:B:107:MET:CB	1.77	1.57
1:B:107:MET:CB	1:B:107:MET:SD	2.31	1.18
1:B:9:ARG:HG2	1:B:9:ARG:HH11	1.01	1.07
1:B:9:ARG:HG2	1:B:9:ARG:NH1	1.77	0.96
1:B:163:ILE:HD13	1:B:163:ILE:O	1.67	0.94
1:B:107:MET:SD	1:B:107:MET:HB2	2.04	0.93
1:B:163:ILE:H	1:B:163:ILE:HD12	1.31	0.93
1:A:2:HIS:HD2	1:A:4:GLN:H	1.14	0.92
1:B:9:ARG:HH11	1:B:9:ARG:CG	1.85	0.90

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:163:ILE:HD12	1:B:163:ILE:N	1.90	0.85
1:A:2:HIS:CD2	1:A:4:GLN:H	1.97	0.83
1:B:163:ILE:O	5:B:718:HOH:O	2.11	0.69
1:B:163:ILE:H	1:B:163:ILE:CD1	2.04	0.67
1:B:38:GLU:OE2	5:B:700:HOH:O	2.11	0.67
1:B:163:ILE:N	1:B:163:ILE:CD1	2.55	0.67
1:B:170:GLU:HG3	5:B:714:HOH:O	1.95	0.66
1:B:164:SER:HA	5:B:718:HOH:O	1.99	0.63
1:B:124:ARG:HG3	5:B:635:HOH:O	2.01	0.61
1:A:39:THR:HG21	1:A:178:GLY:HA3	1.89	0.55
1:A:18:SER:O	1:A:22:GLN:HG3	2.09	0.53
1:B:124:ARG:NH2	5:B:753:HOH:O	2.30	0.49
1:B:162:ASP:OD2	5:B:734:HOH:O	2.20	0.48
1:B:163:ILE:O	1:B:163:ILE:CD1	2.52	0.47
1:B:124:ARG:NE	5:B:753:HOH:O	2.41	0.47
1:A:21:ARG:NH1	5:A:619:HOH:O	2.40	0.47
1:B:107:MET:CG	1:B:107:MET:CA	2.82	0.45
1:B:163:ILE:HD13	1:B:163:ILE:C	2.40	0.44
1:A:121:ARG:NH2	5:A:655:HOH:O	2.51	0.43
1:A:2:HIS:CD2	1:A:4:GLN:N	2.78	0.42
1:A:65:ILE:C	1:A:65:ILE:HD12	2.45	0.42
1:B:24:ILE:CG1	1:B:58:ILE:HD13	2.50	0.41
1:A:121:ARG:HH22	2:A:502:COA:H71	1.86	0.41
1:B:24:ILE:HG12	1:B:58:ILE:HD13	2.01	0.41
1:B:9:ARG:NH1	1:B:9:ARG:CG	2.55	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:502:COA:N6A	5:B:725:HOH:O[2_564]	2.12	0.08

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	179/181 (99%)	178 (99%)	1 (1%)	0	100	100
1	B	174/181 (96%)	172 (99%)	2 (1%)	0	100	100
All	All	353/362 (98%)	350 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	143/143 (100%)	143 (100%)	0	100	100
1	B	138/143 (96%)	135 (98%)	3 (2%)	45	34
All	All	281/286 (98%)	278 (99%)	3 (1%)	65	60

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	9	ARG
1	B	163	ILE
1	B	164	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2	HIS
1	A	22	GLN
1	A	135	HIS
1	B	22	GLN
1	B	66	GLN
1	B	135	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	RIO	B	501	-	32,33,33	1.18	3 (9%)	46,49,49	1.37	6 (13%)
4	PAP	B	600	-	32,33,33	1.63	5 (15%)	50,52,52	1.75	9 (18%)
2	COA	B	503	-	47,50,50	1.51	8 (17%)	69,75,75	1.65	12 (17%)
4	PAP	B	601	-	32,33,33	1.58	6 (18%)	50,52,52	1.79	9 (18%)
2	COA	A	502	-	47,50,50	1.41	6 (12%)	69,75,75	1.69	10 (14%)
3	RIO	A	500	-	32,33,33	1.12	4 (12%)	46,49,49	1.38	7 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	RIO	B	501	-	-	5/12/68/68	0/3/3/3
4	PAP	B	600	-	-	1/21/37/37	0/3/3/3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	COA	B	503	-	-	10/48/64/64	0/3/3/3
4	PAP	B	601	-	-	3/21/37/37	0/3/3/3
2	COA	A	502	-	-	7/48/64/64	0/3/3/3
3	RIO	A	500	-	-	5/12/68/68	0/3/3/3

All (32) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	600	PAP	P-O3'	5.77	1.69	1.59
4	B	601	PAP	P-O3'	4.85	1.68	1.59
2	B	503	COA	C4A-N3A	4.26	1.42	1.34
2	B	503	COA	P3B-O3B	4.04	1.66	1.59
2	A	502	COA	C9P-N8P	3.67	1.42	1.33
2	A	502	COA	CEP-CBP	3.51	1.61	1.53
2	B	503	COA	C2A-N1A	3.14	1.39	1.33
4	B	601	PAP	C2-N1	3.14	1.39	1.33
4	B	600	PAP	PB-O3B	3.03	1.66	1.54
2	A	502	COA	C7P-C6P	2.92	1.61	1.51
4	B	601	PAP	C5-C4	2.88	1.44	1.39
2	B	503	COA	P2A-O3A	-2.73	1.56	1.59
4	B	600	PAP	C5-C4	2.72	1.43	1.39
4	B	601	PAP	C4-N3	2.69	1.39	1.34
3	A	500	RIO	C7-C8	2.64	1.59	1.53
4	B	600	PAP	C4-N9	-2.63	1.32	1.37
4	B	601	PAP	PB-O3B	2.62	1.64	1.54
2	B	503	COA	CEP-CBP	2.60	1.59	1.53
4	B	600	PAP	C2-N1	2.56	1.38	1.33
3	B	501	RIO	C15-N4	2.56	1.51	1.47
2	A	502	COA	C4A-N3A	2.52	1.39	1.34
2	B	503	COA	P1A-O3A	2.40	1.62	1.59
2	A	502	COA	C8A-N9A	-2.36	1.33	1.37
3	A	500	RIO	C9-C4	2.33	1.57	1.52
3	B	501	RIO	C3-C16	-2.28	1.50	1.52
3	A	500	RIO	C15-N4	2.27	1.50	1.47
4	B	601	PAP	C6-N6	2.23	1.39	1.34
3	B	501	RIO	C14-C15	2.14	1.56	1.53
2	B	503	COA	OAP-CAP	2.11	1.46	1.42
2	B	503	COA	P1A-O2A	-2.07	1.45	1.55
3	A	500	RIO	O2-C3	2.07	1.45	1.41
2	A	502	COA	CCP-CBP	2.05	1.56	1.52

All (53) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	502	COA	C4A-N9A-C8A	6.09	112.13	105.74
4	B	600	PAP	C4-N9-C8	6.09	112.13	105.74
4	B	601	PAP	N3-C4-N9	5.99	137.36	127.17
2	B	503	COA	N3A-C4A-N9A	5.98	137.34	127.17
2	A	502	COA	N3A-C4A-N9A	5.82	137.06	127.17
2	B	503	COA	C4A-N9A-C8A	5.54	111.55	105.74
4	B	601	PAP	C4-N9-C8	4.95	110.93	105.74
3	A	500	RIO	O3-C3-O2	-4.83	106.44	111.37
4	B	600	PAP	N3-C4-N9	4.80	135.33	127.17
2	B	503	COA	C5A-C4A-N9A	-4.58	100.81	105.81
4	B	601	PAP	C5-C4-N9	-4.19	101.24	105.81
2	A	502	COA	C5A-C4A-N9A	-4.12	101.32	105.81
4	B	601	PAP	C5-C4-N3	-3.88	121.38	126.72
2	A	502	COA	N9A-C8A-N7A	-3.78	108.58	113.94
2	A	502	COA	C5A-C4A-N3A	-3.70	121.62	126.72
3	B	501	RIO	O3-C3-O2	-3.59	107.70	111.37
4	B	600	PAP	O2B-PB-O3A	3.55	116.53	104.64
2	B	503	COA	C5A-C4A-N3A	-3.54	121.84	126.72
4	B	600	PAP	C5-C4-N9	-3.44	102.06	105.81
4	B	600	PAP	N9-C8-N7	-3.24	109.34	113.94
3	B	501	RIO	O5-C9-C4	3.15	115.42	107.42
2	B	503	COA	N9A-C8A-N7A	-3.08	109.57	113.94
4	B	601	PAP	O4'-C1'-N9	3.03	113.90	108.09
4	B	600	PAP	C5-C4-N3	-2.98	122.61	126.72
3	A	500	RIO	C10-O6-C11	2.95	119.48	113.72
3	B	501	RIO	C10-O5-C9	2.91	124.87	117.98
4	B	600	PAP	C4-N9-C1'	-2.79	120.11	126.63
2	B	503	COA	CAP-C9P-N8P	-2.71	111.35	116.48
3	B	501	RIO	C10-O6-C11	2.60	118.79	113.72
4	B	600	PAP	C2'-C3'-C4'	-2.55	98.78	103.24
3	A	500	RIO	O5-C9-C8	-2.50	103.21	109.18
2	A	502	COA	C5A-C6A-N6A	-2.47	117.17	123.29
2	A	502	COA	C4A-N9A-C1B	-2.46	120.88	126.63
4	B	601	PAP	C5-C6-N6	-2.45	117.22	123.29
3	A	500	RIO	O3-C4-C9	2.43	113.59	107.42
2	A	502	COA	O3B-P3B-O7A	-2.42	100.69	109.33
4	B	601	PAP	N9-C8-N7	-2.42	110.51	113.94
2	B	503	COA	N6A-C6A-N1A	2.38	123.68	118.38
3	A	500	RIO	O5-C9-C4	2.35	113.38	107.42
4	B	601	PAP	N6-C6-N1	2.33	123.56	118.38
3	A	500	RIO	C10-O5-C9	2.32	123.47	117.98
2	A	502	COA	N6A-C6A-N1A	2.29	123.49	118.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	501	RIO	O3-C4-C5	2.26	112.97	107.23
2	B	503	COA	O4B-C1B-N9A	2.24	112.39	108.09
2	B	503	COA	C5A-C6A-N6A	-2.20	117.84	123.29
2	A	502	COA	O2A-P1A-O3A	2.18	113.16	107.27
3	A	500	RIO	C5-C6-N1	-2.11	106.77	110.94
2	B	503	COA	C6P-C5P-N4P	-2.08	112.55	116.34
3	B	501	RIO	C8-C7-C6	-2.08	107.46	111.02
4	B	600	PAP	O4'-C1'-N9	2.06	112.04	108.09
4	B	601	PAP	C6-C5-N7	-2.06	128.13	132.09
2	B	503	COA	C4A-N9A-C1B	-2.03	121.88	126.63
2	B	503	COA	C2B-C1B-N9A	2.01	118.30	113.30

There are no chirality outliers.

All (31) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	502	COA	CCP-O6A-P2A-O5A
2	B	503	COA	CCP-O6A-P2A-O3A
2	B	503	COA	CCP-O6A-P2A-O5A
4	B	601	PAP	O4'-C4'-C5'-O5'
4	B	601	PAP	C3'-C4'-C5'-O5'
3	B	501	RIO	C4-C9-O5-C10
3	A	500	RIO	C9-C4-O3-C3
3	A	500	RIO	C4-C9-O5-C10
3	B	501	RIO	C5-C4-O3-C3
3	B	501	RIO	C9-C4-O3-C3
3	A	500	RIO	C5-C4-O3-C3
3	A	500	RIO	O6-C10-O5-C9
2	A	502	COA	O5P-C5P-N4P-C3P
2	A	502	COA	C2B-C1B-N9A-C8A
2	B	503	COA	C6P-C7P-N8P-C9P
2	B	503	COA	C2B-C1B-N9A-C8A
2	B	503	COA	P2A-O3A-P1A-O1A
3	A	500	RIO	C8-C9-O5-C10
3	B	501	RIO	C8-C9-O5-C10
2	A	502	COA	CCP-O6A-P2A-O3A
2	A	502	COA	CCP-O6A-P2A-O4A
2	A	502	COA	C2B-C1B-N9A-C4A
2	B	503	COA	C3B-O3B-P3B-O8A
3	B	501	RIO	O6-C10-O5-C9
2	B	503	COA	C2B-C1B-N9A-C4A
2	A	502	COA	O4B-C1B-N9A-C8A

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	B	503	COA	O4B-C1B-N9A-C8A
4	B	600	PAP	C3'-O3'-P-O3
4	B	601	PAP	C3'-O3'-P-O2
2	B	503	COA	C3B-O3B-P3B-O7A
2	B	503	COA	P2A-O3A-P1A-O2A


There are no ring outliers.

1 monomer is involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	502	COA	1	1

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand RIO B 501




Bond lengths



Bond angles




Torsions




Rings


Ligand PAP B 600




Ligand COA B 503




Ligand PAP B 601



Ligand COA A 502





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	181/181 (100%)	-0.47	3 (1%) 69 69	9, 13, 27, 37	0
1	B	176/181 (97%)	-0.36	7 (3%) 42 41	8, 14, 33, 45	0
All	All	357/362 (98%)	-0.42	10 (2%) 55 55	8, 13, 30, 45	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	55	HIS	4.1
1	B	124	ARG	3.1
1	B	57	ALA	2.9
1	A	121	ARG	2.9
1	B	162	ASP	2.8
1	B	6	HIS	2.8
1	A	162	ASP	2.5
1	B	18	SER	2.5
1	A	1	MET	2.3
1	B	54	HIS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	PAP	B	601	31/31	0.77	0.14	39,46,54,55	0
4	PAP	B	600	31/31	0.91	0.11	22,32,38,40	0
3	RIO	A	500	31/31	0.96	0.06	11,14,27,29	0
2	COA	A	502	48/48	0.96	0.07	12,21,30,31	0
2	COA	B	503	48/48	0.96	0.10	15,21,55,56	0
3	RIO	B	501	31/31	0.97	0.05	8,12,19,22	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around PAP B 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PAP B 600:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around RIO A 500:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around COA A 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around COA B 503:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around RIO B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.