



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 07:15 AM UTC

PDB ID : 4MIK / pdb_00004mik
Title : Crystal Structure of hPNMT in Complex with bisubstrate inhibitor (2R,3R,4S,5S)-2-(6-amino-9H-purin-9-yl)-5-(((2-(((7-nitro-1,2,3,4-tetrahydroisoquinolin-3-yl)methyl)amino)ethyl)thio)methyl)tetrahydrofuran-3,4-diol
Authors : Bart, A.G.; Scott, E.E.
Deposited on : 2013-08-31
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	289	
1	B	289	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	TRS	A	302	-	X	X	-
3	TRS	B	302	-	X	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8617 atoms, of which 4125 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Phenylethanolamine N-methyltransferase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	255	Total	C	H	N	O	S	0	3	0
			4008	1280	1992	358	369	9			
1	B	265	Total	C	H	N	O	S	0	4	0
			4127	1315	2053	369	381	9			

There are 14 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	283	GLU	-	expression tag	UNP P11086
A	284	HIS	-	expression tag	UNP P11086
A	285	HIS	-	expression tag	UNP P11086
A	286	HIS	-	expression tag	UNP P11086
A	287	HIS	-	expression tag	UNP P11086
A	288	HIS	-	expression tag	UNP P11086
A	289	HIS	-	expression tag	UNP P11086
B	283	GLU	-	expression tag	UNP P11086
B	284	HIS	-	expression tag	UNP P11086
B	285	HIS	-	expression tag	UNP P11086
B	286	HIS	-	expression tag	UNP P11086
B	287	HIS	-	expression tag	UNP P11086
B	288	HIS	-	expression tag	UNP P11086
B	289	HIS	-	expression tag	UNP P11086

- Molecule 2 is 9-{5-S-[2-({[(3S)-7-nitro-1,2,3,4-tetrahydroisoquinolin-3-yl]methyl}amino)ethyl]-5-thio- α -L-lyxofuranosyl}-9H-purin-6-amine (CCD ID: JIL) (formula: C₂₂H₂₈N₈O₅S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total 64	C 22	H 28	N 8	O 5	S 1	0	0
2	B	1	Total 64	C 22	H 28	N 8	O 5	S 1	0	0

- Molecule 3 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: $C_4H_{12}NO_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	H	N	O	0	0
			20	4	12	1	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	H	N	O	0	0
			20	4	12	1	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	147	Total	O	0	0
			147	147		
4	B	167	Total	O	0	0
			167	167		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phenylethanolamine N-methyltransferase



- Molecule 1: Phenylethanolamine N-methyltransferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	94.66Å 94.66Å 188.46Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.62 – 1.95 38.62 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.2 (38.62-1.95) 90.6 (38.62-1.95)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.90 (at 1.95Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1377)	Depositor
R, R_{free}	0.182 , 0.230 0.190 , 0.239	Depositor DCC
R_{free} test set	3199 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	37.1	Xtriage
Anisotropy	0.454	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 31.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8617	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TRS, JIL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.73	1/2082 (0.0%)	0.80	0/2833
1	B	0.85	1/2150 (0.0%)	0.85	2/2926 (0.1%)
All	All	0.79	2/4232 (0.0%)	0.83	2/5759 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	118	PRO	N-CD	5.27	1.55	1.47
1	A	42	PRO	CA-C	5.05	1.54	1.51

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	117	GLU	CA-C-N	-7.31	112.47	119.85
1	B	117	GLU	C-N-CA	-7.31	112.47	119.85

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2016	1992	1972	11	0
1	B	2074	2053	2023	23	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	36	28	28	7	0
2	B	36	28	28	11	0
3	A	8	12	12	6	0
3	B	8	12	11	9	0
4	A	147	0	0	1	0
4	B	167	0	0	2	0
All	All	4492	4125	4074	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:301:JIL:S08	3:A:302:TRS:N	2.28	1.06
3:B:302:TRS:O1	3:B:302:TRS:O3	1.88	0.84
1:A:222:TYR:OH	2:A:301:JIL:H091	1.89	0.73
1:A:139:CYS:SG	1:B:48:CYS:HA	2.32	0.70
1:A:181:ALA:O	3:A:302:TRS:H31	1.96	0.66
1:B:222:TYR:CE1	2:B:301:JIL:H122	2.33	0.62
1:B:35:TYR:OH	2:B:301:JIL:H091	1.99	0.61
1:B:182:PHE:HB3	2:B:301:JIL:H111	1.68	0.58
1:B:35:TYR:OH	2:B:301:JIL:H231	2.06	0.56
1:B:83:THR:CG2	3:B:302:TRS:H32	2.37	0.55
1:B:142:ASP:OD1	1:B:145:ARG:NH2	2.36	0.54
1:B:20:GLN:NE2	4:B:514:HOH:O	2.41	0.53
1:B:80:SER:O	3:B:302:TRS:H21	2.08	0.53
2:A:301:JIL:C07	3:A:302:TRS:N	2.72	0.53
1:B:182:PHE:O	2:B:301:JIL:H092	2.10	0.52
1:A:182:PHE:O	2:A:301:JIL:H102	2.09	0.52
2:B:301:JIL:S08	3:B:302:TRS:N	2.84	0.51
1:A:35:TYR:CE1	2:A:301:JIL:H092	2.45	0.51
1:B:83:THR:OG1	3:B:302:TRS:H32	2.10	0.51
1:B:181:ALA:O	3:B:302:TRS:H12	2.10	0.50
1:A:75:LEU:HD23	1:A:76:ILE:N	2.28	0.49
1:B:80:SER:O	3:B:302:TRS:C2	2.61	0.48
1:B:83:THR:HG21	3:B:302:TRS:H32	1.95	0.48
1:A:48:CYS:CB	1:B:139:CYS:HG	2.26	0.48
1:B:222:TYR:CZ	2:B:301:JIL:H102	2.49	0.48
1:A:181:ALA:O	3:A:302:TRS:O1	2.32	0.46
2:B:301:JIL:C10	2:B:301:JIL:H252	2.46	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:57:LYS:NZ	2:B:301:JIL:O20	2.36	0.45
1:B:75:LEU:HD11	1:B:179:VAL:HG23	2.00	0.43
1:B:128[A]:GLN:HG2	1:B:140:TRP:CD1	2.53	0.43
1:B:80:SER:C	3:B:302:TRS:H22	2.43	0.43
1:B:74:THR:HG22	1:B:96:ASP:HB3	2.01	0.42
1:A:75:LEU:HD23	1:A:75:LEU:C	2.44	0.42
1:B:222:TYR:CD1	2:B:301:JIL:H122	2.55	0.42
1:B:253:LEU:HD13	1:B:275:ALA:HB2	2.02	0.42
1:A:80:SER:C	3:A:302:TRS:H12	2.46	0.41
1:A:56:TRP:CE2	1:A:256:TYR:HB2	2.56	0.41
2:A:301:JIL:H071	3:A:302:TRS:N	2.36	0.41
2:B:301:JIL:H121	4:B:510:HOH:O	2.22	0.40
2:A:301:JIL:H121	4:A:504:HOH:O	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	256/289 (89%)	253 (99%)	3 (1%)	0	100	100
1	B	267/289 (92%)	264 (99%)	3 (1%)	0	100	100
All	All	523/578 (90%)	517 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	214/233 (92%)	213 (100%)	1 (0%)	81	82
1	B	220/233 (94%)	219 (100%)	1 (0%)	81	82
All	All	434/466 (93%)	432 (100%)	2 (0%)	81	82

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	251	ARG
1	B	251	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	129	HIS
1	B	108	GLN
1	B	146	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	TRS	A	302	-	7,7,7	1.30	1 (14%)	9,9,9	2.15	3 (33%)
2	JIL	B	301	-	40,40,40	3.03	12 (30%)	51,57,57	2.66	24 (47%)
3	TRS	B	302	-	7,7,7	2.02	3 (42%)	9,9,9	3.55	7 (77%)
2	JIL	A	301	-	40,40,40	3.04	13 (32%)	51,57,57	2.58	24 (47%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TRS	A	302	-	-	6/9/9/9	-
2	JIL	B	301	-	-	7/15/42/42	0/5/5/5
3	TRS	B	302	-	-	8/9/9/9	-
2	JIL	A	301	-	-	5/15/42/42	0/5/5/5

All (29) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	301	JIL	O20-N19	12.33	1.44	1.22
2	A	301	JIL	O20-N19	11.30	1.42	1.22
2	A	301	JIL	C25-C13	8.06	1.59	1.52
2	B	301	JIL	C25-C13	7.99	1.59	1.52
2	A	301	JIL	C12-N11	-5.36	1.38	1.47
2	B	301	JIL	C31-N32	5.03	1.47	1.34
2	A	301	JIL	C23-C24	4.84	1.47	1.39
2	A	301	JIL	C31-N32	4.72	1.46	1.34
2	A	301	JIL	C13-N14	-4.15	1.40	1.47
2	B	301	JIL	C13-N14	-4.06	1.40	1.47
2	A	301	JIL	C30-N29	4.04	1.39	1.31
2	A	301	JIL	C25-C24	3.96	1.58	1.51
2	B	301	JIL	C30-N29	3.96	1.39	1.31
2	B	301	JIL	C12-N11	-3.90	1.40	1.47
2	B	301	JIL	C25-C24	3.69	1.58	1.51
2	B	301	JIL	C23-C24	3.60	1.45	1.39
2	B	301	JIL	C15-N14	-3.52	1.42	1.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	302	TRS	C3-C	-3.24	1.44	1.53
2	A	301	JIL	C15-N14	-3.15	1.42	1.46
3	B	302	TRS	C2-C	-3.05	1.45	1.53
2	B	301	JIL	C24-C16	-2.51	1.35	1.40
2	A	301	JIL	C24-C16	-2.48	1.35	1.40
2	B	301	JIL	C30-N26	-2.42	1.33	1.37
3	A	302	TRS	C2-C	-2.21	1.47	1.53
3	B	302	TRS	O1-C1	-2.17	1.35	1.42
2	A	301	JIL	C17-C16	2.15	1.42	1.39
2	B	301	JIL	C27-N35	2.13	1.38	1.34
2	A	301	JIL	C09-S08	-2.11	1.73	1.81
2	A	301	JIL	C27-N35	2.03	1.38	1.34

All (58) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	JIL	C27-N26-C04	-5.69	113.33	126.63
2	B	301	JIL	C28-C27-N35	-5.57	119.05	126.72
2	B	301	JIL	N35-C34-N33	-5.49	120.28	128.58
2	A	301	JIL	C27-N26-C30	5.42	111.43	105.74
3	B	302	TRS	O3-C3-C	-5.40	95.81	110.88
2	A	301	JIL	C28-C27-N35	-5.15	119.62	126.72
3	B	302	TRS	C1-C-N	5.09	121.14	108.17
2	B	301	JIL	C27-N26-C04	-5.06	114.79	126.63
2	B	301	JIL	C27-N26-C30	5.04	111.03	105.74
2	A	301	JIL	N35-C34-N33	-4.88	121.19	128.58
2	B	301	JIL	N35-C27-N26	4.85	135.42	127.17
2	A	301	JIL	N35-C27-N26	4.80	135.32	127.17
2	A	301	JIL	N26-C30-N29	-4.49	107.57	113.94
2	B	301	JIL	N26-C30-N29	-4.47	107.60	113.94
2	B	301	JIL	C03-C02-C06	-4.35	94.20	102.61
3	A	302	TRS	O2-C2-C	-4.17	99.26	110.88
3	B	302	TRS	C2-C-C1	-4.00	99.99	110.66
2	A	301	JIL	C07-S08-C09	3.92	113.89	102.26
2	B	301	JIL	O05-C04-N26	3.87	115.53	108.09
2	B	301	JIL	C34-N35-C27	3.84	121.22	111.83
2	B	301	JIL	C16-C15-N14	-3.83	104.10	113.25
2	B	301	JIL	O01-C02-C06	3.75	121.86	111.08
2	A	301	JIL	C04-N26-C30	3.62	135.12	127.09
2	A	301	JIL	C03-C04-N26	-3.61	104.32	113.30
3	A	302	TRS	C3-C-C2	-3.54	101.23	110.66
3	B	302	TRS	O2-C2-C	-3.53	101.04	110.88

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	JIL	C34-N35-C27	3.49	120.35	111.83
2	A	301	JIL	O05-C04-N26	3.35	114.53	108.09
2	B	301	JIL	C07-C06-C02	3.27	123.24	115.06
3	B	302	TRS	C3-C-C1	-3.27	101.95	110.66
2	A	301	JIL	C34-N33-C31	3.22	124.02	118.73
2	B	301	JIL	C28-N29-C30	3.22	108.51	103.45
2	B	301	JIL	C04-N26-C30	3.18	134.15	127.09
2	B	301	JIL	C22-C23-C24	-3.17	117.25	121.39
2	A	301	JIL	C15-N14-C13	3.13	116.85	111.28
2	A	301	JIL	C22-C18-C17	3.11	124.11	120.11
2	A	301	JIL	C22-C23-C24	-3.09	117.36	121.39
2	B	301	JIL	C03-C04-N26	-2.98	105.89	113.30
2	A	301	JIL	C22-C18-N19	-2.98	116.73	119.34
2	B	301	JIL	C34-N33-C31	2.91	123.52	118.73
2	B	301	JIL	C22-C18-C17	2.91	123.86	120.11
3	B	302	TRS	C3-C-C2	2.87	118.31	110.66
2	B	301	JIL	C27-C28-N29	-2.83	107.34	110.58
2	B	301	JIL	C23-C24-C16	2.82	122.44	118.98
2	A	301	JIL	O01-C02-C06	2.73	118.92	111.08
2	A	301	JIL	C23-C24-C16	2.70	122.29	118.98
3	B	302	TRS	O1-C1-C	2.67	118.32	110.88
2	B	301	JIL	C18-C17-C16	-2.62	116.74	119.81
2	A	301	JIL	C28-N29-C30	2.61	107.55	103.45
2	A	301	JIL	C16-C15-N14	-2.58	107.10	113.25
2	B	301	JIL	C15-N14-C13	2.49	115.69	111.28
2	A	301	JIL	C03-C02-C06	-2.45	97.88	102.61
2	A	301	JIL	C18-C17-C16	-2.36	117.05	119.81
2	A	301	JIL	O20-N19-C18	-2.27	115.69	118.82
3	A	302	TRS	C3-C-N	2.21	113.81	108.17
2	B	301	JIL	C22-C18-N19	-2.17	117.44	119.34
2	B	301	JIL	C06-O05-C04	-2.14	104.74	109.47
2	A	301	JIL	C02-C03-C04	2.13	105.48	101.46

There are no chirality outliers.

All (26) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	301	JIL	N11-C12-C13-C25
3	A	302	TRS	N-C-C1-O1
3	A	302	TRS	C1-C-C2-O2
3	A	302	TRS	C3-C-C2-O2
3	A	302	TRS	N-C-C2-O2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	B	302	TRS	N-C-C1-O1
2	A	301	JIL	C09-C10-N11-C12
2	B	301	JIL	C09-C10-N11-C12
3	B	302	TRS	C3-C-C1-O1
3	A	302	TRS	C3-C-C1-O1
3	B	302	TRS	N-C-C2-O2
3	B	302	TRS	N-C-C3-O3
2	A	301	JIL	C13-C12-N11-C10
2	A	301	JIL	C03-C04-N26-C30
2	B	301	JIL	C03-C04-N26-C30
2	B	301	JIL	N11-C12-C13-N14
3	A	302	TRS	C2-C-C1-O1
3	B	302	TRS	C2-C-C1-O1
2	A	301	JIL	C03-C04-N26-C27
2	B	301	JIL	C06-C07-S08-C09
3	B	302	TRS	C1-C-C2-O2
3	B	302	TRS	C3-C-C2-O2
3	B	302	TRS	C2-C-C3-O3
2	A	301	JIL	O05-C04-N26-C30
2	B	301	JIL	O05-C04-N26-C30
2	B	301	JIL	C03-C04-N26-C27


There are no ring outliers.

4 monomers are involved in 29 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	302	TRS	6	0
2	B	301	JIL	11	0
3	B	302	TRS	9	0
2	A	301	JIL	7	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	255/289 (88%)	0.30	8 (3%)	51	58	27, 50, 76, 101	2 (0%)
1	B	265/289 (91%)	0.05	7 (2%)	57	64	24, 45, 65, 103	4 (1%)
All	All	520/578 (89%)	0.17	15 (2%)	53	60	24, 47, 74, 103	6 (1%)

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	139	CYS	5.3
1	B	280	VAL	4.9
1	A	280	VAL	4.8
1	A	26	ALA	4.6
1	B	170	PRO	4.1
1	A	237[A]	GLU	3.4
1	B	16	SER	2.8
1	A	162	PRO	2.5
1	B	118	PRO	2.5
1	A	167	ALA	2.5
1	B	169	SER	2.3
1	B	227	ALA	2.2
1	A	170	PRO	2.2
1	A	168	GLY	2.0
1	B	29	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	TRS	B	302	8/8	0.83	0.14	31,45,79,79	0
3	TRS	A	302	8/8	0.92	0.13	37,58,77,77	0
2	JIL	B	301	36/36	0.94	0.14	33,53,111,226	0
2	JIL	A	301	36/36	0.95	0.16	36,62,261,335	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around JIL A 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.