



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 10:23 AM UTC

PDB ID : 6NAO / pdb_00006nao
Title : Discovery of a high affinity inhibitor of cGAS
Authors : Hall, J.
Deposited on : 2018-12-06
Resolution : 3.23 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.23 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2153 (3.28-3.20)
Clashscore	190562	2275 (3.28-3.20)
Ramachandran outliers	187476	2233 (3.28-3.20)
Sidechain outliers	187428	2232 (3.28-3.20)
RSRZ outliers	180081	2153 (3.28-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	362	7% 87% 10% ...
1	B	362	8% 89% 9% ..

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 10861 atoms, of which 5250 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called CYCLIC GMP-AMP SYNTHASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	358	Total	C	H	N	O	S	0	0	0
			5447	1785	2659	479	510	14			
1	B	361	Total	C	H	N	O	S	0	0	0
			5296	1757	2553	465	508	13			

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		

- Molecule 3 is (1R,2S)-2-[(7-hydroxy-5-phenylpyrazolo[1,5-a]pyrimidine-3-carbonyl)amino]cyclohexane-1-carboxylic acid (CCD ID: KHM) (formula: C₂₀H₂₀N₄O₄) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	H	N	O	0	0
			47	20	19	4	4		
3	B	1	Total	C	H	N	O	0	0
			47	20	19	4	4		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	13	Total	O	0	0
			13	13		
4	B	9	Total	O	0	0
			9	9		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: CYCLIC GMP-AMP SYNTHASE



• Molecule 1: CYCLIC GMP-AMP SYNTHASE



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	219.99Å 45.95Å 90.11Å 90.00° 111.46° 90.00°	Depositor
Resolution (Å)	102.37 – 3.23 102.37 – 3.23	Depositor EDS
% Data completeness (in resolution range)	99.2 (102.37-3.23) 99.2 (102.37-3.23)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.32 (at 3.19Å)	Xtriage
Refinement program	BUSTER 2.11.7	Depositor
R, R_{free}	0.217 , 0.253 0.231 , 0.261	Depositor DCC
R_{free} test set	626 reflections (4.49%)	wwPDB-VP
Wilson B-factor (Å ²)	63.0	Xtriage
Anisotropy	0.276	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 73.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.023 for -h-2*1,-k,l	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	10861	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, KHM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.87	2/2845 (0.1%)	1.24	13/3846 (0.3%)
1	B	0.86	1/2801 (0.0%)	1.23	7/3798 (0.2%)
All	All	0.86	3/5646 (0.1%)	1.24	20/7644 (0.3%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	276	MET	SD-CE	-6.05	1.64	1.79
1	B	276	MET	SD-CE	-5.76	1.65	1.79
1	A	234	VAL	CA-C	5.38	1.56	1.52

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	306	PRO	CA-C-N	8.29	131.74	120.54
1	B	306	PRO	C-N-CA	8.29	131.74	120.54
1	A	521	GLU	N-CA-C	7.39	126.55	110.80
1	B	210	ASN	CA-CB-CG	7.28	119.88	112.60
1	A	234	VAL	CB-CA-C	6.95	116.87	110.13
1	A	210	ASN	CA-CB-CG	6.87	119.47	112.60
1	A	256	ASN	CA-CB-CG	6.53	119.13	112.60
1	A	520	ASP	CA-C-N	6.53	134.00	121.54
1	A	520	ASP	C-N-CA	6.53	134.00	121.54
1	B	520	ASP	CA-CB-CG	6.17	118.77	112.60
1	A	221	SER	N-CA-C	6.02	122.88	113.72
1	A	213	SER	CA-C-N	5.87	128.07	120.44
1	A	213	SER	C-N-CA	5.87	128.07	120.44
1	B	221	SER	N-CA-C	5.71	122.39	113.72
1	B	290	ASP	CA-CB-CG	5.43	118.03	112.60
1	A	290	ASP	CA-CB-CG	5.27	117.87	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	221	SER	CA-C-N	-5.14	117.29	122.74
1	A	221	SER	C-N-CA	-5.14	117.29	122.74
1	A	214	TYR	N-CA-CB	-5.12	102.58	110.01
1	B	336	GLU	CB-CG-CD	5.07	121.21	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2788	2659	2658	10	0
1	B	2743	2553	2555	8	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	28	19	0	0	0
3	B	28	19	0	0	0
4	A	13	0	0	0	0
4	B	9	0	0	2	0
All	All	5611	5250	5213	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:454:GLN:HB2	4:B:701:HOH:O	1.82	0.78
1:B:449:ASN:C	4:B:701:HOH:O	2.47	0.58
1:A:234:VAL:HG23	1:A:237:ILE:HG21	1.85	0.57
1:A:216:GLU:HG2	1:A:384:LYS:HA	1.87	0.55
1:A:183:ALA:HB1	1:A:214:TYR:OH	2.11	0.51
1:A:262:LEU:HD22	1:A:276:MET:HE2	1.92	0.51
1:B:262:LEU:HD22	1:B:276:MET:HE2	1.92	0.51
1:B:419:GLN:HB3	1:B:518:VAL:HG22	1.92	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:336:GLU:HG2	1:B:486:PRO:HG2	1.93	0.49
1:A:234:VAL:CG2	1:A:237:ILE:HG21	2.42	0.48
1:B:216:GLU:HG2	1:B:384:LYS:HA	2.00	0.43
1:A:338:LEU:HD13	1:A:484:PHE:CE2	2.55	0.42
1:B:338:LEU:HD13	1:B:484:PHE:CE2	2.55	0.42
1:A:234:VAL:HG11	1:A:324:LEU:HD22	2.02	0.41
1:A:182:ALA:HB1	1:A:316:ILE:HD12	2.02	0.41
1:A:262:LEU:CD2	1:A:276:MET:HE2	2.50	0.41
1:B:209:LEU:O	1:B:210:ASN:CG	2.64	0.41
1:A:234:VAL:HG23	1:A:237:ILE:CG2	2.50	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	354/362 (98%)	332 (94%)	18 (5%)	4 (1%)	11	40
1	B	359/362 (99%)	333 (93%)	22 (6%)	4 (1%)	11	40
All	All	713/724 (98%)	665 (93%)	40 (6%)	8 (1%)	11	40

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	300	ARG
1	B	300	ARG
1	A	235	PRO
1	A	345	SER
1	B	345	SER
1	B	235	PRO
1	A	256	ASN
1	B	256	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	288/334 (86%)	262 (91%)	26 (9%)	9	32
1	B	275/334 (82%)	254 (92%)	21 (8%)	12	39
All	All	563/668 (84%)	516 (92%)	47 (8%)	10	36

All (47) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	170	GLU
1	A	181	THR
1	A	208	LEU
1	A	209	LEU
1	A	220	ILE
1	A	229	MET
1	A	231	LYS
1	A	233	GLU
1	A	246	ARG
1	A	255	ARG
1	A	256	ASN
1	A	259	GLU
1	A	262	LEU
1	A	266	LEU
1	A	276	MET
1	A	288	ILE
1	A	300	ARG
1	A	316	ILE
1	A	322	LEU
1	A	359	LEU
1	A	371	GLN
1	A	384	LYS
1	A	386	ILE
1	A	438	VAL
1	A	462	LEU
1	A	499	ARG
1	B	170	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	208	LEU
1	B	209	LEU
1	B	220	ILE
1	B	229	MET
1	B	231	LYS
1	B	233	GLU
1	B	246	ARG
1	B	258	LYS
1	B	259	GLU
1	B	262	LEU
1	B	276	MET
1	B	288	ILE
1	B	291	ILE
1	B	322	LEU
1	B	336	GLU
1	B	359	LEU
1	B	384	LYS
1	B	386	ILE
1	B	438	VAL
1	B	462	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	192	HIS
1	A	217	HIS
1	A	451	GLN
1	B	217	HIS
1	B	371	GLN
1	B	451	GLN
1	B	466	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	KHM	A	602	-	30,31,31	2.11	7 (23%)	39,44,44	2.17	14 (35%)
3	KHM	B	602	-	30,31,31	1.99	7 (23%)	39,44,44	2.14	14 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	KHM	A	602	-	-	2/16/27/27	0/4/4/4
3	KHM	B	602	-	-	4/16/27/27	0/4/4/4

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	602	KHM	C14-N15	5.83	1.46	1.34
3	B	602	KHM	C14-N15	5.69	1.46	1.34
3	A	602	KHM	C05-C04	4.53	1.54	1.47
3	A	602	KHM	C26-N27	4.43	1.39	1.32
3	B	602	KHM	C26-N27	4.42	1.39	1.32
3	B	602	KHM	C05-C04	3.88	1.53	1.47
3	B	602	KHM	C26-C13	-3.18	1.37	1.41
3	B	602	KHM	C12-N28	-3.01	1.34	1.38
3	A	602	KHM	C12-N28	-2.89	1.34	1.38
3	A	602	KHM	O01-C02	2.84	1.41	1.32

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	602	KHM	O01-C02	2.73	1.40	1.32
3	A	602	KHM	C26-C13	-2.72	1.37	1.41
3	A	602	KHM	O25-C14	-2.41	1.19	1.23
3	B	602	KHM	O25-C14	-2.23	1.19	1.23

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	602	KHM	C13-C12-N28	5.97	108.71	105.58
3	B	602	KHM	C13-C12-N28	5.77	108.61	105.58
3	B	602	KHM	C26-N27-N28	5.47	107.28	103.43
3	A	602	KHM	C26-N27-N28	5.44	107.25	103.43
3	A	602	KHM	C03-C04-N11	-3.91	118.11	121.83
3	B	602	KHM	C13-C26-N27	-3.37	108.27	112.45
3	A	602	KHM	C04-N11-C12	3.34	121.20	117.78
3	A	602	KHM	C18-C17-C16	3.34	116.59	111.42
3	B	602	KHM	C04-N11-C12	3.23	121.08	117.78
3	A	602	KHM	C13-C26-N27	-3.21	108.47	112.45
3	B	602	KHM	C13-C12-N11	3.05	131.84	123.54
3	B	602	KHM	C12-N28-N27	-2.97	109.11	112.24
3	B	602	KHM	C03-C04-N11	-2.90	119.07	121.83
3	A	602	KHM	C13-C12-N11	2.84	131.28	123.54
3	B	602	KHM	C26-C13-C12	2.76	106.63	104.57
3	A	602	KHM	C12-N28-N27	-2.74	109.35	112.24
3	B	602	KHM	C19-C18-C17	2.67	116.91	111.42
3	B	602	KHM	N28-C12-N11	-2.45	119.43	122.15
3	B	602	KHM	C13-C14-N15	2.45	119.44	116.25
3	B	602	KHM	C17-C16-C21	-2.45	108.04	111.14
3	A	602	KHM	C13-C14-N15	2.35	119.31	116.25
3	B	602	KHM	O23-C22-C21	2.34	120.01	113.93
3	A	602	KHM	O23-C22-C21	2.32	119.97	113.93
3	A	602	KHM	C19-C18-C17	2.30	116.14	111.42
3	B	602	KHM	O01-C02-C03	-2.29	120.77	125.58
3	A	602	KHM	O01-C02-C03	-2.29	120.78	125.58
3	A	602	KHM	C26-C13-C12	2.17	106.19	104.57
3	A	602	KHM	C18-C19-C20	2.05	115.64	111.42

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	602	KHM	C20-C21-C22-O23

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	A	602	KHM	C16-C21-C22-O24
3	B	602	KHM	C16-C21-C22-O23
3	B	602	KHM	C16-C21-C22-O24
3	B	602	KHM	C20-C21-C22-O24
3	A	602	KHM	C16-C21-C22-O23

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	358/362 (98%)	0.42	24 (6%) 24 16	21, 62, 109, 129	0
1	B	361/362 (99%)	0.49	28 (7%) 19 13	30, 66, 112, 139	0
All	All	719/724 (99%)	0.45	52 (7%) 21 14	21, 64, 111, 139	0

All (52) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	305	SER	4.8
1	A	213	SER	4.3
1	A	178	ASP	4.3
1	B	293	ASP	3.7
1	A	185	MET	3.6
1	B	185	MET	3.5
1	A	294	THR	3.4
1	B	298	MET	3.3
1	A	214	TYR	3.2
1	B	368	ASN	3.1
1	B	366	GLU	3.0
1	A	219	LYS	3.0
1	A	366	GLU	2.9
1	A	181	THR	2.8
1	A	218	VAL	2.8
1	B	284	ILE	2.7
1	A	293	ASP	2.7
1	B	172	LEU	2.7
1	A	341	GLN	2.6
1	A	284	ILE	2.6
1	B	370	PHE	2.6
1	A	368	ASN	2.6
1	B	178	ASP	2.5
1	A	522	PHE	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	223	PRO	2.5
1	B	302	ARG	2.5
1	A	220	ILE	2.5
1	B	306	PRO	2.4
1	B	301	LYS	2.3
1	B	367	GLY	2.3
1	A	177	ASP	2.3
1	B	295	ASP	2.3
1	B	256	ASN	2.3
1	A	373	GLU	2.3
1	B	269	GLU	2.3
1	A	179	ILE	2.3
1	B	179	ILE	2.3
1	B	192	HIS	2.3
1	B	483	TYR	2.3
1	B	181	THR	2.2
1	B	373	GLU	2.2
1	A	212	GLY	2.2
1	A	269	GLU	2.2
1	A	367	GLY	2.2
1	B	266	LEU	2.1
1	B	286	GLU	2.1
1	B	182	ALA	2.1
1	A	254	LYS	2.1
1	B	214	TYR	2.1
1	B	294	THR	2.1
1	A	190	VAL	2.0
1	B	288	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,



median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	KHM	B	602	28/28	0.90	0.14	45,57,63,69	0
3	KHM	A	602	28/28	0.91	0.11	45,56,60,61	0
2	ZN	A	601	1/1	0.99	0.07	57,57,57,57	0
2	ZN	B	601	1/1	1.00	0.08	48,48,48,48	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around KHM B 602:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around KHM A 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.