



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 12, 2026 – 03:16 PM UTC

PDB ID : 4NEN / pdb_00004nen
Title : An internal ligand-bound, metastable state of a leukocyte integrin, aXb2
Authors : Sen, M.; Yuki, K.; Springer, T.A.
Deposited on : 2013-10-29
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1094	% 81% 16% .
2	B	686	2% 81% 15% .
3	C	2	50% 50%
3	D	2	50% 50%
3	F	2	50% 50%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	G	2	 100%
3	H	2	 100%
4	E	6	 33% 50% 17%

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 13731 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Integrin alpha-X.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1063	Total	C	N	O	S	0	4	0
			8265	5239	1422	1567	37			

There are 17 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	42	ASP	ASN	engineered mutation	UNP P20702
A	368	ASP	SER	engineered mutation	UNP P20702
A	678	THR	ASN	engineered mutation	UNP P20702
A	885	SER	ASN	engineered mutation	UNP P20702
A	920	CYS	ASN	engineered mutation	UNP P20702
A	1083	PRO	-	expression tag	UNP P20702
A	1084	GLY	-	expression tag	UNP P20702
A	1085	PRO	-	expression tag	UNP P20702
A	1086	ALA	-	expression tag	UNP P20702
A	1087	ALA	-	expression tag	UNP P20702
A	1088	LEU	-	expression tag	UNP P20702
A	1089	GLN	-	expression tag	UNP P20702
A	1090	THR	-	expression tag	UNP P20702
A	1091	LEU	-	expression tag	UNP P20702
A	1092	PHE	-	expression tag	UNP P20702
A	1093	GLN	-	expression tag	UNP P20702
A	1094	GLY	-	expression tag	UNP P20702

- Molecule 2 is a protein called Integrin beta-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	665	Total	C	N	O	S	0	4	1
			5118	3154	912	987	65			

There are 15 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	190	ASP	ASN	engineered mutation	UNP P05107
B	232	LYS	ASN	engineered mutation	UNP P05107
B	674	CYS	VAL	engineered mutation	UNP P05107
B	675	GLY	-	expression tag	UNP P05107
B	676	GLY	-	expression tag	UNP P05107
B	677	PRO	-	expression tag	UNP P05107
B	678	ALA	-	expression tag	UNP P05107
B	679	ALA	-	expression tag	UNP P05107
B	680	LEU	-	expression tag	UNP P05107
B	681	GLN	-	expression tag	UNP P05107
B	682	THR	-	expression tag	UNP P05107
B	683	LEU	-	expression tag	UNP P05107
B	684	PHE	-	expression tag	UNP P05107
B	685	GLN	-	expression tag	UNP P05107
B	686	GLY	-	expression tag	UNP P05107

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	D	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	F	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	G	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	H	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	E	6	Total	C	N	O	0	0	0
			72	40	2	30			

- Molecule 5 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	3	Total	Ca	0	0
			3	3		
5	B	2	Total	Ca	0	0
			2	2		

- Molecule 6 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Mg	0	0
			1	1		
6	B	1	Total	Mg	0	0
			1	1		

- Molecule 7 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Cl	0	0
			1	1		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	74	Total	O	0	0
			74	74		
8	B	54	Total	O	0	0
			54	54		



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C: 50% 50%



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D: 50% 50%



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F: 50% 50%



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G: 100%



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H: 100%



- Molecule 4: alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	127.09Å 119.85Å 182.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.81 – 2.90 47.81 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.2 (47.81-2.90) 99.2 (47.81-2.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.34 (at 2.91Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, R_{free}	0.224 , 0.264 0.227 , 0.267	Depositor DCC
R_{free} test set	3117 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	69.2	Xtriage
Anisotropy	0.264	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 64.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	13731	wwPDB-VP
Average B, all atoms (Å ²)	89.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, MAN, CL, MG, NAG, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.28	0/8465	0.76	4/11502 (0.0%)
2	B	0.27	0/5223	0.78	3/7049 (0.0%)
All	All	0.28	0/13688	0.77	7/18551 (0.0%)

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	542	GLY	CA-C-N	8.70	130.72	119.84
1	A	542	GLY	C-N-CA	8.70	130.72	119.84
1	A	888	PRO	N-CA-CB	7.77	110.72	103.41
1	A	176	ASN	N-CA-C	-5.50	105.93	113.30
2	B	167	ASN	CA-C-N	5.24	124.86	119.56

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8265	0	8095	101	0
2	B	5118	0	4920	60	0
3	C	28	0	25	1	0
3	D	28	0	25	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	F	28	0	25	0	0
3	G	28	0	25	0	0
3	H	28	0	25	0	0
4	E	72	0	61	1	0
5	A	3	0	0	0	0
5	B	2	0	0	0	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
7	A	1	0	0	0	0
8	A	74	0	0	3	0
8	B	54	0	0	1	0
All	All	13731	0	13201	153	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 153 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:952:PHE:HB2	1:A:1011:PHE:HB2	1.68	0.75
1:A:685:ARG:HH12	1:A:694:GLU:HB2	1.53	0.72
2:B:361:ASP:HB2	2:B:390:GLN:HB3	1.75	0.68
1:A:995:ILE:HG21	1:A:1040:ILE:HG23	1.78	0.66
1:A:12:ARG:HG2	1:A:590:GLN:HG2	1.76	0.66

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	1061/1094 (97%)	1010 (95%)	48 (4%)	3 (0%)	36 65

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
2	B	663/686 (97%)	621 (94%)	41 (6%)	1 (0%)	43 72
All	All	1724/1780 (97%)	1631 (95%)	89 (5%)	4 (0%)	43 72

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	543	PRO
1	A	175	SER
1	A	528	GLU
2	B	555	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	909/932 (98%)	896 (99%)	13 (1%)	59 85
2	B	576/590 (98%)	567 (98%)	9 (2%)	55 83
All	All	1485/1522 (98%)	1463 (98%)	22 (2%)	59 84

5 of 22 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	174	LYS
2	B	330	VAL
2	B	303	ARG
2	B	399	ILE
1	A	438	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1039	GLN
1	A	1066	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	498	HIS
2	B	167	ASN
2	B	275	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

16 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	C	1	1,3	14,14,15	0.43	0	17,19,21	0.46	0
3	NAG	C	2	3	14,14,15	0.27	0	17,19,21	0.55	0
3	NAG	D	1	1,3	14,14,15	0.65	0	17,19,21	0.59	0
3	NAG	D	2	3	14,14,15	0.26	0	17,19,21	0.45	0
4	NAG	E	1	1,4	14,14,15	0.54	0	17,19,21	0.43	0
4	NAG	E	2	4	14,14,15	0.21	0	17,19,21	0.38	0
4	BMA	E	3	4	11,11,12	0.65	0	15,15,17	1.18	2 (13%)
4	MAN	E	4	4	11,11,12	0.92	0	15,15,17	1.17	3 (20%)
4	MAN	E	5	4	11,11,12	0.80	0	15,15,17	0.91	1 (6%)
4	MAN	E	6	4	11,11,12	1.02	1 (9%)	15,15,17	0.94	1 (6%)
3	NAG	F	1	1,3	14,14,15	0.58	1 (7%)	17,19,21	0.58	0
3	NAG	F	2	3	14,14,15	0.33	0	17,19,21	0.39	0
3	NAG	G	1	2,3	14,14,15	0.55	0	17,19,21	0.42	0
3	NAG	G	2	3	14,14,15	0.32	0	17,19,21	0.41	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	H	1	2,3	14,14,15	0.58	0	17,19,21	0.66	0
3	NAG	H	2	3	14,14,15	0.36	0	17,19,21	0.48	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	C	2	3	-	1/6/23/26	0/1/1/1
3	NAG	D	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	D	2	3	-	1/6/23/26	0/1/1/1
4	NAG	E	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	E	2	4	-	2/6/23/26	0/1/1/1
4	BMA	E	3	4	-	1/2/19/22	0/1/1/1
4	MAN	E	4	4	-	0/2/19/22	0/1/1/1
4	MAN	E	5	4	-	1/2/19/22	0/1/1/1
4	MAN	E	6	4	-	2/2/19/22	0/1/1/1
3	NAG	F	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	F	2	3	-	2/6/23/26	0/1/1/1
3	NAG	G	1	2,3	-	2/6/23/26	0/1/1/1
3	NAG	G	2	3	-	0/6/23/26	0/1/1/1
3	NAG	H	1	2,3	-	0/6/23/26	0/1/1/1
3	NAG	H	2	3	-	0/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	E	6	MAN	C2-C3	2.32	1.56	1.52
3	F	1	NAG	C1-C2	2.00	1.55	1.52

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	E	3	BMA	C1-O5-C5	2.82	115.96	112.19
4	E	3	BMA	O2-C2-C3	-2.46	105.05	110.15
4	E	6	MAN	O2-C2-C3	-2.45	105.07	110.15
4	E	4	MAN	C1-O5-C5	2.35	115.34	112.19
4	E	4	MAN	O3-C3-C2	2.31	114.76	110.05

There are no chirality outliers.

5 of 18 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	D	1	NAG	O5-C5-C6-O6
3	G	1	NAG	O5-C5-C6-O6
3	F	1	NAG	O5-C5-C6-O6
4	E	2	NAG	O5-C5-C6-O6
4	E	2	NAG	C4-C5-C6-O6


There are no ring outliers.


3 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	1	NAG	1	0
4	E	6	MAN	1	0
3	D	1	NAG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.














5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1063/1094 (97%)	-0.11	13 (1%) 76 69	34, 76, 143, 301	4 (0%)
2	B	665/686 (96%)	-0.02	15 (2%) 61 52	30, 94, 175, 250	5 (0%)
All	All	1728/1780 (97%)	-0.08	28 (1%) 70 62	30, 80, 165, 301	9 (0%)

The worst 5 of 28 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	428	ARG	4.6
2	B	8	VAL	4.2
2	B	37	SER	4.2
1	A	614	PRO	3.6
1	A	650	ARG	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	H	2	14/15	0.45	0.13	139,155,167,175	0
4	MAN	E	6	11/12	0.55	0.12	157,165,172,173	0
4	MAN	E	5	11/12	0.63	0.10	149,155,160,162	0
3	NAG	F	2	14/15	0.68	0.10	141,149,158,159	0
3	NAG	C	2	14/15	0.68	0.13	117,134,152,154	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	G	2	14/15	0.69	0.12	161,170,177,179	0
4	MAN	E	4	11/12	0.72	0.11	141,148,152,154	0
3	NAG	D	2	14/15	0.80	0.10	69,114,123,127	0
3	NAG	F	1	14/15	0.84	0.11	83,106,111,129	0
4	BMA	E	3	11/12	0.85	0.07	102,114,137,143	0
3	NAG	H	1	14/15	0.85	0.09	92,105,122,138	0
3	NAG	G	1	14/15	0.87	0.10	109,120,133,145	0
3	NAG	C	1	14/15	0.89	0.07	50,81,99,101	0
4	NAG	E	2	14/15	0.89	0.09	66,70,89,94	0
3	NAG	D	1	14/15	0.94	0.08	58,77,92,97	0
4	NAG	E	1	14/15	0.95	0.10	57,63,70,77	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain F:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain G:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain H:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	CL	A	1117	1/1	0.86	0.08	85,85,85,85	0
5	CA	A	1115	1/1	0.95	0.08	131,131,131,131	0
5	CA	B	706	1/1	0.97	0.04	92,92,92,92	0
5	CA	B	705	1/1	0.99	0.03	44,44,44,44	0
5	CA	A	1114	1/1	0.99	0.02	65,65,65,65	0
6	MG	B	707	1/1	0.99	0.04	59,59,59,59	0
5	CA	A	1113	1/1	0.99	0.03	54,54,54,54	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	MG	A	1116	1/1	1.00	0.02	72,72,72,72	0

6.5 Other polymers [i](#)

There are no such residues in this entry.