



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 5, 2026 – 03:47 PM EDT

PDB ID : 3NLI / pdb_00003nli
Title : Structure of endothelial nitric oxide synthase N368D mutant heme domain complexed with 6-{{(3'R,4'R)-3'-[2''-(3'''-fluorophenethylamino)ethoxy]pyrrolidin-4'-yl}methyl}-4-methylpyridin-2-amine
Authors : Delker, S.L.; Li, H.; Poulos, T.L.
Deposited on : 2010-06-21
Resolution : 1.98 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1506 (1.98-1.98)
Clashscore	190562	1534 (1.98-1.98)
Ramachandran outliers	187476	1518 (1.98-1.98)
Sidechain outliers	187428	1518 (1.98-1.98)
RSRZ outliers	180081	1506 (1.98-1.98)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	444	
1	B	444	

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 7256 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Nitric oxide synthase, endothelial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	404	Total	C	N	O	S	0	2	0
			3227	2051	568	592	16			
1	B	402	Total	C	N	O	S	0	0	0
			3202	2035	564	587	16			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	100	ARG	CYS	SEE REMARK 999	UNP P29473
A	368	ASP	ASN	engineered mutation	UNP P29473
B	100	ARG	CYS	SEE REMARK 999	UNP P29473
B	368	ASP	ASN	engineered mutation	UNP P29473

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (CCD ID: H4B) (formula: $C_9H_{15}N_5O_3$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			17	9	5	3		
3	B	1	Total	C	N	O	0	0
			17	9	5	3		

- Molecule 4 is ACETATE ION (CCD ID: ACT) (formula: $C_2H_3O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is 6-{{[(3R,4R)-4-(2-{{[2-(3-fluorophenyl)ethyl]amino}ethoxy)pyrrolidin-3-yl]methyl}}-4-methylpyridin-2-amine (CCD ID: JRR) (formula: $C_{21}H_{29}FN_4O$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	F	N	O	0	0
			27	21	1	4	1		
6	B	1	Total	C	F	N	O	0	0
			27	21	1	4	1		

- Molecule 7 is CACODYLIC ACID (CCD ID: CAD) (formula: $C_2H_7AsO_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total 3	As 1	C 2	0	0
7	B	1	Total 3	As 1	C 2	0	0

- Molecule 8 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	1	Total 1	Zn 1	0	0


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	317	Total 317	O 317	0	0
9	B	309	Total 309	O 309	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nitric oxide synthase, endothelial



- Molecule 1: Nitric oxide synthase, endothelial



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	57.96Å 106.80Å 156.63Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.70 – 1.98 42.70 – 1.98	Depositor EDS
% Data completeness (in resolution range)	99.4 (42.70-1.98) 99.5 (42.70-1.98)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.10 (at 1.98Å)	Xtriage
Refinement program	REFMAC 5.5.0089, CNS	Depositor
R, R_{free}	0.169 , 0.203 0.183 , 0.216	Depositor DCC
R_{free} test set	3409 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	28.1	Xtriage
Anisotropy	0.099	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 36.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7256	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, ZN, GOL, CAD, HEM, H4B, JRR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.58	0/3316	0.77	0/4516
1	B	0.57	0/3290	0.79	2/4480 (0.0%)
All	All	0.57	0/6606	0.78	2/8996 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	107	LEU	CA-C-N	-7.31	112.09	119.99
1	B	107	LEU	C-N-CA	-7.31	112.09	119.99

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3227	0	3132	23	0
1	B	3202	0	3107	25	0
2	A	43	0	30	1	0
2	B	43	0	30	2	0
3	A	17	0	15	1	0
3	B	17	0	15	1	0
4	A	4	0	3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	4	0	3	0	0
5	A	6	0	8	0	0
5	B	6	0	8	0	0
6	A	27	0	29	0	0
6	B	27	0	29	1	0
7	A	3	0	0	2	0
7	B	3	0	0	1	0
8	A	1	0	0	0	0
9	A	317	0	0	8	0
9	B	309	0	0	4	0
All	All	7256	0	6409	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:240:ARG:HD3	1:A:298:PRO:HB3	1.68	0.74
1:B:236:ARG:HD3	1:B:351:SER:HB3	1.70	0.74
1:A:378:LEU:HB2	9:A:1008:HOH:O	2.00	0.61
1:A:72:ARG:HD3	9:A:1179:HOH:O	2.01	0.61
1:A:340:ASN:H	1:A:340:ASN:HD22	1.49	0.61
1:A:285:ASN:HB3	9:A:1006:HOH:O	2.03	0.59
1:B:107:LEU:O	9:B:1216:HOH:O	2.17	0.58
1:A:384:CYS:CB	7:A:950:CAD:AS	3.13	0.57
1:B:236:ARG:HD2	1:B:242:ASP:OD1	2.05	0.56
1:B:323:GLU:HG3	9:B:1242:HOH:O	2.07	0.54
1:A:360:MET:HE3	1:A:362:THR:OG1	2.07	0.54
1:B:362:THR:HA	1:B:405:ASN:HD21	1.74	0.53
2:B:500:HEM:HBB2	2:B:500:HEM:HHC	1.91	0.52
1:A:252:ARG:CD	9:A:1247:HOH:O	2.57	0.52
1:A:370:CYS:SG	1:A:378:LEU:HD13	2.51	0.51
1:B:362:THR:HA	1:B:405:ASN:ND2	2.25	0.51
1:B:240:ARG:HD3	1:B:298:PRO:HB3	1.92	0.51
1:B:370:CYS:SG	1:B:378:LEU:HD13	2.51	0.51
1:B:324:TRP:HB2	7:B:950:CAD:C1	2.41	0.51
1:B:233:PHE:HB3	1:B:234:PRO:CD	2.42	0.49
1:B:122:ALA:N	9:B:1159:HOH:O	2.45	0.49
1:A:126:LEU:HD11	1:A:156:GLU:HA	1.94	0.49
1:A:252:ARG:HD2	9:A:1247:HOH:O	2.13	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:360:MET:HE3	1:B:362:THR:OG1	2.12	0.48
1:B:378:LEU:HB2	9:B:1004:HOH:O	2.12	0.48
1:B:455:SER:HB3	1:B:458:LEU:HD22	1.97	0.47
1:B:366:THR:O	1:B:370:CYS:HB2	2.16	0.46
1:A:249:GLN:HB2	1:A:252:ARG:HG2	1.98	0.46
1:A:252:ARG:HD3	9:A:1247:HOH:O	2.14	0.46
1:A:384:CYS:SG	7:A:950:CAD:C1	3.03	0.46
1:A:96:CYS:HB3	1:B:96:CYS:HB3	1.98	0.45
1:B:240:ARG:HD2	1:B:241:GLY:O	2.17	0.45
1:A:240:ARG:NH2	9:A:1018:HOH:O	2.50	0.45
1:A:449:TRP:HA	3:A:600:H4B:N1	2.32	0.45
1:A:427:VAL:HG23	9:A:1111:HOH:O	2.16	0.44
1:B:340:ASN:O	1:B:478:GLN:HG3	2.15	0.44
1:B:472:SER:HA	1:B:473:PRO:C	2.43	0.43
1:A:472:SER:HA	1:A:473:PRO:C	2.43	0.43
1:A:246:TRP:HB2	1:A:294:LEU:HB3	2.00	0.43
1:B:449:TRP:HA	3:B:600:H4B:N1	2.33	0.42
2:A:500:HEM:HBB2	2:A:500:HEM:HHC	2.02	0.42
1:A:141:LYS:HA	1:A:141:LYS:HD3	1.88	0.42
1:B:167:LEU:HG	1:B:348:LEU:HD12	2.02	0.42
1:B:236:ARG:HG3	1:B:349:GLU:HB2	2.02	0.42
2:B:500:HEM:HBA2	6:B:800:JRR:H4	2.02	0.41
1:B:319:HIS:CG	1:B:320:PRO:HD2	2.56	0.41
1:A:231:THR:O	1:A:353:ALA:HA	2.20	0.41
1:B:231:THR:O	1:B:353:ALA:HA	2.21	0.40
1:A:105:LEU:HD22	1:B:465:GLU:HB3	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	402/444 (90%)	398 (99%)	4 (1%)	0	100	100
1	B	398/444 (90%)	391 (98%)	7 (2%)	0	100	100
All	All	800/888 (90%)	789 (99%)	11 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	345/377 (92%)	332 (96%)	13 (4%)	29	19
1	B	342/377 (91%)	326 (95%)	16 (5%)	23	12
All	All	687/754 (91%)	658 (96%)	29 (4%)	26	15

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	69	LYS
1	A	102	LEU
1	A	105	LEU
1	A	123	GLU
1	A	125	LEU
1	A	260	ASP
1	A	293	LEU
1	A	294	LEU
1	A	328	LEU
1	A	340	ASN
1	A	378	LEU
1	A	386	ASP
1	A	458	LEU
1	B	99	ARG
1	B	102	LEU
1	B	105	LEU
1	B	123	GLU
1	B	141	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	153	GLN
1	B	223	ARG
1	B	236	ARG
1	B	240	ARG
1	B	260	ASP
1	B	323	GLU
1	B	330	LEU
1	B	340	ASN
1	B	389	THR
1	B	458	LEU
1	B	478	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	92	GLN
1	A	124	GLN
1	A	153	GLN
1	A	166	HIS
1	A	191	GLN
1	A	207	GLN
1	A	259	GLN
1	A	340	ASN
1	A	376	ASN
1	A	413	GLN
1	A	468	ASN
1	B	124	GLN
1	B	128	GLN
1	B	222	ASN
1	B	225	ASN
1	B	340	ASN
1	B	376	ASN
1	B	405	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 1 is monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	JRR	A	800	-	28,29,29	0.65	0	29,38,38	1.66	6 (20%)
4	ACT	A	860	-	3,3,3	0.81	0	3,3,3	0.73	0
2	HEM	A	500	1	50,50,50	1.94	11 (22%)	67,82,82	1.36	7 (10%)
7	CAD	B	950	1	0,2,4	-	-	0,1,6	-	-
5	GOL	B	880	-	5,5,5	0.39	0	5,5,5	0.56	0
3	H4B	A	600	-	17,18,18	0.98	1 (5%)	14,26,26	1.76	5 (35%)
3	H4B	B	600	-	17,18,18	0.95	1 (5%)	14,26,26	2.06	6 (42%)
5	GOL	A	880	-	5,5,5	0.43	0	5,5,5	0.39	0
7	CAD	A	950	1	0,2,4	-	-	0,1,6	-	-
6	JRR	B	800	-	28,29,29	0.64	0	29,38,38	1.54	3 (10%)
4	ACT	B	860	-	3,3,3	0.81	0	3,3,3	0.77	0
2	HEM	B	500	1	50,50,50	2.09	11 (22%)	67,82,82	1.46	9 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	JRR	A	800	-	-	1/13/23/23	0/3/3/3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	500	1	-	2/14/54/54	-
5	GOL	B	880	-	-	0/4/4/4	-
3	H4B	A	600	-	-	0/8/17/17	0/2/2/2
3	H4B	B	600	-	-	0/8/17/17	0/2/2/2
5	GOL	A	880	-	-	0/4/4/4	-
6	JRR	B	800	-	-	1/13/23/23	0/3/3/3
2	HEM	B	500	1	-	2/14/54/54	-

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	500	HEM	C3D-C2D	7.55	1.53	1.36
2	A	500	HEM	C3D-C2D	7.17	1.52	1.36
2	B	500	HEM	FE-NB	5.27	2.11	1.94
2	A	500	HEM	FE-NB	5.09	2.10	1.94
2	B	500	HEM	FE-ND	4.91	2.10	1.94
2	B	500	HEM	FE-NA	4.82	2.11	1.95
2	B	500	HEM	FE-NC	4.69	2.10	1.95
2	A	500	HEM	FE-NC	4.20	2.09	1.95
2	A	500	HEM	FE-ND	4.13	2.07	1.94
2	A	500	HEM	FE-NA	3.49	2.06	1.95
2	B	500	HEM	CAB-C3B	3.30	1.56	1.47
2	A	500	HEM	CAB-C3B	3.15	1.55	1.47
2	B	500	HEM	CAC-C3C	3.01	1.55	1.47
2	A	500	HEM	CAC-C3C	2.98	1.55	1.47
3	B	600	H4B	C7-C6	2.86	1.54	1.52
2	B	500	HEM	CMD-C2D	2.50	1.55	1.50
3	A	600	H4B	C7-C6	2.46	1.54	1.52
2	A	500	HEM	CMB-C2B	2.36	1.55	1.50
2	A	500	HEM	CMA-C3A	2.32	1.55	1.50
2	B	500	HEM	CMB-C2B	2.25	1.55	1.50
2	A	500	HEM	C2A-C3A	-2.21	1.33	1.38
2	A	500	HEM	CMC-C2C	2.13	1.55	1.50
2	B	500	HEM	CMA-C3A	2.06	1.55	1.50
2	B	500	HEM	CMC-C2C	2.01	1.54	1.50

All (36) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	800	JRR	C6A-N1A-C2A	5.29	122.02	118.07
6	A	800	JRR	C6A-N1A-C2A	5.24	121.99	118.07
2	A	500	HEM	C4D-ND-C1D	5.14	111.29	105.21

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	500	HEM	C4D-ND-C1D	4.63	110.69	105.21
2	B	500	HEM	CBA-CAA-C2A	-4.55	99.96	112.53
3	B	600	H4B	C2-N1-C8A	4.40	121.12	113.36
3	A	600	H4B	C2-N1-C8A	4.06	120.52	113.36
2	B	500	HEM	CBD-CAD-C3D	-3.68	102.36	112.53
2	A	500	HEM	CBA-CAA-C2A	-3.38	103.17	112.53
2	A	500	HEM	CBD-CAD-C3D	-3.14	103.84	112.53
6	B	800	JRR	C5'-N1'-C2'	3.13	113.24	105.45
3	B	600	H4B	C4A-C4-N3	2.89	120.07	112.13
6	A	800	JRR	C7A-C2A-N1A	2.85	120.06	116.57
6	A	800	JRR	C5'-N1'-C2'	2.82	112.47	105.45
6	A	800	JRR	C3A-C2A-N1A	-2.81	119.53	122.73
2	B	500	HEM	C3B-C2B-C1B	2.78	108.50	106.41
2	B	500	HEM	CMD-C2D-C1D	2.69	129.23	125.03
3	B	600	H4B	C4-C4A-N5	2.56	123.26	116.27
3	B	600	H4B	O4-C4-C4A	-2.55	121.12	127.26
6	A	800	JRR	C14-C13-C12	-2.54	119.88	123.23
2	A	500	HEM	CAD-C3D-C4D	2.50	129.06	124.70
3	B	600	H4B	C2-N3-C4	-2.48	120.61	125.11
2	B	500	HEM	C2A-C1A-NA	-2.44	107.44	110.15
2	B	500	HEM	C3B-C4B-NB	-2.43	107.72	109.47
3	A	600	H4B	C4A-C4-N3	2.41	118.74	112.13
2	A	500	HEM	C3B-C4B-NB	-2.38	107.76	109.47
6	B	800	JRR	C14-C13-C12	-2.33	120.16	123.23
2	B	500	HEM	C1B-NB-C4B	2.32	107.96	105.21
3	A	600	H4B	C4-C4A-N5	2.17	122.19	116.27
2	A	500	HEM	C2A-C1A-NA	-2.16	107.76	110.15
6	A	800	JRR	C5A-C4A-C3A	2.16	120.51	118.11
3	B	600	H4B	N2-C2-N3	2.13	121.25	116.76
3	A	600	H4B	O4-C4-C4A	-2.12	122.15	127.26
2	A	500	HEM	CMD-C2D-C1D	2.06	128.25	125.03
3	A	600	H4B	N2-C2-N3	2.04	121.07	116.76
2	B	500	HEM	CAD-C3D-C4D	2.01	128.21	124.70

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	B	800	JRR	C4-C3-N2-C2
6	A	800	JRR	C1-C2-N2-C3
2	B	500	HEM	CAA-CBA-CGA-O2A
2	B	500	HEM	CAA-CBA-CGA-O1A

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	500	HEM	CAA-CBA-CGA-O2A
2	A	500	HEM	CAA-CBA-CGA-O1A


There are no ring outliers.


7 monomers are involved in 8 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	500	HEM	1	0
7	B	950	CAD	1	0
3	A	600	H4B	1	0
3	B	600	H4B	1	0
7	A	950	CAD	2	0
6	B	800	JRR	1	0
2	B	500	HEM	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	404/444 (90%)	0.05	15 (3%)	45 55	11, 28, 49, 66	2 (0%)
1	B	402/444 (90%)	0.04	12 (2%)	52 62	19, 29, 48, 66	0
All	All	806/888 (90%)	0.05	27 (3%)	49 59	11, 28, 49, 66	2 (0%)

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	239	GLY	5.0
1	A	68	PRO	4.2
1	B	108	PRO	4.2
1	A	108	PRO	3.9
1	A	122	ALA	3.9
1	B	122	ALA	3.7
1	A	67	GLY	3.2
1	A	285	ASN	3.0
1	B	323	GLU	2.7
1	B	123	GLU	2.7
1	A	123	GLU	2.6
1	B	107	LEU	2.6
1	A	238	PRO	2.6
1	A	69	LYS	2.6
1	A	477	TYR	2.5
1	B	239	GLY	2.5
1	B	259	GLN	2.4
1	A	259	GLN	2.3
1	B	124	GLN	2.3
1	B	479	PRO	2.3
1	B	109	ARG	2.3
1	B	257	ARG	2.2
1	A	91	GLN	2.2
1	A	99	ARG	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	482	TRP	2.2
1	A	127	SER	2.1
1	B	147	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	GOL	B	880	6/6	0.77	0.16	46,46,47,48	0
5	GOL	A	880	6/6	0.86	0.13	43,44,44,44	0
4	ACT	B	860	4/4	0.92	0.12	36,39,39,40	0
6	JRR	B	800	27/27	0.95	0.07	22,26,29,29	0
3	H4B	A	600	17/17	0.96	0.05	18,20,22,23	0
6	JRR	A	800	27/27	0.96	0.06	22,26,30,30	0
4	ACT	A	860	4/4	0.96	0.07	33,36,37,37	0
3	H4B	B	600	17/17	0.97	0.05	17,20,24,24	0
7	CAD	A	950	3/5	0.97	0.11	50,50,50,51	0
2	HEM	A	500	43/43	0.98	0.05	17,20,25,27	0
7	CAD	B	950	3/5	0.98	0.10	50,50,51,51	0
2	HEM	B	500	43/43	0.99	0.05	17,20,25,26	0
8	ZN	A	900	1/1	1.00	0.01	22,22,22,22	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around JRR B 800:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around JRR A 800:**


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM A 500:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.