



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 10:56 AM UTC

PDB ID : 5NOK / pdb_00005nok
Title : Polysaccharide Lyase BACCELL_00875
Authors : Cartmell, A.; Munoz-Munoz, J.; Terrapon, N.; Basle, A.; Henrissat, B.;
Gilbert, H.J.
Deposited on : 2017-04-12
Resolution : 2.24 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.24 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3416 (2.26-2.22)
Clashscore	190562	3556 (2.26-2.22)
Ramachandran outliers	187476	3500 (2.26-2.22)
Sidechain outliers	187428	3501 (2.26-2.22)
RSRZ outliers	180081	3415 (2.26-2.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	694	5% 88% 8% .
1	B	694	% 89% 7% ..

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 10944 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BACCELL_00875.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	668	Total	C	N	O	S	Se	0	0	0
			5307	3385	906	991	10	15			
1	B	669	Total	C	N	O	S	Se	0	0	0
			5381	3432	919	1005	10	15			

- Molecule 2 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: $C_6H_{13}NO_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	112	Total	O	0	0
			112	112		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	132	Total 132	O 132	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: BACCELL_00875



• Molecule 1: BACCELL_00875



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	116.73Å 116.73Å 228.49Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.48 – 2.24 47.48 – 2.24	Depositor EDS
% Data completeness (in resolution range)	100.0 (47.48-2.24) 100.0 (47.48-2.24)	Depositor EDS
R_{merge}	0.01	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.88 (at 2.24Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.195 , 0.249 0.208 , 0.254	Depositor DCC
R_{free} test set	3842 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	36.8	Xtriage
Anisotropy	0.382	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 23.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	10944	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.78	0/5425	0.87	1/7342 (0.0%)
1	B	0.77	0/5504	0.88	0/7448
All	All	0.77	0/10929	0.87	1/14790 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	52	MSE	CG-SE-CE	-5.08	87.74	98.92

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5307	0	5025	28	0
1	B	5381	0	5124	32	0
2	B	12	0	13	0	0
3	A	112	0	0	0	0
3	B	132	0	0	1	0
All	All	10944	0	10162	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 3.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:588:ASN:HD22	1:A:640:ASN:HD21	1.16	0.90
1:B:588:ASN:HD22	1:B:640:ASN:HD21	1.27	0.78
1:A:446:ASN:HD21	1:A:484:GLN:HE22	1.39	0.71
1:B:588:ASN:ND2	1:B:640:ASN:HD21	1.88	0.71
1:A:588:ASN:ND2	1:A:640:ASN:HD21	1.87	0.69
1:B:657:CYS:HA	1:B:680:GLN:HE21	1.57	0.69
1:B:452:MSE:CE	1:B:478:LEU:HD23	2.25	0.67
1:A:79:VAL:HG23	1:A:137:VAL:CG2	2.25	0.67
1:B:678:ASN:H	1:B:680:GLN:HE22	1.42	0.67
1:B:403:VAL:HG11	1:B:460:GLN:HB3	1.80	0.63
1:B:478:LEU:HD13	1:B:515:LEU:HD11	1.81	0.63
1:B:446:ASN:HD21	1:B:484:GLN:HE22	1.48	0.62
1:B:252:LEU:HD23	1:B:259:LEU:HD21	1.82	0.61
1:A:403:VAL:HG11	1:A:460:GLN:HB3	1.81	0.61
1:A:355:MSE:O	1:A:363:ASP:O	2.19	0.60
1:B:446:ASN:HD21	1:B:484:GLN:NE2	1.98	0.60
1:A:79:VAL:HG23	1:A:137:VAL:HG21	1.85	0.59
1:B:614:TRP:CE3	1:B:617:ILE:HD11	2.39	0.57
1:A:446:ASN:HD21	1:A:484:GLN:NE2	2.01	0.57
1:B:110:ILE:HD11	1:B:137:VAL:HG11	1.87	0.57
1:B:104:LEU:HD11	1:B:111:ARG:HD2	1.87	0.56
1:B:356:ASN:HD22	1:B:420:ARG:HH22	1.54	0.56
1:A:541:GLU:OE2	1:A:543:SER:OG	2.25	0.55
1:B:452:MSE:HG3	1:B:494:ILE:HA	1.89	0.54
1:B:541:GLU:OE1	1:B:543:SER:OG	2.17	0.54
1:B:452:MSE:HE3	1:B:478:LEU:HD23	1.90	0.54
1:B:447:ARG:HD2	1:B:537:GLU:OE2	2.08	0.53
1:A:291:ASN:ND2	1:A:317:ARG:H	2.08	0.52
1:B:452:MSE:CE	1:B:478:LEU:CD2	2.88	0.51
1:A:385:ASN:ND2	1:A:388:ASP:OD2	2.43	0.51
1:A:50:ARG:HE	1:A:160:ASN:HD22	1.58	0.51
1:B:588:ASN:HD22	1:B:640:ASN:ND2	2.04	0.50
1:A:471:VAL:CG1	1:A:503:LEU:HD12	2.43	0.49
1:A:122:ASP:OD2	1:A:255:LYS:NZ	2.43	0.49
1:A:205:ARG:NH2	1:A:594:HIS:O	2.43	0.49
1:A:346:VAL:HG21	1:A:405:GLN:HG2	1.94	0.48
1:B:38:GLY:O	1:B:116:ARG:NH2	2.45	0.46
1:B:156:GLN:HE22	1:B:532:ASN:HD22	1.61	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:152:TYR:HA	1:A:158:CYS:SG	2.56	0.46
1:A:31:MSE:HE1	1:A:118:ILE:HD11	1.99	0.45
1:B:45:LEU:HG	1:B:241:ALA:HB2	1.98	0.45
1:B:452:MSE:SE	1:B:481:MSE:HE2	2.67	0.45
1:B:579:ASN:ND2	3:B:801:HOH:O	2.49	0.45
1:A:31:MSE:CE	1:A:118:ILE:HD11	2.46	0.44
1:A:679:ASP:OD1	1:A:679:ASP:N	2.47	0.44
1:B:497:ARG:H	1:B:554:GLN:HE21	1.66	0.44
1:A:471:VAL:HG13	1:A:503:LEU:HD12	1.99	0.43
1:B:252:LEU:HD12	1:B:257:SER:HB3	2.00	0.43
1:B:588:ASN:ND2	1:B:640:ASN:ND2	2.64	0.43
1:B:190:LEU:C	1:B:190:LEU:HD23	2.43	0.43
1:A:431:THR:HG22	1:A:442:GLN:HE22	1.83	0.43
1:A:662:PRO:O	1:A:670:ALA:HB3	2.19	0.43
1:A:508:MSE:HE2	1:A:511:GLU:HG3	2.00	0.43
1:A:72:TRP:CD1	1:A:72:TRP:C	2.97	0.42
1:A:549:ILE:HD11	1:A:571:GLN:HB3	2.01	0.42
1:B:678:ASN:N	1:B:680:GLN:HE22	2.13	0.42
1:B:542:GLN:HE21	1:B:593:ARG:NH1	2.18	0.41
1:A:156:GLN:HE22	1:A:532:ASN:HA	1.86	0.41
1:B:680:GLN:H	1:B:680:GLN:CD	2.29	0.41
1:A:199:ASN:CG	1:A:663:TYR:HB2	2.46	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	664/694 (96%)	633 (95%)	31 (5%)	0	100	100
1	B	667/694 (96%)	640 (96%)	27 (4%)	0	100	100
All	All	1331/1388 (96%)	1273 (96%)	58 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	548/586 (94%)	538 (98%)	10 (2%)	51	61
1	B	563/586 (96%)	546 (97%)	17 (3%)	36	43
All	All	1111/1172 (95%)	1084 (98%)	27 (2%)	43	51

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	51	SER
1	A	67	LYS
1	A	213	GLN
1	A	234	SER
1	A	287	THR
1	A	395	ARG
1	A	423	ASP
1	A	481	MSE
1	A	679	ASP
1	A	694	LEU
1	B	36	SER
1	B	45	LEU
1	B	137	VAL
1	B	144	VAL
1	B	188	LEU
1	B	243	ASN
1	B	269	LYS
1	B	275	VAL
1	B	306	VAL
1	B	396	LEU
1	B	481	MSE
1	B	508	MSE
1	B	513	THR
1	B	541	GLU
1	B	627	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	651	GLU
1	B	680	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (30) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	156	GLN
1	A	160	ASN
1	A	187	HIS
1	A	211	ASN
1	A	243	ASN
1	A	245	ASN
1	A	291	ASN
1	A	353	GLN
1	A	442	GLN
1	A	484	GLN
1	A	517	ASN
1	A	554	GLN
1	A	588	ASN
1	B	69	ASN
1	B	131	ASN
1	B	211	ASN
1	B	245	ASN
1	B	291	ASN
1	B	353	GLN
1	B	356	ASN
1	B	442	GLN
1	B	446	ASN
1	B	484	GLN
1	B	532	ASN
1	B	542	GLN
1	B	554	GLN
1	B	579	ASN
1	B	588	ASN
1	B	644	ASN
1	B	680	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	MES	B	701	-	12,12,12	2.00	1 (8%)	15,16,16	1.59	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MES	B	701	-	-	4/6/14/14	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	701	MES	C8-S	-6.24	1.68	1.77

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	701	MES	O2S-S-C8	4.47	113.48	106.73
2	B	701	MES	O3S-S-O2S	-2.02	106.35	111.40

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	701	MES	C7-C8-S-O1S
2	B	701	MES	C7-C8-S-O2S
2	B	701	MES	C7-C8-S-O3S
2	B	701	MES	N4-C7-C8-S

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	653/694 (94%)	0.28	33 (5%) 33 32	20, 38, 73, 93	0
1	B	654/694 (94%)	0.04	5 (0%) 82 84	23, 38, 60, 80	0
All	All	1307/1388 (94%)	0.16	38 (2%) 53 54	20, 38, 69, 93	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	502	SER	4.3
1	A	533	TYR	4.2
1	A	24	ALA	4.0
1	A	22	LEU	3.8
1	A	467	ASP	3.3
1	A	503	LEU	3.3
1	A	444	ASN	3.2
1	B	694	LEU	3.2
1	A	500	LEU	3.2
1	A	512	TYR	3.1
1	A	509	LYS	3.1
1	A	361	LEU	3.0
1	A	462	TYR	2.9
1	A	384	CYS	2.9
1	A	506	ALA	2.9
1	A	531	LEU	2.9
1	B	358	PRO	2.8
1	A	487	TYR	2.6
1	A	520	ILE	2.6
1	A	23	SER	2.5
1	A	411	ASP	2.5
1	A	485	PHE	2.4
1	A	489	PHE	2.3
1	A	433	ASN	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	381	THR	2.2
1	A	431	THR	2.2
1	A	440	VAL	2.2
1	A	358	PRO	2.2
1	A	560	GLY	2.2
1	B	118	ILE	2.1
1	A	693	GLY	2.1
1	A	359	SER	2.1
1	A	517	ASN	2.1
1	A	515	LEU	2.1
1	B	26	ASN	2.1
1	A	382	PRO	2.1
1	B	51	SER	2.0
1	A	432	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	MES	B	701	12/12	0.74	0.26	68,82,120,124	0

6.5 Other polymers [i](#)

There are no such residues in this entry.