



wwPDB EM Validation Summary Report ⓘ

Sep 5, 2026 – 01:04 PM EDT

PDB ID : 9NSP / pdb_00009nsp
EMDB ID : EMD-49747
Title : MPXV replisome bound to ssDNA
Authors : Yu, Z.; Abraham, J.
Deposited on : 2025-03-17
Resolution : 4.10 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev133
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 4.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	6458 (3.60 - 4.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	D	785	15% 74% 12% 14%
1	E	785	36% 57% 14% 28%
1	H	785	13% 40% 5% 55%
1	I	785	12% 41% 55%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	J	785	11% 36% 5% 59%
1	K	785	13% 39% 5% 55%
2	A	1006	84% 13%
3	C	426	8% 73% 18% 10%
4	B	218	11% 79% 20%
5	T	29	14% 52% 28% 21%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 34677 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uncoating factor OPG117.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	D	674	Total	C	N	O	S	0	0
			5469	3479	932	1027	31		
1	E	566	Total	C	N	O	S	0	0
			4599	2923	779	869	28		
1	I	351	Total	C	N	O	S	0	0
			2857	1831	482	529	15		
1	H	351	Total	C	N	O	S	0	0
			2855	1832	482	526	15		
1	J	324	Total	C	N	O	S	0	0
			2635	1684	452	485	14		
1	K	351	Total	C	N	O	S	0	0
			2856	1831	483	528	14		

- Molecule 2 is a protein called DNA polymerase.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	A	981	Total	C	N	O	S	0	0
			8031	5134	1340	1504	53		

- Molecule 3 is a protein called DNA polymerase processivity factor component OPG148.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	C	385	Total	C	N	O	S	0	0
			3141	2028	513	590	10		

- Molecule 4 is a protein called Uracil-DNA glycosylase.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	B	218	Total	C	N	O	S	0	0
			1773	1149	292	326	6		

- Molecule 5 is a DNA chain called DNA (5'-D(P*TP*TP*TP*TP*TP*TP*TP*TP*TP*TP*TP*TP*T)-3').

Mol	Chain	Residues	Atoms					AltConf	Trace
5	T	23	Total	C	N	O	P	0	0
			460	230	46	161	23		


- Molecule 6 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
6	D	1	Total	Zn	0
			1	1	


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Uncoating factor OPG117




• Molecule 1: Uncoating factor OPG117





- Molecule 1: Uncoating factor OPG117



ALA TYR SER SER ILE VAL ARG ILE PHE ASP MET ASP VAL ASP ASP LEU LEU ALA ALA LEU LEU GLU ILE ILE ASP ASP TYR TYR THR ALA ALA GLN ASP PHE PHE ILE ILE ILE GLU GLU VAL VAL ASN CYS VAL VAL ALA ALA PHE PHE THR THR GLU GLU CYS GLY GLY ALA ALA ILE HIS GLU ASN VAL ILE LYS SER MET ARG ARG ASN ASN




● Molecule 2: DNA polymerase






• Molecule 3: DNA polymerase processivity factor component OPG148

Chain C:



• Molecule 4: Uracil-DNA glycosylase

Chain B:



• Molecule 5: DNA (5'-D(P*TP*TP*TP*TP*TP*TP*TP*TP*TP*TP*TP*TP*T)-3')

Chain T:



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	56874	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	FEI FALCON IV (4k x 4k)	Depositor
Maximum map value	2.469	Depositor
Minimum map value	-0.006	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.024	Depositor
Recommended contour level	0.04	Depositor
Map size ($\text{\AA}$)	378.88, 378.88, 378.88	wwPDB
Map dimensions	512, 512, 512	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	0.74, 0.74, 0.74	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	D	0.10	0/5579	0.25	0/7542
1	E	0.10	0/4684	0.25	0/6324
1	H	0.08	0/2914	0.23	0/3934
1	I	0.08	0/2915	0.22	0/3935
1	J	0.07	0/2684	0.22	0/3614
1	K	0.07	0/2912	0.21	0/3927
2	A	0.08	0/8199	0.24	1/11070 (0.0%)
3	C	0.09	0/3199	0.24	0/4306
4	B	0.09	0/1823	0.26	0/2479
5	T	0.28	0/503	0.83	0/770
All	All	0.09	0/35412	0.26	1/47901 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	948	LYS	N-CA-C	-5.18	108.71	114.62

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	5469	0	5471	58	0
1	E	4599	0	4592	64	0
1	H	2855	0	2892	20	0
1	I	2857	0	2889	15	0
1	J	2635	0	2674	21	0
1	K	2856	0	2892	22	0
2	A	8031	0	7999	78	0
3	C	3141	0	3151	50	0
4	B	1773	0	1764	26	0
5	T	460	0	279	15	0
6	D	1	0	0	0	0
All	All	34677	0	34603	348	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 348 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:1000:ARG:NH1	5:T:11:DT:C7	1.69	1.56
2:A:1000:ARG:NH1	5:T:11:DT:H72	1.31	1.30
2:A:1000:ARG:NH1	5:T:11:DT:H73	1.45	1.07
1:D:151:LYS:NZ	5:T:12:DT:OP2	2.00	0.94
1:H:681:ILE:HG13	1:H:682:PRO:HD3	1.48	0.94

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	D	666/785 (85%)	648 (97%)	17 (3%)	1 (0%)	43 76
1	E	552/785 (70%)	537 (97%)	15 (3%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	345/785 (44%)	339 (98%)	6 (2%)	0	100	100
1	I	345/785 (44%)	338 (98%)	7 (2%)	0	100	100
1	J	314/785 (40%)	307 (98%)	7 (2%)	0	100	100
1	K	343/785 (44%)	336 (98%)	7 (2%)	0	100	100
2	A	973/1006 (97%)	944 (97%)	28 (3%)	1 (0%)	48	81
3	C	371/426 (87%)	359 (97%)	12 (3%)	0	100	100
4	B	216/218 (99%)	208 (96%)	8 (4%)	0	100	100
All	All	4125/6360 (65%)	4016 (97%)	107 (3%)	2 (0%)	100	100

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	255	ILE
2	A	487	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	624/725 (86%)	620 (99%)	4 (1%)	78	80
1	E	522/725 (72%)	508 (97%)	14 (3%)	39	60
1	H	324/725 (45%)	318 (98%)	6 (2%)	50	67
1	I	324/725 (45%)	318 (98%)	6 (2%)	50	67
1	J	298/725 (41%)	296 (99%)	2 (1%)	76	79
1	K	322/725 (44%)	318 (99%)	4 (1%)	63	73
2	A	909/928 (98%)	901 (99%)	8 (1%)	70	76
3	C	358/396 (90%)	354 (99%)	4 (1%)	65	74
4	B	200/200 (100%)	197 (98%)	3 (2%)	57	70
All	All	3881/5874 (66%)	3830 (99%)	51 (1%)	59	72

5 of 51 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	106	ILE
1	I	507	THR
1	K	535	VAL
3	C	149	VAL
4	B	117	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 46 such sidechains are listed below:

Mol	Chain	Res	Type
4	B	55	GLN
1	H	541	ASN
4	B	147	HIS
1	I	629	HIS
1	H	629	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
5	T	2

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	T	13:DT	O3'	34:DT	P	55.39
1	T	1:DT	O3'	6:DT	P	27.58


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-49747. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y



Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 256




Y Index: 256




Z Index: 256

6.2.2 Raw map



X Index: 256



Y Index: 256



Z Index: 256

The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 239




Y Index: 240




Z Index: 299


6.3.2 Raw map



X Index: 0



Y Index: 0




Z Index: 0


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y



Z


The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



X




Y




Z

The images above show the 3D surface view of the map at the recommended contour level 0.04. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.5.2 Raw map



X



Y



Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_49747_msk_1.map [i](#)



X




Y




Z


6.6.2 emd_49747_msk_2.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 285 nm^3 ; this corresponds to an approximate mass of 258 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of $0.244 \text{ \AA}^{-1}$

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.244 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	4.10	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	6.18	8.64	6.64

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 6.18 differs from the reported value 4.1 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-49747 and PDB model 9NSP. Per-residue inclusion information can be found in section [3](#) on page [6](#).


9.1 Map-model overlay [i](#)



X



Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.04 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.04).

9.4 Atom inclusion [i](#)



At the recommended contour level, 81% of all backbone atoms, 64% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.04) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.6430	0.1520
A	0.8400	0.3470
B	0.6870	0.1980
C	0.7120	0.1090
D	0.6600	0.1590
E	0.4030	0.0360
H	0.5540	0.0640
I	0.5790	0.0700
J	0.5800	0.0640
K	0.5640	0.0560
T	0.5960	0.1200

1.0

0.0

<0.0