



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2026 – 10:19 PM UTC

PDB ID : 1O46 / pdb_00001o46
Title : CRYSTAL STRUCTURE OF SH2 IN COMPLEX WITH RU90395.
Authors : Lange, G.; Loenze, P.; Liesum, A.
Deposited on : 2003-06-15
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	108	 53% 39% 6% ..

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 1045 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTO-ONCOGENE TYROSINE-PROTEIN KINASE SRC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	106	Total	C	N	O	S	0	0	0
			855	538	153	161	3			

- Molecule 2 is 2-{4-[2-ACETYLAMINO-2-(1-BIPHENYL-4-YLMETHYL-2-OXO-AZEPAN-3-YLCARBAMOYL)-ETHYL]-2-METHOXYCARBONYL-PHENYL}-2-FLUORO-MALONIC ACID (CCD ID: 903) (formula: C₃₅H₃₆FN₃O₉).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	0	0
			48	35	1	3	9		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	142	Total	O	0	0
			142	142		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: PROTO-ONCOGENE TYROSINE-PROTEIN KINASE SRC

Chain A: 



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	26.59Å 58.55Å 64.19Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 2.00	Depositor
% Data completeness (in resolution range)	95.3 (8.00-2.00)	Depositor
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.192 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	1045	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 903

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.56	8/872 (0.9%)	1.83	29/1175 (2.5%)

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	47	VAL	CA-CB	6.77	1.64	1.54
1	A	14	ARG	CA-C	6.06	1.60	1.52
1	A	43	TYR	CA-C	-5.82	1.45	1.52
1	A	17	SER	C-O	5.74	1.30	1.24
1	A	59	LYS	CA-C	-5.61	1.45	1.52
1	A	19	ARG	C-O	-5.54	1.17	1.24
1	A	26	ASN	C-N	5.18	1.39	1.33
1	A	103	VAL	CA-C	5.03	1.58	1.52

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	25	GLU	CA-CB-CG	-12.83	88.44	114.10
1	A	83	GLN	CA-CB-CG	-10.69	92.71	114.10
1	A	104	CYS	CB-CA-C	-8.89	95.66	109.62
1	A	67	ASP	CB-CA-C	8.11	125.29	109.72
1	A	68	SER	N-CA-C	-7.98	102.81	112.54
1	A	21	LEU	N-CA-C	7.69	120.74	111.82
1	A	69	GLY	N-CA-C	7.59	125.09	114.85
1	A	59	LYS	CA-CB-CG	-7.19	99.73	114.10
1	A	54	LYS	CB-CA-C	-6.69	96.88	109.72
1	A	38	THR	N-CA-C	6.27	121.99	113.97
1	A	105	PRO	CB-CA-C	-6.09	102.98	110.95
1	A	91	LYS	CB-CA-C	-6.04	98.93	109.07
1	A	14	ARG	NE-CZ-NH1	-5.91	115.59	121.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	23	ASN	CB-CA-C	5.81	119.11	109.53
1	A	78	GLN	N-CA-C	5.69	118.96	110.14
1	A	19	ARG	CA-C-O	-5.67	114.41	120.42
1	A	31	PHE	N-CA-C	5.52	117.23	108.79
1	A	16	GLU	CA-CB-CG	5.50	125.10	114.10
1	A	18	GLU	CA-CB-CG	-5.47	103.16	114.10
1	A	11	LYS	N-CA-C	5.36	117.91	109.39
1	A	51	ASP	CB-CA-C	5.34	122.58	110.67
1	A	27	PRO	CA-C-O	-5.31	115.54	122.12
1	A	88	TYR	CA-C-O	5.29	126.37	120.82
1	A	19	ARG	N-CA-C	-5.26	105.63	111.36
1	A	70	GLY	N-CA-C	5.24	119.71	112.37
1	A	14	ARG	CA-CB-CG	-5.23	103.64	114.10
1	A	5	GLU	CB-CA-C	-5.21	102.03	109.84
1	A	2	ILE	N-CA-C	5.12	116.45	110.62
1	A	43	TYR	CB-CA-C	-5.08	100.23	110.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	855	0	845	17	0
2	A	48	0	35	8	0
3	A	142	0	0	5	0
All	All	1045	0	880	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:300:903:C40	2:A:300:903:F80	1.59	1.38

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:83:GLN:HE21	1:A:83:GLN:H	1.17	0.88
2:A:300:903:F80	2:A:300:903:C41	2.15	0.83
1:A:64:ARG:HD3	3:A:339:HOH:O	1.80	0.82
1:A:44:CYS:SG	2:A:300:903:H3	2.20	0.81
1:A:102:THR:HB	3:A:424:HOH:O	1.90	0.71
1:A:59:LYS:HE2	3:A:336:HOH:O	1.90	0.70
2:A:300:903:H811	2:A:300:903:H152	1.80	0.63
1:A:83:GLN:H	1:A:83:GLN:NE2	1.92	0.61
1:A:65:LYS:HE2	1:A:66:LEU:O	2.01	0.60
1:A:81:SER:HB2	1:A:83:GLN:NE2	2.21	0.56
2:A:300:903:C6	2:A:300:903:N12	2.69	0.55
2:A:300:903:H152	2:A:300:903:C81	2.37	0.54
2:A:300:903:N12	2:A:300:903:H6	2.25	0.52
2:A:300:903:F80	2:A:300:903:O43	2.18	0.51
1:A:72:TYR:HB3	1:A:78:GLN:HG2	1.96	0.48
1:A:32:LEU:HD12	1:A:32:LEU:C	2.39	0.47
1:A:1:SER:N	3:A:433:HOH:O	2.48	0.46
1:A:8:TYR:CZ	1:A:10:GLY:HA2	2.50	0.46
1:A:39:THR:HB	1:A:42:ALA:HB3	1.97	0.46
1:A:51:ASP:O	1:A:55:GLY:N	2.51	0.43
1:A:50:PHE:HA	1:A:55:GLY:O	2.21	0.41
1:A:105:PRO:HG2	3:A:358:HOH:O	2.21	0.40
1:A:71:PHE:CD2	1:A:82:LEU:HD13	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	104/108 (96%)	98 (94%)	6 (6%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	93/95 (98%)	89 (96%)	4 (4%)	26	25

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	SER
1	A	36	SER
1	A	83	GLN
1	A	102	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	78	GLN
1	A	83	GLN
1	A	92	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	903	A	300	-	48,51,51	3.98	27 (56%)	61,72,72	3.79	30 (49%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	903	A	300	-	-	7/44/63/63	0/4/4/4

All (27) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	300	903	C17-N18	18.50	1.53	1.35
2	A	300	903	C5-C4	6.26	1.50	1.40
2	A	300	903	C16-C17	5.59	1.61	1.53
2	A	300	903	C3-C4	5.26	1.46	1.39
2	A	300	903	C6-C1	5.23	1.48	1.39
2	A	300	903	O42-C41	4.92	1.48	1.30
2	A	300	903	C15-C13	4.86	1.60	1.50
2	A	300	903	C32-C31	4.60	1.48	1.39
2	A	300	903	C5-C37	4.40	1.59	1.50
2	A	300	903	O76-C75	-4.35	1.14	1.30
2	A	300	903	O38-C81	4.30	1.54	1.45
2	A	300	903	C30-C23	4.03	1.46	1.38
2	A	300	903	C3-C2	3.98	1.45	1.38
2	A	300	903	C9-N10	3.77	1.42	1.34
2	A	300	903	C35-C36	3.53	1.44	1.38
2	A	300	903	O19-C17	3.29	1.28	1.22
2	A	300	903	C30-C29	3.11	1.43	1.38
2	A	300	903	F80-C40	3.00	1.59	1.41

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	300	903	C27-C26	2.90	1.43	1.38
2	A	300	903	C22-C23	-2.85	1.46	1.51
2	A	300	903	C8-N12	2.78	1.51	1.45
2	A	300	903	C35-C34	2.72	1.44	1.38
2	A	300	903	C36-C31	2.51	1.44	1.39
2	A	300	903	C33-C32	2.50	1.43	1.38
2	A	300	903	O11-C9	2.33	1.27	1.23
2	A	300	903	O77-C75	2.24	1.29	1.22
2	A	300	903	O38-C37	2.23	1.38	1.33

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	300	903	C81-O38-C37	18.93	152.19	115.81
2	A	300	903	C25-C16-C17	10.05	124.47	111.65
2	A	300	903	F80-C40-C4	9.13	123.95	105.86
2	A	300	903	C23-C22-N18	-6.10	103.56	113.15
2	A	300	903	C35-C36-C31	-5.56	114.21	120.54
2	A	300	903	C25-C16-N10	-5.50	99.69	109.76
2	A	300	903	C16-N10-C9	-5.11	110.66	121.65
2	A	300	903	C7-C1-C6	-3.88	113.77	120.43
2	A	300	903	C8-C9-N10	-3.26	109.68	116.63
2	A	300	903	C27-C26-C23	-3.25	116.73	121.00
2	A	300	903	O76-C75-O77	-2.90	114.58	123.86
2	A	300	903	C6-C5-C4	-2.86	116.61	120.23
2	A	300	903	C3-C2-C1	-2.84	117.27	121.00
2	A	300	903	C34-C35-C36	2.82	123.72	120.24
2	A	300	903	C17-C16-N10	2.81	114.65	108.73
2	A	300	903	C36-C31-C32	2.79	122.68	117.68
2	A	300	903	C2-C1-C6	2.71	122.28	118.55
2	A	300	903	C20-N18-C17	2.64	125.50	122.42
2	A	300	903	C4-C5-C37	2.62	129.53	124.49
2	A	300	903	C1-C7-C8	-2.61	106.42	113.36
2	A	300	903	O38-C37-O39	2.56	128.43	123.46
2	A	300	903	C29-C28-C31	-2.53	116.90	121.25
2	A	300	903	C30-C23-C26	2.52	121.98	118.23
2	A	300	903	O11-C9-N10	2.41	127.27	122.96
2	A	300	903	C9-C8-N12	-2.37	104.70	111.11
2	A	300	903	O38-C37-C5	-2.34	108.42	112.31
2	A	300	903	C6-C5-C37	-2.34	113.78	118.41
2	A	300	903	O19-C17-C16	2.33	125.64	121.04
2	A	300	903	O42-C41-O43	2.26	131.10	123.86

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	300	903	C26-C27-C28	2.10	123.81	121.12

There are no chirality outliers.

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	300	903	C5-C4-C40-F80
2	A	300	903	C3-C4-C40-F80
2	A	300	903	C3-C4-C40-C41
2	A	300	903	C4-C40-C75-O77
2	A	300	903	O38-C37-C5-C6
2	A	300	903	O38-C37-C5-C4
2	A	300	903	C4-C40-C75-O76

There are no ring outliers.

1 monomer is involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	300	903	8	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.