



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 15, 2026 – 12:57 AM UTC

PDB ID : 4OBU / pdb_00004obu
Title : Ruminococcus gnavus tryptophan decarboxylase RUMGNA_01526 (apo)
Authors : Van Benschoten, A.H.; Fraser, J.S.
Deposited on : 2014-01-07
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	490	
1	B	490	
1	C	490	
1	E	490	
1	F	490	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	490	 85%9%6%
1	H	490	 84%10%6%
1	U	490	 86%9%6%

2 Entry composition


There are 3 unique types of molecules in this entry. The entry contains 29598 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pyridoxal-dependent decarboxylase domain protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	E	462	Total	C	N	O	S	0	0	0
			3633	2310	618	687	18			
1	A	462	Total	C	N	O	S	0	0	0
			3633	2310	618	687	18			
1	U	462	Total	C	N	O	S	0	0	0
			3633	2310	618	687	18			
1	G	462	Total	C	N	O	S	0	0	0
			3633	2310	618	687	18			
1	H	462	Total	C	N	O	S	0	0	0
			3633	2310	618	687	18			
1	F	462	Total	C	N	O	S	0	0	0
			3633	2310	618	687	18			
1	B	462	Total	C	N	O	S	0	0	0
			3633	2310	618	687	18			
1	C	462	Total	C	N	O	S	0	0	0
			3633	2310	618	687	18			

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (CCD ID: PLP) (formula: C₈H₁₀NO₆P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	E	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	A	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	U	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	G	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	H	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	F	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	B	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	C	1	Total	C	N	O	P	0	0
			15	8	1	5	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	E	56	Total	O	0	0
			56	56		
3	A	49	Total	O	0	0
			49	49		
3	U	57	Total	O	0	0
			57	57		
3	G	46	Total	O	0	0
			46	46		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	H	52	Total 52	O 52	0	0
3	F	73	Total 73	O 73	0	0
3	B	45	Total 45	O 45	0	0
3	C	36	Total 36	O 36	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Pyridoxal-dependent decarboxylase domain protein




- Molecule 1: Pyridoxal-dependent decarboxylase domain protein



- Molecule 1: Pyridoxal-dependent decarboxylase domain protein




- Molecule 1: Pyridoxal-dependent decarboxylase domain protein






- Molecule 1: Pyridoxal-dependent decarboxylase domain protein

Chain H: 84% 10% 6%





- Molecule 1: Pyridoxal-dependent decarboxylase domain protein

Chain F: 84% 10% 6%





- Molecule 1: Pyridoxal-dependent decarboxylase domain protein

Chain B: 84% 10% 6%



- Molecule 1: Pyridoxal-dependent decarboxylase domain protein

Chain C: 82% 12% 6%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	58.63Å 145.77Å 165.07Å 72.85° 88.84° 88.30°	Depositor
Resolution (Å)	79.37 – 2.80 79.37 – 2.80	Depositor EDS
% Data completeness (in resolution range)	94.7 (79.37-2.80) 80.1 (79.37-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.88 (at 2.82Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.1_1168)	Depositor
R, R_{free}	0.231 , 0.257 0.232 , 0.258	Depositor DCC
R_{free} test set	6088 reflections (4.76%)	wwPDB-VP
Wilson B-factor (Å ²)	35.7	Xtriage
Anisotropy	0.146	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 21.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	0.105 for h,-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	29598	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 24.28 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.9546e-03.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.28	0/3713	0.71	1/5031 (0.0%)
1	B	0.28	0/3713	0.71	2/5031 (0.0%)
1	C	0.29	0/3713	0.71	4/5031 (0.1%)
1	E	0.28	0/3713	0.71	0/5031
1	F	0.28	0/3713	0.71	2/5031 (0.0%)
1	G	0.29	0/3713	0.72	3/5031 (0.1%)
1	H	0.28	0/3713	0.71	3/5031 (0.1%)
1	U	0.28	0/3713	0.71	3/5031 (0.1%)
All	All	0.28	0/29704	0.71	18/40248 (0.0%)

There are no bond length outliers.

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	72	VAL	N-CA-C	9.55	119.56	110.30
1	C	423	GLU	N-CA-C	5.72	117.19	111.07
1	U	72	VAL	N-CA-C	5.62	115.82	110.42
1	C	461	HIS	CA-C-N	5.44	125.11	119.56
1	C	461	HIS	C-N-CA	5.44	125.11	119.56

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3633	0	3604	21	0
1	B	3633	0	3604	25	1
1	C	3633	0	3604	36	0
1	E	3633	0	3604	25	0
1	F	3633	0	3604	25	1
1	G	3633	0	3604	25	1
1	H	3633	0	3604	31	0
1	U	3633	0	3604	22	0
2	A	15	0	7	0	0
2	B	15	0	7	1	0
2	C	15	0	7	0	0
2	E	15	0	7	1	0
2	F	15	0	7	2	0
2	G	15	0	7	0	0
2	H	15	0	7	1	0
2	U	15	0	7	1	0
3	A	49	0	0	1	0
3	B	45	0	0	1	0
3	C	36	0	0	7	0
3	E	56	0	0	1	0
3	F	73	0	0	1	1
3	G	46	0	0	1	0
3	H	52	0	0	3	0
3	U	57	0	0	0	0
All	All	29598	0	28888	203	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 203 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:421:THR:O	3:C:629:HOH:O	1.89	0.89
1:B:326:PHE:O	1:B:330:HIS:ND1	2.20	0.74
1:H:326:PHE:O	1:H:330:HIS:ND1	2.20	0.74
1:F:326:PHE:O	1:F:330:HIS:ND1	2.23	0.71
1:G:326:PHE:O	1:G:330:HIS:ND1	2.23	0.70

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:23:THR:OG1	1:B:35:ASP:OD2[1_655]	2.09	0.11
1:G:335:TYR:OH	3:F:611:HOH:O[1_455]	2.12	0.08

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	458/490 (94%)	441 (96%)	17 (4%)	0	100	100
1	B	458/490 (94%)	441 (96%)	17 (4%)	0	100	100
1	C	458/490 (94%)	441 (96%)	17 (4%)	0	100	100
1	E	458/490 (94%)	441 (96%)	17 (4%)	0	100	100
1	F	458/490 (94%)	441 (96%)	17 (4%)	0	100	100
1	G	458/490 (94%)	441 (96%)	17 (4%)	0	100	100
1	H	458/490 (94%)	442 (96%)	16 (4%)	0	100	100
1	U	458/490 (94%)	442 (96%)	16 (4%)	0	100	100
All	All	3664/3920 (94%)	3530 (96%)	134 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	395/422 (94%)	392 (99%)	3 (1%)	73	90

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	395/422 (94%)	393 (100%)	2 (0%)	81	93
1	C	395/422 (94%)	392 (99%)	3 (1%)	73	90
1	E	395/422 (94%)	392 (99%)	3 (1%)	73	90
1	F	395/422 (94%)	392 (99%)	3 (1%)	73	90
1	G	395/422 (94%)	392 (99%)	3 (1%)	73	90
1	H	395/422 (94%)	392 (99%)	3 (1%)	73	90
1	U	395/422 (94%)	391 (99%)	4 (1%)	68	88
All	All	3160/3376 (94%)	3136 (99%)	24 (1%)	73	90

5 of 24 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	317	VAL
1	F	317	VAL
1	F	119	ILE
1	F	351	ILE
1	U	119	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 35 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	310	GLN
1	B	385	GLN
1	C	219	GLN
1	U	407	GLN
1	U	385	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PLP	B	501	-	15,15,16	1.04	1 (6%)	21,22,23	1.42	2 (9%)
2	PLP	G	501	-	15,15,16	1.02	1 (6%)	21,22,23	1.35	2 (9%)
2	PLP	A	501	-	15,15,16	1.06	1 (6%)	21,22,23	1.38	2 (9%)
2	PLP	C	501	-	15,15,16	1.04	1 (6%)	21,22,23	1.35	2 (9%)
2	PLP	H	501	-	15,15,16	1.04	1 (6%)	21,22,23	1.35	2 (9%)
2	PLP	E	501	-	15,15,16	1.05	1 (6%)	21,22,23	1.45	2 (9%)
2	PLP	F	501	-	15,15,16	1.04	1 (6%)	21,22,23	1.43	2 (9%)
2	PLP	U	501	-	15,15,16	1.05	1 (6%)	21,22,23	1.28	2 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLP	B	501	-	-	2/6/6/8	0/1/1/1
2	PLP	G	501	-	-	3/6/6/8	0/1/1/1
2	PLP	A	501	-	-	3/6/6/8	0/1/1/1
2	PLP	C	501	-	-	3/6/6/8	0/1/1/1
2	PLP	H	501	-	-	3/6/6/8	0/1/1/1
2	PLP	E	501	-	-	3/6/6/8	0/1/1/1
2	PLP	F	501	-	-	2/6/6/8	0/1/1/1
2	PLP	U	501	-	-	2/6/6/8	0/1/1/1

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	501	PLP	C2-N1	2.47	1.38	1.33
2	E	501	PLP	C2-N1	2.46	1.38	1.33
2	A	501	PLP	C2-N1	2.46	1.38	1.33
2	U	501	PLP	C2-N1	2.46	1.38	1.33
2	C	501	PLP	C2-N1	2.42	1.38	1.33

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	501	PLP	C4A-C4-C3	-3.91	114.01	120.52
2	E	501	PLP	C4A-C4-C3	-3.73	114.31	120.52
2	H	501	PLP	C4A-C4-C3	-3.64	114.46	120.52
2	B	501	PLP	O4P-C5A-C5	3.61	116.13	109.36
2	C	501	PLP	C4A-C4-C3	-3.53	114.63	120.52

There are no chirality outliers.

5 of 21 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	E	501	PLP	C4-C5-C5A-O4P
2	E	501	PLP	C6-C5-C5A-O4P
2	A	501	PLP	C4-C5-C5A-O4P
2	A	501	PLP	C6-C5-C5A-O4P
2	G	501	PLP	C4-C5-C5A-O4P

There are no ring outliers.

5 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	501	PLP	1	0
2	H	501	PLP	1	0
2	E	501	PLP	1	0
2	F	501	PLP	2	0
2	U	501	PLP	1	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	462/490 (94%)	-1.61	0 100 100	1, 14, 63, 107	0
1	B	462/490 (94%)	-1.60	0 100 100	1, 14, 63, 110	0
1	C	462/490 (94%)	-1.57	0 100 100	1, 15, 60, 109	0
1	E	462/490 (94%)	-1.61	0 100 100	1, 15, 62, 113	0
1	F	462/490 (94%)	-1.60	0 100 100	1, 13, 62, 112	0
1	G	462/490 (94%)	-1.62	0 100 100	1, 14, 60, 111	0
1	H	462/490 (94%)	-1.59	0 100 100	1, 16, 64, 105	0
1	U	462/490 (94%)	-1.58	0 100 100	1, 15, 64, 110	0
All	All	3696/3920 (94%)	-1.60	0 100 100	1, 15, 63, 113	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PLP	E	501	15/16	0.99	0.04	24,44,79,80	0
2	PLP	A	501	15/16	0.99	0.04	2,49,70,78	0
2	PLP	U	501	15/16	0.99	0.04	4,38,74,80	0
2	PLP	B	501	15/16	0.99	0.04	8,43,64,67	0
2	PLP	H	501	15/16	1.00	0.03	7,28,73,74	0
2	PLP	F	501	15/16	1.00	0.03	6,31,65,66	0
2	PLP	G	501	15/16	1.00	0.03	6,32,73,76	0
2	PLP	C	501	15/16	1.00	0.03	10,35,66,66	0

6.5 Other polymers [i](#)

There are no such residues in this entry.