



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 03:59 PM UTC

PDB ID : 3ORZ / pdb_00003orz
Title : PDK1 mutant bound to allosteric disulfide fragment activator 2A2
Authors : Sadowsky, J.D.; Wells, J.A.
Deposited on : 2010-09-08
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	316	
1	B	316	
1	C	316	
1	D	316	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 9976 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-phosphoinositide-dependent protein kinase 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	277	Total	C	N	O	P	S	0	1	0
			2238	1456	362	411	1	8			
1	B	279	Total	C	N	O	P	S	0	0	0
			2241	1457	365	410	1	8			
1	C	277	Total	C	N	O	P	S	0	0	0
			2229	1451	362	407	1	8			
1	D	279	Total	C	N	O	P	S	0	0	0
			2251	1461	367	414	1	8			

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	44	GLY	-	expression tag	UNP O15530
A	45	ALA	-	expression tag	UNP O15530
A	46	MET	-	expression tag	UNP O15530
A	47	ASP	-	expression tag	UNP O15530
A	48	PRO	-	expression tag	UNP O15530
A	49	GLU	-	expression tag	UNP O15530
A	50	PHE	-	expression tag	UNP O15530
A	148	CYS	THR	engineered mutation	UNP O15530
B	44	GLY	-	expression tag	UNP O15530
B	45	ALA	-	expression tag	UNP O15530
B	46	MET	-	expression tag	UNP O15530
B	47	ASP	-	expression tag	UNP O15530
B	48	PRO	-	expression tag	UNP O15530
B	49	GLU	-	expression tag	UNP O15530
B	50	PHE	-	expression tag	UNP O15530
B	148	CYS	THR	engineered mutation	UNP O15530
C	44	GLY	-	expression tag	UNP O15530
C	45	ALA	-	expression tag	UNP O15530
C	46	MET	-	expression tag	UNP O15530
C	47	ASP	-	expression tag	UNP O15530
C	48	PRO	-	expression tag	UNP O15530

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
C	49	GLU	-	expression tag	UNP O15530
C	50	PHE	-	expression tag	UNP O15530
C	148	CYS	THR	engineered mutation	UNP O15530
D	44	GLY	-	expression tag	UNP O15530
D	45	ALA	-	expression tag	UNP O15530
D	46	MET	-	expression tag	UNP O15530
D	47	ASP	-	expression tag	UNP O15530
D	48	PRO	-	expression tag	UNP O15530
D	49	GLU	-	expression tag	UNP O15530
D	50	PHE	-	expression tag	UNP O15530
D	148	CYS	THR	engineered mutation	UNP O15530

- Molecule 2 is 4-[4-(3-chlorophenyl)piperazin-1-yl]-4-oxobutane-1-thiol (CCD ID: 2A2) (formula: C₁₄H₁₉ClN₂OS).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total 19	C 14	Cl 1	N 2	O 1	S 1	0	0
2	B	1	Total 19	C 14	Cl 1	N 2	O 1	S 1	0	0
2	C	1	Total 19	C 14	Cl 1	N 2	O 1	S 1	0	0
2	D	1	Total 19	C 14	Cl 1	N 2	O 1	S 1	0	0

- Molecule 3 is 3-(1H-INDOL-3-YL)-4-{1-[2-(1-METHYLPYRROLIDIN-2-YL)ETHYL]-1H-INDOL-3-YL}-1H-PYRROLE-2,5-DIONE (CCD ID: BI4) (formula: C₂₇H₂₆N₄O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			33	27	4	2		
3	B	1	Total	C	N	O	0	0
			33	27	4	2		
3	C	1	Total	C	N	O	0	0
			33	27	4	2		
3	D	1	Total	C	N	O	0	0
			33	27	4	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	209	Total	O	0	0
			209	209		
4	B	203	Total	O	0	0
			203	203		
4	C	200	Total	O	0	0
			200	200		
4	D	197	Total	O	0	0
			197	197		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 3-phosphoinositide-dependent protein kinase 1



- Molecule 1: 3-phosphoinositide-dependent protein kinase 1




- Molecule 1: 3-phosphoinositide-dependent protein kinase 1



- Molecule 1: 3-phosphoinositide-dependent protein kinase 1

Chain D: 70% 16% • 12%



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	243.13Å 116.12Å 37.70Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.98 – 2.00 41.98 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.3 (41.98-2.00) 98.8 (41.98-2.00)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.14 (at 2.00Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 2010_01_09_2330)	Depositor
R, R_{free}	0.218 , 0.265 0.216 , 0.260	Depositor DCC
R_{free} test set	3528 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	19.5	Xtriage
Anisotropy	0.474	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 30.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.39$, $\langle L^2 \rangle = 0.23$	Xtriage
Estimated twinning fraction	0.458 for -h,-k,l	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	9976	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 99.49 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.7895e-13. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 2A2, BI4, SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.63	4/2286 (0.2%)	0.80	2/3087 (0.1%)
1	B	0.60	3/2284 (0.1%)	0.78	1/3082 (0.0%)
1	C	0.59	3/2274 (0.1%)	0.82	5/3071 (0.2%)
1	D	0.58	1/2294 (0.0%)	0.81	1/3095 (0.0%)
All	All	0.60	11/9138 (0.1%)	0.80	9/12335 (0.1%)

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	120	LYS	C-O	-6.25	1.16	1.24
1	C	350	LEU	C-O	-6.06	1.17	1.24
1	C	150	GLN	C-O	-5.90	1.16	1.23
1	B	214	ASN	C-O	-5.70	1.17	1.23
1	A	203	HIS	C-O	-5.59	1.17	1.24

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	124	VAL	CB-CA-C	-10.12	103.62	114.35
1	B	124	VAL	CB-CA-C	-8.75	105.08	114.35
1	C	348	GLU	CA-C-N	-8.45	111.71	123.03
1	C	348	GLU	C-N-CA	-8.45	111.71	123.03
1	A	153	GLU	N-CA-C	6.40	118.34	111.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2238	0	2221	28	0
1	B	2241	0	2223	41	1
1	C	2229	0	2213	34	0
1	D	2251	0	2235	37	1
2	A	19	0	18	0	0
2	B	19	0	18	3	0
2	C	19	0	18	0	0
2	D	19	0	18	1	0
3	A	33	0	26	5	0
3	B	33	0	26	5	0
3	C	33	0	26	7	0
3	D	33	0	26	6	0
4	A	209	0	0	1	0
4	B	203	0	0	3	0
4	C	200	0	0	2	0
4	D	197	0	0	4	0
All	All	9976	0	9068	164	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 164 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:73:GLN:HG2	1:B:74:PRO:HD2	1.50	0.91
1:A:202:ILE:HD12	1:A:228:LYS:HD2	1.67	0.76
1:C:214:ASN:ND2	1:C:216:ASP:OD2	2.21	0.72
1:C:276:VAL:O	1:C:304:LYS:HE2	1.91	0.71
1:B:129:ARG:HH11	1:B:229:VAL:H	1.39	0.70

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:287:GLU:OE1	1:D:287:GLU:OE1[4_554]	2.15	0.05

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	274/316 (87%)	262 (96%)	12 (4%)	0	100	100
1	B	274/316 (87%)	266 (97%)	8 (3%)	0	100	100
1	C	273/316 (86%)	266 (97%)	7 (3%)	0	100	100
1	D	274/316 (87%)	266 (97%)	8 (3%)	0	100	100
All	All	1095/1264 (87%)	1060 (97%)	35 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	239/273 (88%)	228 (95%)	11 (5%)	24	22
1	B	238/273 (87%)	224 (94%)	14 (6%)	18	14
1	C	237/273 (87%)	224 (94%)	13 (6%)	19	17
1	D	241/273 (88%)	226 (94%)	15 (6%)	16	13
All	All	955/1092 (88%)	902 (94%)	53 (6%)	19	17

5 of 53 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	167	LEU
1	C	286	ASN
1	D	289	LEU
1	C	196	LEU
1	C	249	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	117	HIS
1	D	247	GLN
1	D	286	ASN
1	D	240	ASN
1	B	286	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	SEP	D	241	1	8,9,10	1.64	1 (12%)	7,12,14	1.21	1 (14%)
1	SEP	A	241	1	8,9,10	1.68	1 (12%)	7,12,14	1.58	1 (14%)
1	SEP	B	241	1	8,9,10	1.67	1 (12%)	7,12,14	1.22	1 (14%)
1	SEP	C	241	1	8,9,10	1.69	1 (12%)	7,12,14	1.45	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	D	241	1	-	0/6/8/10	-
1	SEP	A	241	1	-	2/6/8/10	-
1	SEP	B	241	1	-	3/6/8/10	-
1	SEP	C	241	1	-	4/6/8/10	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	241	SEP	P-O1P	3.65	1.61	1.50
1	C	241	SEP	P-O1P	3.63	1.61	1.50
1	A	241	SEP	P-O1P	3.61	1.61	1.50
1	D	241	SEP	P-O1P	3.61	1.61	1.50

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	241	SEP	OG-CB-CA	3.63	111.67	108.14
1	C	241	SEP	OG-CB-CA	3.24	111.30	108.14
1	B	241	SEP	OG-CB-CA	2.64	110.71	108.14
1	D	241	SEP	OG-CB-CA	2.55	110.63	108.14

There are no chirality outliers.

5 of 9 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	241	SEP	N-CA-CB-OG
1	A	241	SEP	C-CA-CB-OG
1	B	241	SEP	CB-OG-P-O2P
1	C	241	SEP	C-CA-CB-OG
1	C	241	SEP	CB-OG-P-O1P

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	BI4	B	360	-	38,38,38	1.32	5 (13%)	47,56,56	3.12	17 (36%)
2	2A2	D	1	1	20,20,20	0.80	1 (5%)	26,26,26	1.41	3 (11%)
3	BI4	A	360	-	38,38,38	1.31	4 (10%)	47,56,56	3.05	16 (34%)
2	2A2	C	1	1	20,20,20	0.83	1 (5%)	26,26,26	1.45	5 (19%)
3	BI4	C	360	-	38,38,38	1.31	4 (10%)	47,56,56	3.06	16 (34%)
3	BI4	D	360	-	38,38,38	1.31	4 (10%)	47,56,56	3.14	17 (36%)
2	2A2	A	1	1	20,20,20	0.82	1 (5%)	26,26,26	1.46	5 (19%)
2	2A2	B	1	1	20,20,20	0.81	1 (5%)	26,26,26	1.45	5 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BI4	B	360	-	-	3/13/39/39	0/6/6/6
2	2A2	D	1	1	-	0/12/22/22	0/2/2/2
3	BI4	A	360	-	-	4/13/39/39	0/6/6/6
2	2A2	C	1	1	-	0/12/22/22	0/2/2/2
3	BI4	C	360	-	-	4/13/39/39	0/6/6/6
3	BI4	D	360	-	-	4/13/39/39	0/6/6/6
2	2A2	A	1	1	-	0/12/22/22	0/2/2/2
2	2A2	B	1	1	-	2/12/22/22	0/2/2/2

The worst 5 of 21 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	360	BI4	C23-C22	-4.41	1.37	1.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	360	BI4	C23-C22	-4.39	1.37	1.45
3	D	360	BI4	C15-C16	-4.19	1.38	1.45
3	B	360	BI4	C23-C22	-4.16	1.38	1.45
3	C	360	BI4	C15-C16	-4.14	1.38	1.45

The worst 5 of 84 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	360	BI4	C15-C16-C8	10.72	112.06	106.15
3	A	360	BI4	C15-C16-C8	10.65	112.02	106.15
3	C	360	BI4	C15-C16-C8	10.64	112.02	106.15
3	B	360	BI4	C15-C16-C8	10.44	111.90	106.15
3	C	360	BI4	C16-C8-N9	-8.39	104.54	110.22

There are no chirality outliers.

5 of 17 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	360	BI4	C25-C24-N6-C7
3	B	360	BI4	C20-C21-C22-C23
3	B	360	BI4	C25-C24-N6-C7
3	C	360	BI4	C25-C24-N6-C7
3	D	360	BI4	C20-C21-C22-C23


There are no ring outliers.

6 monomers are involved in 27 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	360	BI4	5	0
2	D	1	2A2	1	0
3	A	360	BI4	5	0
3	C	360	BI4	7	0
3	D	360	BI4	6	0
2	B	1	2A2	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be


highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand BI4 A 360




Bond lengths



Bond angles




Torsions



Rings


Ligand 2A2 C 1




Bond lengths




Bond angles




Torsions




Rings




Ligand BI4 D 360




Bond lengths



Bond angles




Torsions



Rings


Ligand 2A2 A 1




Bond lengths




Bond angles



Torsions



Rings



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	276/316 (87%)	-1.36	0 100 100	10, 20, 40, 49	1 (0%)
1	B	278/316 (87%)	-1.35	0 100 100	11, 21, 42, 52	0
1	C	276/316 (87%)	-1.36	0 100 100	10, 20, 40, 48	0
1	D	278/316 (87%)	-1.35	0 100 100	11, 21, 43, 52	0
All	All	1108/1264 (87%)	-1.35	0 100 100	10, 20, 42, 52	1 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	SEP	C	241	10/11	0.96	0.06	33,49,69,78	0
1	SEP	A	241	10/11	0.97	0.05	32,49,66,75	0
1	SEP	B	241	10/11	0.98	0.04	31,42,64,65	0
1	SEP	D	241	10/11	0.98	0.04	33,43,66,67	0

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	2A2	A	1	19/19	0.99	0.03	31,36,43,55	0
2	2A2	B	1	19/19	0.99	0.04	29,35,41,51	0
2	2A2	C	1	19/19	0.99	0.04	31,35,44,54	0
2	2A2	D	1	19/19	0.99	0.04	30,35,42,51	0
3	BI4	A	360	33/33	0.99	0.03	13,19,30,32	0
3	BI4	B	360	33/33	0.99	0.03	15,19,30,36	0
3	BI4	C	360	33/33	0.99	0.03	13,19,31,33	0
3	BI4	D	360	33/33	0.99	0.03	15,19,29,34	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around 2A2 B 1:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 2A2 C 1:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 2A2 D 1:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BI4 A 360:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BI4 B 360:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BI4 C 360:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around BI4 D 360:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.