



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 19, 2026 – 09:16 PM UTC

PDB ID : 6Q20 / pdb_00006q20
Title : Crystal structure of human 1E01 Fab in complex with influenza virus neuraminidase from A/Japan/305/1957 (H2N2)
Authors : Zhu, X.; Wilson, I.A.
Deposited on : 2019-08-06
Resolution : 2.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1190 (2.46-2.46)
Clashscore	190562	1229 (2.46-2.46)
Ramachandran outliers	187476	1218 (2.46-2.46)
Sidechain outliers	187428	1218 (2.46-2.46)
RSRZ outliers	180081	1190 (2.46-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	393	2% 86% 13% .
2	L	216	6% 83% 17%
3	H	240	8% 75% 20% 5%
4	B	7	29% 71%
5	C	2	100%

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 6596 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Neuraminidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	388	3014	1862	541	588	23	0	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	77	GLY	-	expression tag	UNP C7RWN0
A	78	SER	-	expression tag	UNP C7RWN0
A	79	PRO	-	expression tag	UNP C7RWN0
A	80	SER	-	expression tag	UNP C7RWN0
A	81	ARG	-	expression tag	UNP C7RWN0


- Molecule 2 is a protein called 1E01 Fab kappa light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	L	215	1661	1051	274	331	5	0	0	0

- Molecule 3 is a protein called 1E01 Fab IgG1 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	H	227	1731	1096	294	335	6	0	0	0

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	B	7	Total	C	N	O	0	0	0
			83	46	2	35			

- Molecule 5 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	C	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 7 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	2	Total	Ca	0	0
			2	2		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	41	Total 41	O 41	0	0
8	L	10	Total 10	O 10	0	0
8	H	12	Total 12	O 12	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Neuraminidase



- Molecule 2: 1E01 Fab kappa light chain



- Molecule 3: 1E01 Fab IgG1 heavy chain



- Molecule 4: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



MAG1	MAG2	MAN4	MAN5	MAN6	MAN7
	BR/A3				

- Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:  100%

MAG1	MAG2
------	------

4 Data and refinement statistics

Property	Value	Source
Space group	P 4 21 2	Depositor
Cell constants a, b, c, α , β , γ	160.45Å 160.45Å 84.17Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.04 – 2.45 47.04 – 2.45	Depositor EDS
% Data completeness (in resolution range)	99.3 (47.04-2.45) 99.5 (47.04-2.45)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.11 (at 2.45Å)	Xtriage
Refinement program	PHENIX (1.12_2829)	Depositor
R, R_{free}	0.217 , 0.272 0.220 , 0.274	Depositor DCC
R_{free} test set	2157 reflections (5.27%)	wwPDB-VP
Wilson B-factor (Å ²)	50.8	Xtriage
Anisotropy	0.882	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 45.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6596	wwPDB-VP
Average B, all atoms (Å ²)	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, CA, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.30	2/3088 (0.1%)	0.44	0/4193
2	L	0.11	0/1700	0.33	0/2311
3	H	0.14	0/1774	0.41	0/2415
All	All	0.23	2/6562 (0.0%)	0.40	0/8919

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	339	ASP	C-O	-7.24	1.17	1.24
1	A	339	ASP	CA-C	-5.98	1.46	1.52

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3014	0	2828	29	0
2	L	1661	0	1620	28	0
3	H	1731	0	1686	44	0
4	B	83	0	70	0	0
5	C	28	0	25	0	0
6	A	14	0	13	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	2	0	0	0	0
8	A	41	0	0	0	0
8	H	12	0	0	2	0
8	L	10	0	0	0	0
All	All	6596	0	6242	94	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (94) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:2:ILE:HD11	2:L:90:HIS:CE1	2.07	0.89
1:A:97:PHE:H	1:A:395:GLN:HE22	1.26	0.83
2:L:42:ASN:HA	3:H:105:ARG:HE	1.52	0.74
3:H:140:LEU:HD21	3:H:225:VAL:HG11	1.71	0.73
3:H:61:ASP:HA	3:H:64:LYS:HD2	1.72	0.72
3:H:52:ASN:ND2	3:H:55:ASP:OD1	2.24	0.71
2:L:108:ARG:NH1	2:L:109:THR:O	2.24	0.70
3:H:211:ASN:OD1	3:H:218:LYS:NZ	2.21	0.70
2:L:50:ALA:HB3	2:L:53:LEU:HD23	1.73	0.69
3:H:100(F):GLU:OE2	3:H:100(F):GLU:N	2.21	0.69
3:H:105:ARG:O	3:H:105:ARG:NH1	2.23	0.69
2:L:41:GLY:C	3:H:105:ARG:HH21	2.02	0.68
2:L:136:LEU:HB2	2:L:175:LEU:HB3	1.75	0.68
1:A:97:PHE:N	1:A:395:GLN:HE22	1.90	0.68
1:A:294:ASN:O	1:A:346:ASN:ND2	2.28	0.66
1:A:177:ALA:HB2	1:A:193:CYS:HB3	1.78	0.64
1:A:426:LEU:HD21	1:A:444:VAL:HG12	1.79	0.63
1:A:199:ARG:HG3	3:H:98:TRP:CZ2	2.33	0.62
2:L:78:LEU:HD22	2:L:82:ASP:HB2	1.81	0.61
1:A:296:LYS:O	1:A:340:PRO:HB2	2.02	0.60
1:A:208:ASP:O	1:A:210:ARG:NH1	2.34	0.60
3:H:68:THR:HB	3:H:81:GLN:HB3	1.84	0.60
1:A:238:THR:HG21	1:A:305:ILE:HG21	1.84	0.59
3:H:211:ASN:HD22	3:H:212:HIS:N	2.01	0.57
1:A:226:GLN:HE21	1:A:240:VAL:H	1.51	0.57
2:L:42:ASN:HA	3:H:105:ARG:NE	2.18	0.57
3:H:119:PRO:HB3	3:H:147:TYR:HB3	1.87	0.56
3:H:100(F):GLU:H	3:H:100(F):GLU:CD	2.14	0.55
3:H:162:ASN:HA	3:H:207:ILE:HG22	1.89	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:H:87:THR:HG23	3:H:110:THR:HA	1.89	0.54
1:A:153:ILE:HG12	1:A:156:ARG:HG3	1.89	0.54
1:A:132:PHE:HB3	1:A:158:LEU:HD11	1.89	0.54
3:H:211:ASN:CG	3:H:218:LYS:HG3	2.33	0.53
3:H:193:VAL:HG11	3:H:206:TYR:CZ	2.43	0.52
1:A:279:SER:HB3	1:A:409:ILE:HG22	1.91	0.52
2:L:2:ILE:HD11	2:L:90:HIS:ND1	2.25	0.52
3:H:105:ARG:NH2	8:H:301:HOH:O	2.37	0.52
2:L:6:GLN:HB3	2:L:102:THR:HG23	1.92	0.51
3:H:157:TRP:CZ3	3:H:208:CYS:HB3	2.45	0.51
1:A:359:ASP:OD1	1:A:380:ILE:HA	2.10	0.50
2:L:163:VAL:HG22	2:L:175:LEU:HD23	1.93	0.50
2:L:164:THR:HG22	2:L:165:GLU:O	2.12	0.50
2:L:90:HIS:ND1	2:L:93:SER:HB2	2.27	0.49
3:H:171:VAL:HG22	3:H:191:VAL:HG23	1.94	0.49
1:A:82:VAL:HG22	1:A:187:LYS:HG2	1.95	0.49
2:L:48:ILE:HG21	2:L:64:ALA:HB3	1.95	0.48
2:L:149:LYS:HB2	2:L:193:ALA:HB3	1.96	0.48
1:A:217:SER:OG	1:A:243:ASP:OD2	2.23	0.48
2:L:66:GLY:HA3	2:L:71:PHE:CD2	2.49	0.47
1:A:199:ARG:HG3	3:H:98:TRP:HZ2	1.79	0.47
3:H:213:LYS:NZ	3:H:218:LYS:HZ1	2.12	0.47
2:L:108:ARG:HH12	2:L:111:ALA:HB2	1.79	0.47
3:H:105:ARG:HH11	3:H:105:ARG:HG3	1.79	0.47
3:H:140:LEU:HD21	3:H:225:VAL:CG1	2.43	0.47
3:H:152:VAL:HG12	3:H:212:HIS:HB2	1.97	0.47
2:L:149:LYS:HG2	2:L:154:LEU:HD23	1.96	0.47
2:L:93:SER:O	2:L:95(A):LEU:N	2.48	0.46
1:A:97:PHE:H	1:A:395:GLN:NE2	2.05	0.46
3:H:2:VAL:HA	3:H:26:GLY:HA3	1.97	0.46
1:A:293:ASP:HB2	1:A:301:PRO:HG3	1.98	0.46
2:L:48:ILE:HD13	2:L:54:LEU:HA	1.97	0.46
1:A:226:GLN:HG3	1:A:239:VAL:HG23	1.98	0.45
3:H:60:THR:C	3:H:64:LYS:HZ2	2.24	0.45
1:A:93:GLN:OE1	1:A:356:ASN:ND2	2.40	0.45
3:H:172:HIS:HB2	3:H:190:VAL:HG22	1.99	0.45
2:L:78:LEU:HD23	2:L:78:LEU:HA	1.87	0.45
2:L:187:GLU:OE2	2:L:211:ARG:NH1	2.49	0.45
1:A:89:LYS:HB2	1:A:418:ILE:HG12	1.99	0.45
3:H:162:ASN:HA	3:H:207:ILE:CG2	2.47	0.45
3:H:40:PRO:HG2	3:H:43:LYS:HE3	1.99	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:183:LYS:O	2:L:187:GLU:HG2	2.17	0.44
1:A:103:ASP:OD2	1:A:105:SER:OG	2.24	0.44
3:H:43:LYS:NZ	8:H:304:HOH:O	2.51	0.44
2:L:95(A):LEU:HD21	3:H:61:ASP:OD1	2.18	0.44
2:L:32:TYR:O	2:L:90:HIS:HD2	2.01	0.43
3:H:162:ASN:OD1	3:H:207:ILE:HG22	2.18	0.43
3:H:39:GLN:HB2	3:H:45:LEU:HD23	2.01	0.43
1:A:169:LEU:HD23	1:A:169:LEU:HA	1.86	0.43
2:L:197:THR:HG22	2:L:204:PRO:HB3	2.01	0.43
3:H:105:ARG:NH1	3:H:105:ARG:HG3	2.33	0.43
2:L:6:GLN:HG3	2:L:100:PRO:HD2	2.00	0.42
1:A:235:GLY:O	1:A:258:LYS:HG3	2.20	0.42
1:A:295:TRP:CG	1:A:296:LYS:HG3	2.55	0.42
3:H:154:VAL:HA	3:H:209:ASN:O	2.20	0.42
3:H:52:ASN:OD1	3:H:54:GLY:N	2.45	0.41
1:A:286:ASP:HA	1:A:307:MET:HG2	2.01	0.41
3:H:38:ARG:HD2	3:H:48:VAL:HG12	2.03	0.41
3:H:6:GLU:OE1	3:H:6:GLU:N	2.49	0.41
3:H:63:VAL:HG13	3:H:67:PHE:CG	2.56	0.41
1:A:296:LYS:HD2	3:H:52(A):TRP:CZ2	2.56	0.40
3:H:40:PRO:HA	3:H:41:PRO:HD3	1.97	0.40
1:A:313:ASP:OD1	1:A:314:SER:N	2.54	0.40
3:H:60:THR:O	3:H:63:VAL:HB	2.20	0.40
2:L:145:LYS:HB3	2:L:197:THR:OG1	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	386/393 (98%)	364 (94%)	19 (5%)	3 (1%)	16 20

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	L	213/216 (99%)	195 (92%)	18 (8%)	0	100	100
3	H	223/240 (93%)	209 (94%)	13 (6%)	1 (0%)	30	37
All	All	822/849 (97%)	768 (93%)	50 (6%)	4 (0%)	24	32

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	200	ASN
1	A	347	PRO
1	A	222	ILE
3	H	151	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	338/342 (99%)	338 (100%)	0	100	100
2	L	189/190 (100%)	189 (100%)	0	100	100
3	H	193/205 (94%)	193 (100%)	0	100	100
All	All	720/737 (98%)	720 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	93	GLN
1	A	173	GLN
1	A	226	GLN
1	A	334	ASN
1	A	356	ASN
1	A	395	GLN
1	A	432	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	465	ASN
2	L	90	HIS
2	L	199	GLN
3	H	179	GLN
3	H	209	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

9 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	B	1	1,4	14,14,15	0.86	1 (7%)	17,19,21	0.91	1 (5%)
4	NAG	B	2	4	14,14,15	0.32	0	17,19,21	0.48	0
4	BMA	B	3	4	11,11,12	0.63	0	15,15,17	0.71	0
4	MAN	B	4	4	11,11,12	0.73	0	15,15,17	1.13	1 (6%)
4	MAN	B	5	4	11,11,12	0.90	0	15,15,17	1.53	2 (13%)
4	MAN	B	6	4	11,11,12	0.68	0	15,15,17	1.30	2 (13%)
4	MAN	B	7	4	11,11,12	0.62	0	15,15,17	0.99	2 (13%)
5	NAG	C	1	1,5	14,14,15	0.21	0	17,19,21	0.47	0
5	NAG	C	2	5	14,14,15	0.30	0	17,19,21	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	B	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	B	2	4	-	0/6/23/26	0/1/1/1
4	BMA	B	3	4	-	0/2/19/22	0/1/1/1
4	MAN	B	4	4	-	2/2/19/22	0/1/1/1
4	MAN	B	5	4	-	1/2/19/22	0/1/1/1
4	MAN	B	6	4	-	2/2/19/22	1/1/1/1
4	MAN	B	7	4	-	2/2/19/22	0/1/1/1
5	NAG	C	1	1,5	-	0/6/23/26	0/1/1/1
5	NAG	C	2	5	-	2/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	1	NAG	O5-C1	-2.20	1.40	1.43

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	5	MAN	C1-O5-C5	4.22	117.85	112.19
4	B	6	MAN	C1-O5-C5	3.98	117.52	112.19
4	B	5	MAN	O2-C2-C3	-3.39	103.13	110.15
4	B	4	MAN	C1-O5-C5	3.33	116.65	112.19
4	B	1	NAG	C1-O5-C5	2.75	115.87	112.19
4	B	7	MAN	C1-O5-C5	2.67	115.77	112.19
4	B	7	MAN	O2-C2-C3	-2.17	105.65	110.15
4	B	6	MAN	O2-C2-C3	-2.16	105.68	110.15

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	2	NAG	O5-C5-C6-O6
4	B	7	MAN	O5-C5-C6-O6
4	B	7	MAN	C4-C5-C6-O6
5	C	2	NAG	C4-C5-C6-O6
4	B	6	MAN	C4-C5-C6-O6
4	B	6	MAN	O5-C5-C6-O6
4	B	5	MAN	O5-C5-C6-O6
4	B	4	MAN	C4-C5-C6-O6

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	B	4	MAN	O5-C5-C6-O6


All (1) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	6	MAN	C1-C2-C3-C4-C5-O5

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	NAG	A	510	1	14,14,15	0.22	0	17,19,21	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	A	510	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	388/393 (98%)	0.30	7 (1%) 67 69	37, 56, 75, 99	0
2	L	215/216 (99%)	0.72	12 (5%) 30 27	54, 80, 106, 133	0
3	H	227/240 (94%)	0.80	20 (8%) 15 13	45, 81, 116, 126	0
All	All	830/849 (97%)	0.54	39 (4%) 36 35	37, 66, 108, 133	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	H	140	LEU	4.2
1	A	403	TRP	3.6
3	H	1	GLU	3.5
2	L	21	ILE	3.3
3	H	219	VAL	3.1
3	H	100(G)	ILE	3.1
2	L	19	VAL	3.1
1	A	93	GLN	3.0
3	H	205	THR	2.8
2	L	10	PHE	2.8
3	H	207	ILE	2.7
3	H	154	VAL	2.7
2	L	67	SER	2.6
3	H	165	ALA	2.6
2	L	214	CYS	2.6
3	H	110	THR	2.4
1	A	376	THR	2.4
2	L	2	ILE	2.4
3	H	105	ARG	2.4
3	H	227	PRO	2.4
3	H	83	ARG	2.3
2	L	66	GLY	2.3
2	L	95(A)	LEU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	H	225	VAL	2.3
2	L	154	LEU	2.2
3	H	120	SER	2.2
3	H	166	LEU	2.2
3	H	144	VAL	2.2
1	A	356	ASN	2.1
2	L	34	ALA	2.1
2	L	164	THR	2.1
3	H	164	GLY	2.1
1	A	339	ASP	2.1
2	L	127	SER	2.1
1	A	255	LEU	2.1
3	H	82(C)	LEU	2.1
1	A	424	VAL	2.1
3	H	228	LYS	2.1
3	H	143	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	B	1	14/15	-	-	47,59,64,65	0
4	NAG	B	2	14/15	-	-	52,60,71,72	0
4	BMA	B	3	11/12	-	-	66,71,83,102	0
4	MAN	B	4	11/12	-	-	70,78,81,85	0
4	MAN	B	5	11/12	-	-	90,96,102,107	0
4	MAN	B	6	11/12	-	-	103,110,115,117	0
4	MAN	B	7	11/12	-	-	111,122,126,129	0
5	NAG	C	1	14/15	-	-	74,86,96,97	0
5	NAG	C	2	14/15	-	-	98,106,112,116	0




The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain B:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain C:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	CA	A	512	1/1	0.72	0.32	109,109,109,109	1
6	NAG	A	510	14/15	0.78	0.11	106,109,115,118	0
7	CA	A	511	1/1	0.81	0.15	96,96,96,96	0

6.5 Other polymers [i](#)

There are no such residues in this entry.