



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:26 PM UTC

PDB ID : 2Q45 / pdb_00002q45
Title : Ensemble refinement of the protein crystal structure of putative tropinone reductase from Arabidopsis thaliana gene At1g07440
Authors : Levin, E.J.; Kondrashov, D.A.; Wesenberg, G.E.; Phillips Jr., G.N.; Center for Eukaryotic Structural Genomics (CESG)
Deposited on : 2007-05-31
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	1-A	266	36% 80% 18%
1	2-A	266	80% 18%
1	3-A	266	80% 18%
1	4-A	266	80% 18%
1	5-A	266	79% 18%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	6-A	266	 79% • 18%
1	7-A	266	 79% • 18%
1	8-A	266	 79% • 18%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 13512 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative tropinone reductase homolog At1g07440.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	1-A	218	Total	C	N	O	S	0	0	0
			1632	1030	278	313	11			
1	2-A	218	Total	C	N	O	S	0	0	0
			1632	1030	278	313	11			
1	3-A	218	Total	C	N	O	S	0	0	0
			1632	1030	278	313	11			
1	4-A	218	Total	C	N	O	S	0	0	0
			1632	1030	278	313	11			
1	5-A	218	Total	C	N	O	S	0	0	0
			1632	1030	278	313	11			
1	6-A	218	Total	C	N	O	S	0	0	0
			1632	1030	278	313	11			
1	7-A	218	Total	C	N	O	S	0	0	0
			1632	1030	278	313	11			
1	8-A	218	Total	C	N	O	S	0	0	0
			1632	1030	278	313	11			

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	1-A	57	Total	O	0	0
			57	57		
2	2-A	57	Total	O	0	0
			57	57		
2	3-A	57	Total	O	0	0
			57	57		
2	4-A	57	Total	O	0	0
			57	57		
2	5-A	57	Total	O	0	0
			57	57		
2	6-A	57	Total	O	0	0
			57	57		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	7-A	57	Total	O	0	0
			57	57		
2	8-A	57	Total	O	0	0
			57	57		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Putative tropinone reductase homolog At1g07440




- Molecule 1: Putative tropinone reductase homolog At1g07440




- Molecule 1: Putative tropinone reductase homolog At1g07440




- Molecule 1: Putative tropinone reductase homolog At1g07440






- Molecule 1: Putative tropinone reductase homolog At1g07440

Chain 5-A: 79% 18%




- Molecule 1: Putative tropinone reductase homolog At1g07440

Chain 6-A: 79% 18%




- Molecule 1: Putative tropinone reductase homolog At1g07440

Chain 7-A: 79% 18%



- Molecule 1: Putative tropinone reductase homolog At1g07440

Chain 8-A: 79% 18%



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	56.34Å 76.60Å 112.13Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.30 – 2.10 38.30 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.5 (38.30-2.10) 99.5 (38.30-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.50 (at 2.10Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.234 , 0.295 0.238 , 0.289	Depositor DCC
R_{free} test set	729 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	32.9	Xtriage
Anisotropy	0.297	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.04 , 37.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	13512	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.78% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	1-A	0.25	0/1660	0.50	0/2243
1	2-A	0.25	0/1660	0.50	0/2243
1	3-A	0.25	0/1660	0.50	0/2243
1	4-A	0.25	0/1660	0.50	0/2243
1	5-A	0.25	0/1660	0.50	0/2243
1	6-A	0.25	0/1660	0.50	0/2243
1	7-A	0.25	0/1660	0.50	0/2243
1	8-A	0.25	0/1660	0.50	0/2243
All	All	0.25	0/13280	0.50	0/17944

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	1-A	1632	0	1609	0	0
1	2-A	1632	0	1609	0	0
1	3-A	1632	0	1609	0	0
1	4-A	1632	0	1609	0	0
1	5-A	1632	0	1609	0	0
1	6-A	1632	0	1609	0	0
1	7-A	1632	0	1609	0	0
1	8-A	1632	0	1609	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	1-A	57	0	0	0	0
2	2-A	57	0	0	0	0
2	3-A	57	0	0	0	0
2	4-A	57	0	0	0	0
2	5-A	57	0	0	0	0
2	6-A	57	0	0	0	0
2	7-A	57	0	0	0	0
2	8-A	57	0	0	0	0
All	All	13512	0	12872	0	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

There are no clashes within the asymmetric unit.

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	1-A	210/266 (79%)	202 (96%)	6 (3%)	2 (1%)	12	9
1	2-A	210/266 (79%)	201 (96%)	7 (3%)	2 (1%)	12	9
1	3-A	210/266 (79%)	198 (94%)	11 (5%)	1 (0%)	24	22
1	4-A	210/266 (79%)	200 (95%)	7 (3%)	3 (1%)	9	5
1	5-A	210/266 (79%)	198 (94%)	9 (4%)	3 (1%)	9	5
1	6-A	210/266 (79%)	200 (95%)	7 (3%)	3 (1%)	9	5
1	7-A	210/266 (79%)	196 (93%)	9 (4%)	5 (2%)	4	2
1	8-A	210/266 (79%)	202 (96%)	5 (2%)	3 (1%)	9	5
All	All	1680/2128 (79%)	1597 (95%)	61 (4%)	22 (1%)	9	6

All (22) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	1-A	72	ALA
1	4-A	162	SER
1	7-A	72	ALA
1	7-A	162	SER
1	7-A	195	ALA
1	8-A	162	SER
1	1-A	162	SER
1	4-A	149	MET
1	5-A	72	ALA
1	5-A	143	CYS
1	6-A	162	SER
1	7-A	194	PRO
1	8-A	72	ALA
1	2-A	72	ALA
1	5-A	162	SER
1	7-A	21	GLY
1	6-A	251	VAL
1	2-A	251	VAL
1	3-A	251	VAL
1	8-A	251	VAL
1	4-A	251	VAL
1	6-A	254	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	1-A	173/211 (82%)	169 (98%)	4 (2%)	44	51
1	2-A	173/211 (82%)	169 (98%)	4 (2%)	44	51
1	3-A	173/211 (82%)	169 (98%)	4 (2%)	44	51
1	4-A	173/211 (82%)	169 (98%)	4 (2%)	44	51
1	5-A	173/211 (82%)	169 (98%)	4 (2%)	44	51
1	6-A	173/211 (82%)	169 (98%)	4 (2%)	44	51
1	7-A	173/211 (82%)	169 (98%)	4 (2%)	44	51
1	8-A	173/211 (82%)	169 (98%)	4 (2%)	44	51

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	1384/1688 (82%)	1352 (98%)	32 (2%)	44 51

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	1-A	24	LYS
1	1-A	145	ASN
1	1-A	149	MET
1	1-A	227	GLU
1	2-A	24	LYS
1	2-A	145	ASN
1	2-A	149	MET
1	2-A	227	GLU
1	3-A	24	LYS
1	3-A	145	ASN
1	3-A	149	MET
1	3-A	227	GLU
1	4-A	24	LYS
1	4-A	145	ASN
1	4-A	149	MET
1	4-A	227	GLU
1	5-A	24	LYS
1	5-A	145	ASN
1	5-A	149	MET
1	5-A	227	GLU
1	6-A	24	LYS
1	6-A	145	ASN
1	6-A	149	MET
1	6-A	227	GLU
1	7-A	24	LYS
1	7-A	145	ASN
1	7-A	149	MET
1	7-A	227	GLU
1	8-A	24	LYS
1	8-A	145	ASN
1	8-A	149	MET
1	8-A	227	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (63) such sidechains are listed below:

Mol	Chain	Res	Type
1	1-A	64	GLN
1	1-A	83	GLN
1	1-A	123	ASN
1	1-A	173	GLN
1	1-A	177	ASN
1	1-A	258	ASN
1	2-A	52	ASN
1	2-A	83	GLN
1	2-A	123	ASN
1	2-A	135	HIS
1	2-A	145	ASN
1	2-A	173	GLN
1	2-A	177	ASN
1	2-A	258	ASN
1	3-A	52	ASN
1	3-A	64	GLN
1	3-A	83	GLN
1	3-A	123	ASN
1	3-A	173	GLN
1	3-A	177	ASN
1	3-A	258	ASN
1	4-A	64	GLN
1	4-A	83	GLN
1	4-A	119	HIS
1	4-A	123	ASN
1	4-A	132	GLN
1	4-A	145	ASN
1	4-A	173	GLN
1	4-A	258	ASN
1	5-A	52	ASN
1	5-A	59	GLN
1	5-A	64	GLN
1	5-A	83	GLN
1	5-A	123	ASN
1	5-A	135	HIS
1	5-A	173	GLN
1	5-A	258	ASN
1	6-A	64	GLN
1	6-A	83	GLN
1	6-A	132	GLN
1	6-A	145	ASN
1	6-A	173	GLN
1	6-A	258	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	7-A	52	ASN
1	7-A	59	GLN
1	7-A	64	GLN
1	7-A	83	GLN
1	7-A	98	ASN
1	7-A	123	ASN
1	7-A	132	GLN
1	7-A	135	HIS
1	7-A	145	ASN
1	7-A	173	GLN
1	7-A	258	ASN
1	8-A	8	GLN
1	8-A	47	ASN
1	8-A	52	ASN
1	8-A	64	GLN
1	8-A	123	ASN
1	8-A	145	ASN
1	8-A	173	GLN
1	8-A	177	ASN
1	8-A	258	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	1-A	218/266 (81%)	2.05	97 (44%) 0 1	2, 4, 6, 7	218 (100%)
1	2-A	0/266	-	-	-	-
1	3-A	0/266	-	-	-	-
1	4-A	0/266	-	-	-	-
1	5-A	0/266	-	-	-	-
1	6-A	0/266	-	-	-	-
1	7-A	0/266	-	-	-	-
1	8-A	0/266	-	-	-	-
All	All	218/2128 (10%)	2.05	97 (44%) 0 1	2, 4, 6, 7	218 (100%)

All (97) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	1-A	180	CYS	6.6
1	1-A	150	SER	6.4
1	1-A	121	SER	6.4
1	1-A	109	LEU	6.4
1	1-A	255	LEU	6.2
1	1-A	49	TYR	6.1
1	1-A	257	VAL	6.0
1	1-A	261	SER	5.8
1	1-A	265	GLN	5.4
1	1-A	110	ASP	5.3
1	1-A	258	ASN	5.3
1	1-A	7	SER	5.0
1	1-A	102	ALA	5.0
1	1-A	115	ASP	4.9
1	1-A	119	HIS	4.8
1	1-A	165	SER	4.7
1	1-A	51	LEU	4.5
1	1-A	120	ILE	4.5
1	1-A	164	TYR	4.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	1-A	122	THR	4.3
1	1-A	39	ALA	4.2
1	1-A	88	MET	4.1
1	1-A	256	THR	4.1
1	1-A	55	LEU	4.0
1	1-A	264	PRO	4.0
1	1-A	136	PRO	3.9
1	1-A	38	GLY	3.8
1	1-A	197	ILE	3.7
1	1-A	83	GLN	3.7
1	1-A	259	GLY	3.6
1	1-A	198	ALA	3.6
1	1-A	184	SER	3.6
1	1-A	111	TYR	3.4
1	1-A	177	ASN	3.4
1	1-A	228	GLU	3.4
1	1-A	70	CYS	3.4
1	1-A	59	GLN	3.3
1	1-A	149	MET	3.3
1	1-A	125	GLU	3.2
1	1-A	187	ILE	3.2
1	1-A	134	ALA	3.1
1	1-A	9	ARG	3.1
1	1-A	50	GLU	3.1
1	1-A	227	GLU	3.1
1	1-A	60	LYS	3.1
1	1-A	142	GLY	3.1
1	1-A	161	GLY	3.1
1	1-A	167	THR	3.1
1	1-A	224	GLY	3.1
1	1-A	53	GLU	3.1
1	1-A	253	GLY	3.1
1	1-A	86	SER	3.0
1	1-A	54	CYS	2.9
1	1-A	87	SER	2.9
1	1-A	178	LEU	2.9
1	1-A	135	HIS	2.8
1	1-A	143	CYS	2.7
1	1-A	225	GLU	2.7
1	1-A	260	PHE	2.7
1	1-A	185	ASP	2.7
1	1-A	100	LEU	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	1-A	137	LEU	2.6
1	1-A	71	ASP	2.6
1	1-A	90	GLY	2.6
1	1-A	75	ARG	2.6
1	1-A	65	VAL	2.6
1	1-A	99	ASN	2.6
1	1-A	14	ALA	2.6
1	1-A	181	GLU	2.5
1	1-A	101	GLY	2.5
1	1-A	123	ASN	2.5
1	1-A	19	VAL	2.5
1	1-A	36	GLY	2.5
1	1-A	196	VAL	2.4
1	1-A	132	GLN	2.4
1	1-A	73	SER	2.4
1	1-A	162	SER	2.4
1	1-A	68	SER	2.4
1	1-A	24	LYS	2.4
1	1-A	170	ALA	2.4
1	1-A	179	ALA	2.4
1	1-A	173	GLN	2.3
1	1-A	147	ILE	2.3
1	1-A	176	ARG	2.3
1	1-A	74	LEU	2.3
1	1-A	263	GLN	2.2
1	1-A	118	PHE	2.2
1	1-A	52	ASN	2.2
1	1-A	226	PRO	2.2
1	1-A	254	GLY	2.2
1	1-A	126	SER	2.2
1	1-A	58	TRP	2.2
1	1-A	128	TYR	2.1
1	1-A	163	ILE	2.1
1	1-A	124	LEU	2.1
1	1-A	57	LYS	2.0
1	1-A	262	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.