



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 4, 2026 – 08:15 AM EDT

PDB ID : 2QBO / pdb_00002qbo
Title : Crystal structure of the P450cam G248V mutant in the cyanide bound state
Authors : von Koenig, K.; Makris, T.M.; Sligar, S.D.; Schlichting, I.
Deposited on : 2007-06-18
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.015 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	421	

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 3802 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Cytochrome P450-cam.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	405	3225	2045	562	599	19	0	2	0

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	248	VAL	GLY	engineered mutation	UNP P00183
A	415	HIS	-	expression tag	UNP P00183
A	416	HIS	-	expression tag	UNP P00183
A	417	HIS	-	expression tag	UNP P00183
A	418	HIS	-	expression tag	UNP P00183
A	419	HIS	-	expression tag	UNP P00183
A	420	HIS	-	expression tag	UNP P00183

- Molecule 2 is CYANIDE ION (CCD ID: CYN) (formula: CN).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	N	0	0
			2	1	1		

- Molecule 3 is POTASSIUM ION (CCD ID: K) (formula: K).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total K 1 1	0	0

- Molecule 4 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 5 is CAMPHOR (CCD ID: CAM) (formula: $\text{C}_{10}\text{H}_{16}\text{O}$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			11	10	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	520	Total	O	0	0
			520	520		

4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	63.68Å 63.68Å 241.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.90 20.00 – 1.90	Depositor EDS
% Data completeness (in resolution range)	98.8 (20.00-1.90) 98.8 (20.00-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.29 (at 1.90Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.201 , 0.252 0.197 , 0.245	Depositor DCC
R_{free} test set	1995 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	30.1	Xtriage
Anisotropy	0.305	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 35.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3802	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CAM, K, HEM, CYN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.41	0/3304	0.89	9/4490 (0.2%)

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	105	PRO	N-CA-C	-7.74	101.25	110.70
1	A	18	VAL	N-CA-C	6.17	114.68	107.77
1	A	109	ARG	N-CA-C	5.72	117.97	111.11
1	A	324	LEU	N-CA-C	5.70	118.43	111.82
1	A	251	ASP	N-CA-C	5.42	119.39	112.12
1	A	48	SER	N-CA-C	5.21	117.63	111.33
1	A	150	PHE	N-CA-C	5.07	117.47	111.33
1	A	85	CYS	CA-C-N	5.03	125.85	120.47
1	A	85	CYS	C-N-CA	5.03	125.85	120.47

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3225	0	3176	76	0
2	A	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	1	0	0	0	0
4	A	43	0	30	2	0
5	A	11	0	16	1	0
6	A	520	0	0	24	0
All	All	3802	0	3222	77	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:69:GLN:HG2	6:A:728:HOH:O	1.72	0.89
1:A:49:ASN:HD22	1:A:49:ASN:H	1.23	0.83
1:A:158:PHE:HB3	1:A:159:PRO:HD3	1.59	0.82
1:A:380:ASP:HB2	6:A:734:HOH:O	1.79	0.82
1:A:181:THR:HA	1:A:184:MET:HE3	1.64	0.78
1:A:306:GLU:HB3	6:A:933:HOH:O	1.89	0.72
1:A:49:ASN:HD22	1:A:49:ASN:N	1.85	0.72
1:A:198:GLU:HG2	6:A:741:HOH:O	1.89	0.72
1:A:85:CYS:HB2	6:A:940:HOH:O	1.88	0.72
1:A:223:VAL:HG22	1:A:241[B]:MET:HE2	1.74	0.70
1:A:277:ARG:HD3	1:A:280:ARG:CZ	2.24	0.67
1:A:49:ASN:H	1:A:49:ASN:ND2	1.91	0.67
1:A:28:MET:HE1	1:A:395:ILE:HD13	1.77	0.66
1:A:327:LEU:HD22	1:A:336:MET:HE1	1.76	0.66
1:A:209:GLU:O	1:A:213:GLN:HG2	1.96	0.66
1:A:241[B]:MET:HG3	6:A:745:HOH:O	2.00	0.62
1:A:294:LEU:HD23	1:A:294:LEU:H	1.64	0.61
1:A:128:GLU:HG3	1:A:365:ARG:NH2	2.16	0.60
1:A:281:ILE:HB	1:A:282:PRO:HD3	1.82	0.60
1:A:343:GLN:HG3	6:A:899:HOH:O	2.01	0.59
1:A:223:VAL:CG2	1:A:241[B]:MET:HE2	2.36	0.56
1:A:111:PHE:HB3	6:A:880:HOH:O	2.04	0.56
1:A:281:ILE:HD11	6:A:936:HOH:O	2.05	0.56
1:A:114:LEU:HG	1:A:241[B]:MET:HE1	1.88	0.56
1:A:414:VAL:HA	6:A:764:HOH:O	2.05	0.55
1:A:68:GLY:HA3	1:A:331:GLU:OE2	2.06	0.55
1:A:69:GLN:NE2	6:A:733:HOH:O	2.39	0.55
1:A:294:LEU:HD23	1:A:294:LEU:N	2.23	0.53
1:A:303:SER:HA	1:A:314:LYS:HB2	1.91	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:192:THR:OG1	1:A:195:GLU:HG3	2.08	0.53
1:A:382:SER:HB2	6:A:563:HOH:O	2.08	0.52
1:A:223:VAL:HG22	1:A:241[B]:MET:CE	2.38	0.52
1:A:128:GLU:CG	1:A:365:ARG:HH22	2.22	0.51
1:A:400:GLN:HG3	6:A:730:HOH:O	2.09	0.51
1:A:344:LYS:HB2	6:A:949:HOH:O	2.11	0.51
1:A:356:LEU:HD22	1:A:360:GLN:NE2	2.26	0.50
1:A:223:VAL:HG13	1:A:241[B]:MET:HE2	1.92	0.50
1:A:386:GLY:HA2	6:A:722:HOH:O	2.11	0.50
1:A:158:PHE:CB	1:A:159:PRO:HD3	2.38	0.49
1:A:212:ARG:HG3	1:A:224:ALA:HB1	1.95	0.49
1:A:79:ARG:HD3	6:A:842:HOH:O	2.13	0.49
1:A:390:GLN:HB3	6:A:766:HOH:O	2.14	0.48
1:A:151:THR:O	1:A:156:GLU:HG3	2.15	0.47
1:A:390:GLN:HG3	6:A:856:HOH:O	2.14	0.47
1:A:141:SER:HB3	6:A:623:HOH:O	2.13	0.47
4:A:441:HEM:HBA1	5:A:442:CAM:H4	1.97	0.47
1:A:356:LEU:HD22	1:A:360:GLN:HE22	1.81	0.47
1:A:412:LYS:HE2	6:A:762:HOH:O	2.15	0.46
1:A:413:ALA:O	1:A:414:VAL:HB	2.14	0.46
1:A:233:ILE:HD11	1:A:241[A]:MET:CE	2.46	0.46
1:A:99:ILE:HG21	1:A:241[A]:MET:HE2	1.97	0.45
1:A:345:VAL:HG21	6:A:528:HOH:O	2.15	0.44
1:A:39:GLN:NE2	1:A:39:GLN:H	2.15	0.44
1:A:184:MET:HE1	1:A:200:LEU:HD22	2.00	0.44
1:A:245:LEU:HA	1:A:248:VAL:HG22	1.99	0.44
1:A:294:LEU:N	1:A:294:LEU:CD2	2.81	0.43
1:A:327:LEU:HD22	1:A:336:MET:CE	2.47	0.43
1:A:99:ILE:HA	1:A:100:PRO:HA	1.82	0.43
1:A:28:MET:CE	1:A:395:ILE:HD13	2.47	0.43
1:A:121:MET:N	1:A:122:PRO:CD	2.81	0.43
1:A:151:THR:HA	1:A:155:ALA:HB3	2.01	0.43
1:A:170:PRO:HG2	1:A:173:ASP:OD1	2.19	0.43
1:A:208:ILE:CD1	1:A:239:LYS:HA	2.49	0.42
1:A:252:THR:HB	4:A:441:HEM:CAB	2.50	0.42
1:A:146:GLY:HA2	1:A:406:TRP:NE1	2.34	0.42
1:A:179:TYR:O	1:A:183:GLN:HG2	2.19	0.42
1:A:223:VAL:CG1	1:A:241[B]:MET:HE2	2.50	0.42
1:A:132:GLN:HE22	1:A:373:GLU:CD	2.28	0.41
1:A:88:ILE:HD11	1:A:317:GLN:HB3	2.01	0.41
1:A:372:LYS:HG3	6:A:935:HOH:O	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:128:GLU:CB	1:A:365:ARG:HH22	2.33	0.41
1:A:383:ILE:HG22	1:A:387:ALA:HB3	2.02	0.41
1:A:143:ARG:CZ	1:A:413:ALA:HB2	2.50	0.41
1:A:274:LEU:HB3	6:A:936:HOH:O	2.19	0.41
1:A:181:THR:HG22	1:A:247:VAL:HG22	2.01	0.41
1:A:372:LYS:HD2	6:A:815:HOH:O	2.20	0.40
1:A:201:TYR:O	1:A:205:ILE:HG13	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	405/421 (96%)	392 (97%)	12 (3%)	1 (0%)	43 36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	321	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	353/366 (96%)	346 (98%)	7 (2%)	48 46

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	39	GLN
1	A	49	ASN
1	A	108	GLN
1	A	228	VAL
1	A	252	THR
1	A	344	LYS
1	A	356	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	30	ASN
1	A	39	GLN
1	A	46	GLN
1	A	49	ASN
1	A	108	GLN
1	A	117	GLN
1	A	129	ASN
1	A	132	GLN
1	A	210	GLN
1	A	225	ASN
1	A	229	ASN
1	A	317	GLN
1	A	343	GLN
1	A	361	HIS
1	A	388	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	HEM	A	441	1	50,50,50	1.34	5 (10%)	67,82,82	0.68	0
2	CYN	A	430	-	1,1,1	0.18	0	-		
5	CAM	A	442	-	12,12,12	1.56	2 (16%)	20,21,21	1.35	2 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	HEM	A	441	1	-	3/14/54/54	-
5	CAM	A	442	-	-	-	0/3/2/2

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	441	HEM	FE-NC	3.76	2.07	1.95
5	A	442	CAM	C1-C2	3.30	1.57	1.52
4	A	441	HEM	CAC-C3C	-3.06	1.39	1.47
4	A	441	HEM	CAB-C3B	-3.03	1.39	1.47
4	A	441	HEM	FE-NA	2.86	2.04	1.95
4	A	441	HEM	FE-ND	2.74	2.03	1.94
5	A	442	CAM	C3-C4	2.29	1.59	1.53

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	442	CAM	C7-C1-C2	3.77	106.75	100.33
5	A	442	CAM	O-C2-C1	2.59	128.44	125.33

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	441	HEM	C2B-C3B-CAB-CBB
4	A	441	HEM	C2C-C3C-CAC-CBC
4	A	441	HEM	C4C-C3C-CAC-CBC

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	441	HEM	2	0
5	A	442	CAM	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	405/421 (96%)	0.14	7 (1%) 69 72	14, 31, 40, 44	2 (0%)

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	414	VAL	4.0
1	A	341	SER	2.9
1	A	413	ALA	2.6
1	A	386	GLY	2.4
1	A	306	GLU	2.3
1	A	380	ASP	2.3
1	A	388	GLN	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	CAM	A	442	11/11	0.89	0.09	32,35,37,38	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CYN	A	430	2/2	0.95	0.17	31,31,31,34	0
4	HEM	A	441	43/43	0.98	0.05	22,25,29,29	0
3	K	A	440	1/1	1.00	0.03	28,28,28,28	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.