



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 01:24 PM UTC

PDB ID : 1QPJ / pdb_00001qpj
Title : CRYSTAL STRUCTURE OF THE LYMPHOCYTE-SPECIFIC KINASE
LCK IN COMPLEX WITH STAUROSPORINE.
Authors : Zhu, X.; Kim, J.L.; Rose, P.E.; Stover, D.R.; Toledo, L.M.
Deposited on : 1999-05-25
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	279	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2963 atoms, of which 675 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called LCK TYROSINE KINASE.

Mol	Chain	Residues	Atoms								ZeroOcc	AltConf	Trace
			Total	C	H	N	O	P	S				
1	A	266	2591	1368	453	357	400	1	12		453	0	0

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	394	PTR	TYR	modified residue	UNP P06239

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0

- Molecule 3 is STAUROSPORINE (CCD ID: STU) (formula: C₂₈H₂₆N₄O₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	H	N	O	2	0
			37	28	2	4	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	110	Total	H	O	220	0
			330	220	110		

4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	61.54Å 69.04Å 73.73Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	6.00 – 2.20	Depositor
% Data completeness (in resolution range)	(Not available) (6.00-2.20)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 98.0	Depositor
R, R_{free}	0.214 , 0.267	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2963	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: STU, SO4, PTR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.54	32/2169 (1.5%)	1.18	17/2941 (0.6%)

All (32) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	483	ASP	CA-CB	8.66	1.68	1.53
1	A	405	LYS	CA-CB	7.53	1.65	1.53
1	A	291	LEU	CA-CB	7.50	1.65	1.53
1	A	342	LEU	C-O	-7.26	1.15	1.24
1	A	465	CYS	C-O	-7.14	1.15	1.24
1	A	261	MET	CA-CB	-6.81	1.42	1.53
1	A	371	LEU	C-O	-6.78	1.15	1.23
1	A	327	PHE	CA-CB	6.49	1.63	1.53
1	A	478	LYS	CA-CB	-6.40	1.42	1.53
1	A	270	VAL	C-O	-6.29	1.16	1.23
1	A	432	GLU	C-O	-6.29	1.16	1.24
1	A	321	ASN	CA-CB	6.18	1.63	1.53
1	A	390	GLU	CA-CB	-5.95	1.44	1.53
1	A	392	ASN	CA-CB	5.93	1.63	1.53
1	A	402	PHE	CA-CB	5.91	1.62	1.54
1	A	252	GLY	C-O	-5.79	1.18	1.23
1	A	466	PRO	CA-CB	5.75	1.61	1.53
1	A	265	ASN	CA-CB	5.72	1.62	1.53
1	A	376	LEU	CB-CG	5.67	1.64	1.53
1	A	427	GLY	C-O	-5.63	1.17	1.23
1	A	350	GLU	CA-CB	5.47	1.61	1.53
1	A	247	LEU	CA-CB	-5.42	1.44	1.53
1	A	385	LEU	CA-CB	5.38	1.63	1.53
1	A	305	ALA	C-O	-5.34	1.17	1.23
1	A	352	MET	C-O	-5.31	1.17	1.24
1	A	476	CYS	CA-CB	5.29	1.62	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	451	GLN	CA-CB	5.29	1.61	1.53
1	A	255	GLN	CA-CB	5.28	1.61	1.53
1	A	243	GLU	C-O	-5.27	1.17	1.24
1	A	251	LEU	CA-CB	5.09	1.61	1.53
1	A	472	LEU	CB-CG	5.08	1.63	1.53
1	A	452	ASN	CG-OD1	-5.07	1.14	1.23

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	392	ASN	N-CA-C	8.02	127.88	110.80
1	A	320	GLU	N-CA-C	7.32	120.12	111.71
1	A	391	ASP	N-CA-C	-7.25	97.86	109.76
1	A	404	ILE	N-CA-C	7.18	117.90	110.36
1	A	402	PHE	CA-C-N	6.91	128.48	119.84
1	A	402	PHE	C-N-CA	6.91	128.48	119.84
1	A	398	GLU	N-CA-C	6.83	119.57	111.71
1	A	267	HIS	N-CA-C	6.12	121.03	113.50
1	A	408	ALA	CA-C-N	5.52	124.98	119.24
1	A	408	ALA	C-N-CA	5.52	124.98	119.24
1	A	302	ARG	N-CA-C	5.39	118.33	109.76
1	A	256	PHE	N-CA-C	5.27	119.41	112.92
1	A	271	ALA	N-CA-C	-5.12	101.35	109.59
1	A	434	VAL	N-CA-C	5.12	117.45	111.05
1	A	441	TYR	CA-C-N	5.11	125.39	119.93
1	A	441	TYR	C-N-CA	5.11	125.39	119.93
1	A	240	VAL	N-CA-C	-5.08	105.02	109.19

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2138	453	2101	31	0
2	A	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	35	2	26	2	0
4	A	110	220	0	0	0
All	All	2288	675	2127	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:392:ASN:HD22	1:A:392:ASN:H	1.34	0.75
1:A:392:ASN:H	1:A:392:ASN:ND2	1.82	0.74
3:A:902:STU:H261	3:A:902:STU:H16	1.68	0.74
1:A:256:PHE:CD1	1:A:280:MET:HG3	2.26	0.70
1:A:358:ARG:HB3	1:A:358:ARG:NH2	2.11	0.66
1:A:294:GLN:O	1:A:358:ARG:NH1	2.32	0.63
1:A:242:ARG:NH1	1:A:311:PRO:O	2.34	0.60
1:A:438:ARG:CZ	1:A:442:PRO:HB3	2.32	0.59
1:A:282:PRO:HB3	1:A:312:ILE:HD11	1.85	0.59
1:A:358:ARG:HB3	1:A:358:ARG:HH21	1.70	0.57
1:A:261:MET:HG3	1:A:318:TYR:CE1	2.42	0.54
1:A:280:MET:HE2	1:A:284:ALA:HB1	1.89	0.54
1:A:354:PHE:HA	1:A:357:GLU:HG2	1.90	0.54
1:A:419:ILE:O	1:A:423:VAL:HG23	2.08	0.53
1:A:285:PHE:CE2	1:A:314:ILE:HG13	2.45	0.52
1:A:315:ILE:HD12	1:A:315:ILE:N	2.24	0.52
1:A:407:THR:HG22	1:A:411:ALA:HB3	1.92	0.52
1:A:331:PRO:O	1:A:335:LYS:HG3	2.10	0.51
1:A:301:VAL:HG21	1:A:381:ALA:HB2	1.90	0.51
1:A:325:VAL:HG12	1:A:368:ALA:HA	1.96	0.47
1:A:307:VAL:HB	1:A:313:TYR:HB2	1.97	0.47
1:A:330:THR:HB	1:A:331:PRO:CD	2.45	0.47
1:A:402:PHE:HA	1:A:403:PRO:HD3	1.81	0.45
1:A:434:VAL:HG11	1:A:469:LEU:HD23	1.98	0.45
1:A:392:ASN:ND2	1:A:392:ASN:N	2.58	0.44
1:A:451:GLN:O	1:A:455:ARG:HG3	2.18	0.43
1:A:366:ARG:HB3	1:A:406:TRP:CD1	2.53	0.43
1:A:253:ALA:HA	1:A:258:GLU:HA	2.02	0.42
1:A:446:ASN:HB2	1:A:447:PRO:HD3	2.02	0.42
3:A:902:STU:H261	3:A:902:STU:C16	2.43	0.42
1:A:423:VAL:HG13	1:A:490:LEU:HD11	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:243:GLU:H	1:A:243:GLU:CD	2.29	0.41
1:A:438:ARG:NH2	1:A:442:PRO:HB3	2.35	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	263/279 (94%)	254 (97%)	8 (3%)	1 (0%)	30 34

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	392	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	226/242 (93%)	220 (97%)	6 (3%)	39 53

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	249	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	282	PRO
1	A	294	GLN
1	A	392	ASN
1	A	398	GLU
1	A	479	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	255	GLN
1	A	265	ASN
1	A	294	GLN
1	A	392	ASN
1	A	451	GLN
1	A	452	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	PTR	A	394	1	15,16,17	2.61	3 (20%)	17,22,24	1.02	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PTR	A	394	1	-	1/10/11/13	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	394	PTR	P-OH	8.94	1.76	1.59
1	A	394	PTR	CE2-CD2	-2.73	1.34	1.38
1	A	394	PTR	CB-CA	-2.20	1.49	1.53

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	394	PTR	OH-P-O1P	-2.02	102.73	109.48

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	394	PTR	CE1-CZ-OH-P

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	STU	A	902	-	39,42,42	2.53	17 (43%)	50,68,68	2.91	16 (32%)
2	SO4	A	901	-	4,4,4	0.88	0	6,6,6	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	STU	A	902	-	-	0/4/42/42	-

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	902	STU	C18-N2	6.04	1.46	1.40
3	A	902	STU	C8-N1	-5.17	1.31	1.35
3	A	902	STU	C24-C23	4.76	1.60	1.53
3	A	902	STU	C23-N4	4.46	1.52	1.47
3	A	902	STU	C14-C13	4.06	1.45	1.38
3	A	902	STU	C9-C10	-3.81	1.47	1.50
3	A	902	STU	C6-C19	3.54	1.47	1.41
3	A	902	STU	C11-C18	2.97	1.47	1.41
3	A	902	STU	C5-C20	-2.91	1.36	1.41
3	A	902	STU	C13-C12	2.80	1.44	1.40
3	A	902	STU	C19-C18	2.76	1.46	1.40
3	A	902	STU	C12-C11	-2.72	1.40	1.46
3	A	902	STU	C3-C2	2.60	1.44	1.38
3	A	902	STU	C6-C7	-2.51	1.36	1.40
3	A	902	STU	C19-N3	-2.15	1.35	1.39
3	A	902	STU	C22-C23	2.07	1.55	1.53
3	A	902	STU	C9-N1	2.04	1.47	1.45

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	902	STU	C11-C12-C17	10.44	117.01	106.49
3	A	902	STU	C9-C10-C7	-6.76	107.40	109.88
3	A	902	STU	C6-C5-C20	6.74	113.28	106.49
3	A	902	STU	C12-C11-C18	-5.94	98.10	106.44
3	A	902	STU	C12-C11-C10	5.72	142.73	132.33
3	A	902	STU	C12-C17-N2	-4.77	105.81	108.94
3	A	902	STU	C5-C6-C19	-4.71	99.83	106.44

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	902	STU	C3-C2-C1	-3.82	115.53	120.24
3	A	902	STU	C11-C18-N2	3.77	113.47	109.38
3	A	902	STU	C13-C12-C11	-3.38	125.45	133.49
3	A	902	STU	C11-C10-C7	2.80	123.09	121.25
3	A	902	STU	C19-C18-N2	-2.59	129.18	131.38
3	A	902	STU	C4-C5-C6	-2.45	127.66	133.49
3	A	902	STU	C5-C6-C7	2.26	142.02	134.87
3	A	902	STU	C6-C7-C10	2.23	122.32	120.76
3	A	902	STU	C2-C1-C20	2.13	122.58	118.28

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	902	STU	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.