



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 01:33 PM UTC

PDB ID : 3QQW / pdb_00003qqw
Title : Crystal structure of a putative lyase (Reut_B4148) from *Ralstonia eutropha* JMP134 at 2.44 Å resolution
Authors : Joint Center for Structural Genomics (JCSG)
Deposited on : 2011-02-16
Resolution : 2.44 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.44 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2340 (2.46-2.42)
Clashscore	190562	2400 (2.46-2.42)
Ramachandran outliers	187476	2379 (2.46-2.42)
Sidechain outliers	187428	2379 (2.46-2.42)
RSRZ outliers	180081	2340 (2.46-2.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	332	
1	B	332	
1	C	332	
1	D	332	
1	E	332	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	332	6% 78% 7% 15%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 13240 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative citrate lyase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	284	Total	C	N	O	S	Se	0	4	0
			2119	1334	384	386	5	10			
1	B	294	Total	C	N	O	S	Se	0	3	0
			2181	1376	389	401	5	10			
1	C	298	Total	C	N	O	S	Se	0	3	0
			2209	1387	396	412	5	9			
1	D	281	Total	C	N	O	S	Se	0	1	0
			2071	1302	372	382	5	10			
1	E	299	Total	C	N	O	S	Se	0	4	0
			2259	1427	403	414	5	10			
1	F	283	Total	C	N	O	S	Se	0	0	0
			2097	1321	378	384	5	9			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP Q46TN2
B	0	GLY	-	expression tag	UNP Q46TN2
C	0	GLY	-	expression tag	UNP Q46TN2
D	0	GLY	-	expression tag	UNP Q46TN2
E	0	GLY	-	expression tag	UNP Q46TN2
F	0	GLY	-	expression tag	UNP Q46TN2

- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cl	0	0
			1	1		
2	B	2	Total	Cl	0	0
			2	2		
2	C	2	Total	Cl	0	0
			2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	D	1	Total 1	Cl 1	0	0
2	E	2	Total 2	Cl 2	0	0
2	F	1	Total 1	Cl 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	41	Total 41	O 41	0	0
3	B	34	Total 34	O 34	0	0
3	C	60	Total 60	O 60	0	0
3	D	41	Total 41	O 41	0	1
3	E	81	Total 82	O 82	0	2
3	F	37	Total 37	O 37	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Putative citrate lyase




- Molecule 1: Putative citrate lyase





- Molecule 1: Putative citrate lyase





- Molecule 1: Putative citrate lyase



- Molecule 1: Putative citrate lyase



- Molecule 1: Putative citrate lyase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	226.40Å 73.83Å 149.25Å 90.00° 125.31° 90.00°	Depositor
Resolution (Å)	29.79 – 2.44 29.79 – 2.44	Depositor EDS
% Data completeness (in resolution range)	(Not available) (29.79-2.44) 99.4 (29.79-2.44)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.87 (at 2.45Å)	Xtriage
Refinement program	BUSTER	Depositor
R, R_{free}	0.202 , 0.231 0.212 , 0.241	Depositor DCC
R_{free} test set	3766 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	44.0	Xtriage
Anisotropy	0.238	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 52.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	13240	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 33.21 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.2994e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.83	0/2167	1.31	3/2933 (0.1%)
1	B	0.80	0/2228	1.31	5/3018 (0.2%)
1	C	0.87	0/2255	1.29	3/3056 (0.1%)
1	D	0.85	0/2109	1.31	6/2858 (0.2%)
1	E	0.91	1/2311 (0.0%)	1.32	9/3126 (0.3%)
1	F	0.83	0/2133	1.30	6/2888 (0.2%)
All	All	0.85	1/13203 (0.0%)	1.31	32/17879 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	54	CYS	CA-C	-5.60	1.47	1.52

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	269	ASP	CA-CB-CG	-6.66	105.94	112.60
1	C	99	ASP	CA-CB-CG	5.81	118.41	112.60
1	D	271	THR	CA-C-N	5.78	128.02	120.28
1	D	271	THR	C-N-CA	5.78	128.02	120.28
1	D	99	ASP	CA-CB-CG	5.75	118.35	112.60
1	F	42	VAL	CA-C-N	5.69	128.17	120.65
1	F	42	VAL	C-N-CA	5.69	128.17	120.65
1	B	137	LYS	CA-C-N	5.62	128.38	120.28
1	B	137	LYS	C-N-CA	5.62	128.38	120.28
1	F	231	ASP	CA-CB-CG	5.61	118.21	112.60
1	F	99	ASP	CA-CB-CG	5.54	118.14	112.60
1	E	27	PHE	CA-CB-CG	-5.53	108.27	113.80
1	C	121	ASN	CA-C-N	5.52	127.67	120.28
1	C	121	ASN	C-N-CA	5.52	127.67	120.28

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	99	ASP	CA-CB-CG	5.47	118.07	112.60
1	E	99	ASP	CA-CB-CG	5.47	118.07	112.60
1	E	54	CYS	N-CA-C	-5.45	105.68	112.93
1	D	260	GLN	OE1-CD-NE2	-5.39	117.21	122.60
1	E	117	PRO	CA-C-N	5.39	130.01	122.08
1	E	117	PRO	C-N-CA	5.39	130.01	122.08
1	E	56	ASP	CA-C-N	5.27	125.79	119.94
1	E	56	ASP	C-N-CA	5.27	125.79	119.94
1	D	260	GLN	CG-CD-NE2	5.27	124.30	116.40
1	A	99	ASP	CA-CB-CG	5.24	117.84	112.60
1	B	54	CYS	CA-C-N	5.20	127.24	120.28
1	B	54	CYS	C-N-CA	5.20	127.24	120.28
1	E	121	ASN	CA-C-N	5.16	127.19	120.28
1	E	121	ASN	C-N-CA	5.16	127.19	120.28
1	F	97	ARG	CA-C-N	5.10	127.11	120.28
1	F	97	ARG	C-N-CA	5.10	127.11	120.28
1	A	77	ASP	CA-C-N	5.04	127.03	120.28
1	A	77	ASP	C-N-CA	5.04	127.03	120.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2119	0	2040	23	0
1	B	2181	0	2080	17	0
1	C	2209	0	2101	15	0
1	D	2071	0	1972	16	0
1	E	2259	0	2189	17	0
1	F	2097	0	2014	12	0
2	A	1	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	1	0	0	0	0
2	E	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	F	1	0	0	0	0
3	A	41	0	0	1	0
3	B	34	0	0	0	0
3	C	60	0	0	1	0
3	D	41	0	0	0	0
3	E	82	0	0	2	0
3	F	37	0	0	0	0
All	All	13240	0	12396	96	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (96) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:60:ALA:HB1	1:D:234:VAL:HG22	1.45	0.99
1:E:110:ARG:HG2	1:E:110:ARG:HH11	1.38	0.88
1:C:110:ARG:HH11	1:C:110:ARG:HG2	1.38	0.86
1:D:110:ARG:HG2	1:D:110:ARG:HH11	1.41	0.85
1:B:110:ARG:HH11	1:B:110:ARG:HG2	1.43	0.83
1:F:110:ARG:HH11	1:F:110:ARG:HG2	1.43	0.83
1:A:110:ARG:HG2	1:A:110:ARG:HH11	1.42	0.82
1:B:243[B]:ARG:HD3	1:B:266:MSE:SE	2.40	0.72
1:F:268:PRO:HD2	1:F:318:MSE:SE	2.42	0.69
1:A:177:MSE:HG3	1:A:192[B]:MSE:SE	2.43	0.68
1:A:60:ALA:CB	1:D:234:VAL:HG22	2.23	0.68
1:A:177:MSE:HE3	1:A:192[B]:MSE:SE	2.44	0.67
1:F:38:MSE:HE3	1:F:78:ARG:HG2	1.75	0.67
1:B:177[B]:MSE:H	1:B:177[B]:MSE:SE	2.29	0.66
1:E:268:PRO:HD2	1:E:318:MSE:SE	2.45	0.66
1:A:268:PRO:HD2	1:A:318:MSE:SE	2.50	0.62
1:B:268:PRO:HD2	1:B:318:MSE:SE	2.50	0.62
1:B:177[B]:MSE:HE1	1:B:224:ASN:HB2	1.82	0.61
1:E:243[A]:ARG:HD3	1:E:266:MSE:SE	2.52	0.60
1:A:177:MSE:CE	1:A:192[B]:MSE:SE	2.99	0.59
1:E:20:VAL:HG11	1:E:311:GLN:HB3	1.85	0.59
1:D:177:MSE:HB2	1:D:192[B]:MSE:HE1	1.85	0.59
1:A:180:VAL:CG1	1:A:192[A]:MSE:HE2	2.33	0.58
1:B:117:PRO:HA	1:B:150:LEU:HB2	1.85	0.58
1:D:268:PRO:HD2	1:D:318:MSE:SE	2.54	0.58
1:C:268:PRO:HD2	1:C:318:MSE:SE	2.54	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:117:PRO:HA	1:D:150:LEU:HB2	1.87	0.56
1:A:120:THR:O	1:A:155:GLY:HA3	2.06	0.56
1:C:55:GLU:HB2	1:C:87:ARG:HG2	1.90	0.53
1:D:177:MSE:HA	1:D:192[B]:MSE:SE	2.59	0.53
1:D:120:THR:O	1:D:155:GLY:HA3	2.09	0.53
1:F:120:THR:O	1:F:155:GLY:HA3	2.10	0.52
1:A:192[B]:MSE:HE1	3:A:652:HOH:O	2.09	0.52
1:C:120:THR:O	1:C:155:GLY:HA3	2.10	0.52
1:A:177:MSE:HA	1:A:192[B]:MSE:SE	2.59	0.51
1:A:269:ASP:HB3	1:A:272:GLU:HB2	1.93	0.51
1:E:110:ARG:HH11	1:E:110:ARG:CG	2.17	0.51
1:A:117:PRO:HA	1:A:150:LEU:HB2	1.91	0.51
1:C:110:ARG:HH11	1:C:110:ARG:CG	2.16	0.51
1:A:186:ALA:HA	1:B:243[B]:ARG:HH12	1.76	0.51
1:B:120:THR:O	1:B:155:GLY:HA3	2.11	0.51
1:E:109:GLY:HA2	3:E:686:HOH:O	2.11	0.50
1:E:120:THR:O	1:E:155:GLY:HA3	2.11	0.50
1:D:110:ARG:HH11	1:D:110:ARG:CG	2.19	0.49
1:F:117:PRO:HA	1:F:150:LEU:HB2	1.94	0.49
1:C:20:VAL:HG11	1:C:311:GLN:HB3	1.95	0.48
1:E:110:ARG:HG2	1:E:110:ARG:NH1	2.17	0.48
1:C:54:CYS:HB2	1:C:87:ARG:O	2.13	0.48
1:A:180:VAL:HG11	1:A:192[A]:MSE:HE2	1.96	0.48
1:A:180:VAL:HG12	1:A:192[A]:MSE:HE2	1.96	0.47
1:B:20:VAL:HG11	1:B:311:GLN:HB3	1.96	0.47
1:E:142:ASP:HA	3:E:688:HOH:O	2.14	0.46
1:C:107:ALA:HB3	3:C:556:HOH:O	2.14	0.46
1:A:243:ARG:HD3	1:A:266:MSE:SE	2.65	0.46
1:C:59:ALA:HB3	1:C:62:GLN:HB2	1.98	0.46
1:E:117:PRO:HA	1:E:150:LEU:HB2	1.97	0.45
1:F:54:CYS:HB2	1:F:87:ARG:O	2.16	0.45
1:B:53:ASP:O	1:B:66:HIS:HE1	2.00	0.45
1:E:59:ALA:HB3	1:E:62:GLN:HB2	1.98	0.44
1:B:269:ASP:HB3	1:B:272:GLU:HB2	1.98	0.44
1:D:20:VAL:HG11	1:D:311:GLN:HB3	1.99	0.44
1:E:53:ASP:O	1:E:66:HIS:HE1	2.00	0.44
1:D:211:VAL:HG11	1:D:246:PHE:O	2.17	0.44
1:B:110:ARG:HH11	1:B:110:ARG:CG	2.20	0.44
1:D:53:ASP:O	1:D:66:HIS:HE1	2.01	0.44
1:D:69:MSE:C	1:D:69:MSE:SE	3.12	0.43
1:D:59:ALA:HB3	1:D:62:GLN:HB2	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:302:THR:O	1:E:305:TYR:HB3	2.19	0.43
1:F:34:ILE:HG22	1:F:38:MSE:HE2	2.00	0.43
1:E:280:LEU:HD21	1:E:307:TRP:HA	2.01	0.43
1:F:269:ASP:HB3	1:F:272:GLU:HB2	2.01	0.43
1:C:117:PRO:HA	1:C:150:LEU:HB2	2.00	0.42
1:C:192:MSE:HE2	1:C:192:MSE:HB3	1.96	0.42
1:A:225:VAL:HG11	1:A:254:ILE:HG23	2.02	0.42
1:C:152:GLU:HB3	1:C:175:GLY:HA3	2.02	0.42
1:E:177:MSE:HG3	1:E:192[A]:MSE:HE1	2.02	0.42
1:B:23:ALA:HB1	1:B:266:MSE:HG3	2.02	0.41
1:B:152:GLU:HB3	1:B:175:GLY:HA3	2.02	0.41
1:B:280:LEU:HD21	1:B:307:TRP:HA	2.02	0.41
1:F:225:VAL:HG11	1:F:254:ILE:HG23	2.03	0.41
1:A:110:ARG:HH11	1:A:110:ARG:CG	2.20	0.41
1:C:55:GLU:HG3	1:C:117:PRO:HG3	2.02	0.41
1:E:225:VAL:HG11	1:E:254:ILE:HG23	2.02	0.41
1:B:225:VAL:HG11	1:B:254:ILE:HG23	2.02	0.41
1:C:23:ALA:HB1	1:C:266:MSE:HG2	2.02	0.41
1:A:152:GLU:HB3	1:A:175:GLY:HA3	2.02	0.41
1:F:110:ARG:HH11	1:F:110:ARG:CG	2.20	0.41
1:D:152:GLU:HB3	1:D:175:GLY:HA3	2.02	0.41
1:F:110:ARG:HG2	1:F:110:ARG:NH1	2.22	0.41
1:A:154:HIS:CG	1:B:216:ALA:HA	2.56	0.41
1:A:23:ALA:HB1	1:A:266:MSE:HG3	2.03	0.41
1:C:110:ARG:HG2	1:C:110:ARG:NH1	2.18	0.40
1:F:152:GLU:HB3	1:F:175:GLY:HA3	2.03	0.40
1:E:321:PRO:O	1:E:323:GLU:N	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	284/332 (86%)	275 (97%)	8 (3%)	1 (0%)	30	36
1	B	293/332 (88%)	278 (95%)	15 (5%)	0	100	100
1	C	297/332 (90%)	285 (96%)	11 (4%)	1 (0%)	36	44
1	D	278/332 (84%)	268 (96%)	9 (3%)	1 (0%)	30	36
1	E	299/332 (90%)	285 (95%)	12 (4%)	2 (1%)	18	21
1	F	279/332 (84%)	268 (96%)	10 (4%)	1 (0%)	30	36
All	All	1730/1992 (87%)	1659 (96%)	65 (4%)	6 (0%)	36	44

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	322	ALA
1	E	60	ALA
1	A	60	ALA
1	C	60	ALA
1	D	60	ALA
1	F	60	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	206/244 (84%)	205 (100%)	1 (0%)	81	87
1	B	209/244 (86%)	204 (98%)	5 (2%)	43	56
1	C	212/244 (87%)	210 (99%)	2 (1%)	70	80
1	D	199/244 (82%)	196 (98%)	3 (2%)	57	69
1	E	222/244 (91%)	216 (97%)	6 (3%)	39	52
1	F	203/244 (83%)	202 (100%)	1 (0%)	81	87
All	All	1251/1464 (86%)	1233 (99%)	18 (1%)	59	70

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	110	ARG
1	B	68	GLU
1	B	101	ASP
1	B	110	ARG
1	B	260	GLN
1	B	280	LEU
1	C	55	GLU
1	C	110	ARG
1	D	65	GLU
1	D	110	ARG
1	D	269	ASP
1	E	110	ARG
1	E	192[A]	MSE
1	E	192[B]	MSE
1	E	274	GLU
1	E	280	LEU
1	E	323	GLU
1	F	110	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (28) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	81	HIS
1	A	168	ASN
1	A	311	GLN
1	B	62	GLN
1	B	66	HIS
1	B	81	HIS
1	B	168	ASN
1	B	260	GLN
1	B	264	ASN
1	B	311	GLN
1	C	81	HIS
1	C	162	GLN
1	C	168	ASN
1	C	184	HIS
1	C	244	ASN
1	C	260	GLN
1	C	311	GLN
1	D	81	HIS
1	D	264	ASN
1	D	311	GLN
1	E	81	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	244	ASN
1	E	260	GLN
1	E	264	ASN
1	E	311	GLN
1	F	81	HIS
1	F	162	GLN
1	F	311	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 9 ligands modelled in this entry, 9 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	275/332 (82%)	0.43	14 (5%)	33 30	35, 59, 97, 115	3 (1%)
1	B	285/332 (85%)	0.32	22 (7%)	19 16	27, 59, 100, 115	2 (0%)
1	C	289/332 (87%)	0.20	14 (4%)	35 33	26, 48, 97, 117	3 (1%)
1	D	272/332 (81%)	0.31	19 (6%)	22 19	33, 56, 107, 130	0
1	E	290/332 (87%)	-0.07	6 (2%)	63 64	19, 41, 81, 96	3 (1%)
1	F	274/332 (82%)	0.40	20 (7%)	21 18	32, 59, 109, 124	0
All	All	1685/1992 (84%)	0.26	95 (5%)	30 27	19, 54, 100, 130	11 (0%)

All (95) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	12	PHE	5.5
1	B	282	ALA	5.1
1	D	302	THR	4.4
1	D	303	TYR	4.4
1	A	302	THR	4.2
1	A	107	ALA	4.0
1	D	59	ALA	3.9
1	F	302	THR	3.8
1	D	279	ILE	3.8
1	A	303	TYR	3.8
1	A	308	GLU	3.8
1	D	306	PHE	3.7
1	D	304	ARG	3.6
1	D	321	PRO	3.6
1	C	184	HIS	3.6
1	A	279	ILE	3.5
1	D	58	ALA	3.5
1	C	273	VAL	3.4
1	B	280	LEU	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	281	VAL	3.4
1	F	310	LEU	3.4
1	F	309	VAL	3.3
1	F	136	ALA	3.3
1	F	321	PRO	3.3
1	C	108	GLY	3.3
1	B	236	ALA	3.3
1	E	302	THR	3.2
1	D	107	ALA	3.2
1	A	58	ALA	3.1
1	F	304	ARG	3.1
1	A	304	ARG	3.1
1	B	304	ARG	3.0
1	D	307	TRP	3.0
1	F	59	ALA	3.0
1	C	142[A]	ASP	3.0
1	C	7	PRO	2.9
1	C	302	THR	2.9
1	B	321	PRO	2.8
1	C	321	PRO	2.8
1	F	107	ALA	2.8
1	D	60	ALA	2.8
1	A	306	PHE	2.8
1	B	303	TYR	2.8
1	B	235	ILE	2.7
1	A	106	GLY	2.7
1	D	19	PRO	2.7
1	C	10	ALA	2.7
1	F	135	VAL	2.6
1	C	277	ALA	2.6
1	A	269	ASP	2.6
1	E	317	GLY	2.6
1	A	108	GLY	2.5
1	C	280	LEU	2.5
1	D	56	ASP	2.5
1	A	60	ALA	2.5
1	F	313	ALA	2.5
1	A	277	ALA	2.5
1	F	303	TYR	2.4
1	F	305	TYR	2.4
1	B	59	ALA	2.4
1	D	310	LEU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	8	ASN	2.3
1	B	11	LEU	2.3
1	C	6	HIS	2.3
1	B	270	PHE	2.3
1	B	107	ALA	2.3
1	F	29	GLY	2.2
1	B	308	GLU	2.2
1	D	270	PHE	2.2
1	F	58	ALA	2.2
1	F	227	LEU	2.2
1	E	13	ALA	2.2
1	D	305	TYR	2.2
1	F	320	VAL	2.1
1	E	60	ALA	2.1
1	E	108	GLY	2.1
1	B	10	ALA	2.1
1	B	190	ALA	2.1
1	A	20	VAL	2.1
1	B	305	TYR	2.1
1	B	9	ASP	2.1
1	C	227	LEU	2.1
1	D	61	GLY	2.1
1	F	109	GLY	2.1
1	D	308	GLU	2.1
1	B	315	VAL	2.0
1	B	239	ALA	2.0
1	B	242	ALA	2.0
1	F	60	ALA	2.0
1	B	140	GLY	2.0
1	C	317	GLY	2.0
1	D	233	GLU	2.0
1	F	279	ILE	2.0
1	E	6	HIS	2.0
1	F	270	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CL	B	404	1/1	0.79	0.15	98,98,98,98	0
2	CL	B	400	1/1	0.84	0.15	88,88,88,88	0
2	CL	E	403	1/1	0.91	0.08	81,81,81,81	0
2	CL	D	402	1/1	0.94	0.08	62,62,62,62	0
2	CL	A	405	1/1	0.94	0.11	60,60,60,60	0
2	CL	E	407	1/1	0.94	0.15	46,46,46,46	0
2	CL	C	401	1/1	0.95	0.07	47,47,47,47	0
2	CL	F	408	1/1	0.95	0.12	58,58,58,58	0
2	CL	C	406	1/1	0.97	0.06	56,56,56,56	0

6.5 Other polymers [i](#)

There are no such residues in this entry.