



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 09:06 PM UTC

PDB ID : 6REN / pdb_00006ren
Title : Crystal structure of 3fPizza6-SH with Zn²⁺
Authors : Noguchi, H.; Clarke, D.E.; Gryspeerdt, J.L.; Feyter, S.D.; Voet, A.R.D.
Deposited on : 2019-04-12
Resolution : 1.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1553 (1.30-1.30)
Clashscore	190562	1595 (1.30-1.30)
Ramachandran outliers	187476	1551 (1.30-1.30)
Sidechain outliers	187428	1551 (1.30-1.30)
RSRZ outliers	180081	1549 (1.30-1.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	256	
1	B	256	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	IPA	A	306	-	-	X	-
3	IPA	B	305	-	-	X	-
3	IPA	B	307	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4297 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 3fPizza6-SH.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	251	Total	C	N	O	0	12	0
			1883	1167	339	377			
1	B	248	Total	C	N	O	0	7	0
			1822	1130	327	365			

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	4	Total	Zn	0	0
			4	4		
2	B	4	Total	Zn	0	0
			4	4		

- Molecule 3 is ISOPROPYL ALCOHOL (CCD ID: IPA) (formula: C₃H₈O).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 3 1	0	0
3	A	1	Total C O 4 3 1	0	0
3	A	1	Total C O 4 3 1	0	0
3	A	1	Total C O 4 3 1	0	0
3	A	1	Total C O 4 3 1	0	0
3	A	1	Total C O 4 3 1	0	0
3	A	1	Total C O 4 3 1	0	0
3	B	1	Total C O 4 3 1	0	0
3	B	1	Total C O 4 3 1	0	0
3	B	1	Total C O 4 3 1	0	0
3	B	1	Total C O 4 3 1	0	0
3	B	1	Total C O 4 3 1	0	0
3	B	1	Total C O 4 3 1	0	0

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	297	Total	O	0	0
			297	297		
5	B	229	Total	O	0	0
			229	229		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 3fPizza6-SH

Chain A: 



- Molecule 1: 3fPizza6-SH

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	44.78Å 66.68Å 70.53Å 90.00° 97.92° 90.00°	Depositor
Resolution (Å)	22.09 – 1.30 22.09 – 1.30	Depositor EDS
% Data completeness (in resolution range)	99.6 (22.09-1.30) 99.5 (22.09-1.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.24 (at 1.30Å)	Xtriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, R_{free}	0.134 , 0.167 0.134 , 0.167	Depositor DCC
R_{free} test set	4935 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	13.1	Xtriage
Anisotropy	0.381	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 48.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	4297	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, GOL, IPA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.52	0/1915	0.71	2/2642 (0.1%)
1	B	0.48	0/1854	0.69	0/2556
All	All	0.50	0/3769	0.70	2/5198 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	73	HIS	N-CA-CB	5.10	118.18	110.28
1	A	31	HIS	CA-CB-CG	5.06	118.86	113.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1883	0	1865	17	0
1	B	1822	0	1798	16	0
2	A	4	0	0	0	0
2	B	4	0	0	0	0
3	A	28	0	51	10	0
3	B	24	0	45	12	0
4	A	6	0	6	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	297	0	0	0	0
5	B	229	0	0	1	0
All	All	4297	0	3765	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:305:IPA:H2	3:B:308:IPA:H13	1.55	0.89
1:B:241:HIS:CE1	3:B:305:IPA:H31	2.24	0.72
1:A:62[B]:VAL:HG12	1:A:68:VAL:HG22	1.70	0.72
1:A:36:VAL:HG21	1:A:70[B]:VAL:HG11	1.71	0.72
3:A:306:IPA:H12	4:A:312:GOL:H32	1.76	0.67
3:A:306:IPA:H12	4:A:312:GOL:H11	1.79	0.65
3:A:307:IPA:H11	3:A:310:IPA:H2	1.81	0.62
1:A:115:HIS:CE1	3:A:309:IPA:H32	2.36	0.61
1:B:31:HIS:CE1	3:B:307:IPA:H11	2.35	0.61
1:A:199:HIS:CE1	3:A:306:IPA:H11	2.36	0.60
3:A:307:IPA:C1	3:A:310:IPA:H2	2.32	0.60
3:B:307:IPA:H2	3:B:308:IPA:O2	2.02	0.60
1:A:73:HIS:HB3	1:A:99:PRO:O	2.06	0.56
1:A:79[B]:VAL:HG12	1:A:90:VAL:HG22	1.90	0.54
1:A:31:HIS:CE1	3:A:307:IPA:H11	2.43	0.54
1:B:241:HIS:HE1	3:B:305:IPA:H31	1.73	0.53
1:A:36:VAL:HG21	1:A:70[A]:VAL:HG21	1.91	0.52
1:B:199:HIS:CE1	3:B:305:IPA:H32	2.44	0.52
1:A:29[B]:THR:HG21	1:A:57:PRO:C	2.35	0.52
1:A:27:TYR:CD1	1:A:62[A]:VAL:HG21	2.45	0.51
1:A:113:THR:HG23	1:A:144:VAL:HB	1.93	0.51
1:B:31:HIS:HE1	3:B:307:IPA:H11	1.76	0.49
1:B:197[B]:THR:HG22	1:B:204:VAL:HG22	1.95	0.48
1:A:115:HIS:CE1	3:A:305:IPA:H11	2.50	0.46
1:A:69:TYR:CE1	1:A:104[B]:VAL:HG21	2.51	0.46
1:B:199:HIS:HE1	3:B:305:IPA:H32	1.80	0.45
1:B:162:VAL:HG21	1:B:196:VAL:HG21	1.99	0.44
1:A:27:TYR:CE1	1:A:62[B]:VAL:HG11	2.52	0.44
1:B:120:VAL:HG21	1:B:154:VAL:HG21	1.99	0.44
1:B:171:THR:OG1	5:B:401:HOH:O	2.21	0.44
1:B:73:HIS:CE1	3:B:307:IPA:H13	2.54	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:28:VAL:HG21	1:B:246:VAL:HG21	2.00	0.43
1:B:29:THR:HG23	1:B:60:VAL:HB	2.02	0.41
3:A:306:IPA:C1	4:A:312:GOL:H11	2.47	0.41
1:A:69:TYR:CD1	1:A:104[B]:VAL:HG21	2.56	0.41
1:B:115:HIS:CE1	3:B:306:IPA:H11	2.56	0.41
3:B:305:IPA:H13	3:B:309:IPA:H31	2.02	0.41
1:B:243[B]:ASN:OD1	1:B:245:ARG:HG2	2.19	0.41
1:A:182:THR:HG21	3:A:311:IPA:H33	2.02	0.41
1:B:73:HIS:CE1	3:B:310:IPA:H33	2.56	0.41
1:A:204:VAL:HG21	1:A:238:VAL:HG21	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	261/256 (102%)	250 (96%)	11 (4%)	0	100	100
1	B	253/256 (99%)	241 (95%)	12 (5%)	0	100	100
All	All	514/512 (100%)	491 (96%)	23 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	210/201 (104%)	210 (100%)	0	100	100
1	B	202/201 (100%)	202 (100%)	0	100	100
All	All	412/402 (102%)	412 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	201	ASN
1	B	86	ASN
1	B	202	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 22 ligands modelled in this entry, 8 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	IPA	A	307	-	3,3,3	0.36	0	3,3,3	0.74	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	IPA	B	305	2	3,3,3	0.72	0	3,3,3	0.70	0
3	IPA	A	305	2	3,3,3	0.94	0	3,3,3	1.31	0
3	IPA	A	306	2	3,3,3	0.76	0	3,3,3	1.10	0
3	IPA	A	311	-	3,3,3	0.48	0	3,3,3	0.44	0
3	IPA	B	310	-	3,3,3	0.56	0	3,3,3	0.27	0
3	IPA	B	309	-	3,3,3	0.54	0	3,3,3	0.27	0
3	IPA	B	306	2	3,3,3	0.45	0	3,3,3	0.40	0
3	IPA	A	309	-	3,3,3	0.47	0	3,3,3	0.43	0
4	GOL	A	312	-	5,5,5	7.34	1 (20%)	5,5,5	0.97	0
3	IPA	B	308	-	3,3,3	0.54	0	3,3,3	0.39	0
3	IPA	B	307	2	3,3,3	0.49	0	3,3,3	0.41	0
3	IPA	A	308	-	3,3,3	0.54	0	3,3,3	0.33	0
3	IPA	A	310	2	3,3,3	0.47	0	3,3,3	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	312	-	-	3/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	312	GOL	O2-C2	-16.36	0.95	1.43

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	312	GOL	C1-C2-C3-O3
4	A	312	GOL	O2-C2-C3-O3
4	A	312	GOL	O1-C1-C2-O2

There are no ring outliers.

13 monomers are involved in 22 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	307	IPA	3	0
3	B	305	IPA	6	0
3	A	305	IPA	1	0
3	A	306	IPA	4	0
3	A	311	IPA	1	0
3	B	310	IPA	1	0
3	B	309	IPA	1	0
3	B	306	IPA	1	0
3	A	309	IPA	1	0
4	A	312	GOL	3	0
3	B	308	IPA	2	0
3	B	307	IPA	4	0
3	A	310	IPA	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	251/256 (98%)	-0.27	7 (2%) 55 61	5, 14, 24, 30	12 (4%)
1	B	248/256 (96%)	-0.11	6 (2%) 59 65	6, 17, 28, 40	7 (2%)
All	All	499/512 (97%)	-0.19	13 (2%) 57 63	5, 15, 26, 40	19 (3%)

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	3	THR	4.3
1	B	168	GLY	3.6
1	A	2	ASN	3.5
1	A	1	SER	3.2
1	B	170	ASN	2.9
1	B	84	GLY	2.8
1	B	83	ALA	2.8
1	A	90	VAL	2.6
1	A	86	ASN	2.4
1	B	171	THR	2.2
1	A	126	GLY	2.1
1	B	251	ALA	2.1
1	A	87	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	IPA	A	308	4/4	0.73	0.14	28,28,29,29	0
3	IPA	B	309	4/4	0.74	0.16	27,27,30,34	0
3	IPA	B	310	4/4	0.77	0.16	25,27,29,32	0
3	IPA	A	311	4/4	0.79	0.15	25,27,27,29	0
3	IPA	A	309	4/4	0.80	0.14	25,30,30,34	0
3	IPA	A	307	4/4	0.84	0.16	29,29,30,31	0
3	IPA	B	308	4/4	0.86	0.12	28,29,29,32	0
3	IPA	A	310	4/4	0.88	0.16	23,27,27,30	0
3	IPA	B	307	4/4	0.91	0.10	23,25,26,27	0
3	IPA	B	305	4/4	0.91	0.11	16,18,19,23	0
4	GOL	A	312	6/6	0.93	0.10	15,22,35,36	0
3	IPA	B	306	4/4	0.96	0.09	22,25,25,26	0
3	IPA	A	306	4/4	0.96	0.08	14,16,19,23	0
3	IPA	A	305	4/4	0.97	0.07	11,15,17,18	0
2	ZN	B	303	1/1	1.00	0.01	13,13,13,13	0
2	ZN	B	304	1/1	1.00	0.04	15,15,15,15	0
2	ZN	A	301	1/1	1.00	0.01	11,11,11,11	0
2	ZN	A	302	1/1	1.00	0.01	12,12,12,12	0
2	ZN	A	303	1/1	1.00	0.00	11,11,11,11	0
2	ZN	A	304	1/1	1.00	0.04	14,14,14,14	0
2	ZN	B	301	1/1	1.00	0.01	15,15,15,15	0
2	ZN	B	302	1/1	1.00	0.03	17,17,17,17	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around ZN B 303:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN B 304:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN A 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN A 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN A 303:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN A 304:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN B 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ZN B 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.