



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 12:46 AM UTC

PDB ID : 4RGA / pdb_00004rga
Title : Phage 1358 receptor binding protein in complex with the trisaccharide GlcNAc-Galf-GlcOMe
Authors : Spinelli, S.; McCabe, O.; Farenc, C.; Tremblay, D.; Blangy, S.; Oscarson, S.; Moineau, S.; Cambillau, C.
Deposited on : 2014-09-29
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	393	14% 88% 10% ..
1	B	393	15% 88% 10% ..
2	C	3	33% 67%
2	D	3	33% 67%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 6995 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Phage 1358 receptor binding protein (ORF20).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	390	Total	C	N	O	S	0	1	0
			3039	1908	526	596	9			
1	B	390	Total	C	N	O	S	0	1	0
			3039	1908	526	596	9			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	expression tag	UNP D3W0F1
B	1	GLY	-	expression tag	UNP D3W0F1

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-beta-D-galactofuranose-(1-6)-methyl alpha-D-glucopyranoside.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	3	Total	C	N	O	0	0	0
			38	21	1	16			
2	D	3	Total	C	N	O	0	0	0
			38	21	1	16			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	423	Total	O	0	0
			423	423		
3	B	418	Total	O	0	0
			418	418		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phage 1358 receptor binding protein (ORF20)




- Molecule 1: Phage 1358 receptor binding protein (ORF20)



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-beta-D-galactofuranose-(1-6)-methyl alpha-D-glucopyranoside



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-beta-D-galactofuranose-(1-6)-methyl alpha-D-glucopyranoside



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 3	Depositor
Cell constants a, b, c, α , β , γ	166.37Å 166.37Å 166.37Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.00 – 2.10 48.00 – 2.10	Depositor EDS
% Data completeness (in resolution range)	100.0 (48.00-2.10) 100.0 (48.00-2.10)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.23 (at 2.10Å)	Xtriage
Refinement program	BUSTER 2.11.5	Depositor
R, R_{free}	0.180 , 0.192 0.184 , 0.199	Depositor DCC
R_{free} test set	4462 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	35.3	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 81.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.034 for l,-k,h	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6995	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.87% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GZL, GYP, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.81	0/3105	1.13	10/4230 (0.2%)
1	B	0.81	0/3105	1.14	10/4230 (0.2%)
All	All	0.81	0/6210	1.14	20/8460 (0.2%)

There are no bond length outliers.

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	106	THR	N-CA-C	-8.04	98.56	110.46
1	A	106	THR	N-CA-C	-7.82	98.89	110.46
1	A	340	ASP	CA-CB-CG	7.14	119.74	112.60
1	A	104	VAL	N-CA-C	-6.70	100.05	108.89
1	B	104	VAL	N-CA-C	-6.42	100.42	108.89
1	B	178	GLY	CA-C-N	5.97	129.41	122.85
1	B	178	GLY	C-N-CA	5.97	129.41	122.85
1	A	338	LYS	CG-CD-CE	5.91	124.90	111.30
1	B	177	SER	N-CA-C	-5.61	103.23	110.19
1	A	256	ASP	CA-CB-CG	5.34	117.94	112.60
1	B	122	ASP	CA-CB-CG	5.31	117.91	112.60
1	B	340	ASP	CA-CB-CG	5.21	117.81	112.60
1	B	338	LYS	CG-CD-CE	5.20	123.27	111.30
1	B	256	ASP	CA-CB-CG	5.11	117.71	112.60
1	A	122	ASP	CA-CB-CG	5.11	117.71	112.60
1	B	391	GLY	N-CA-C	5.10	120.33	114.16
1	A	106	THR	CA-C-N	-5.02	114.91	122.24
1	A	106	THR	C-N-CA	-5.02	114.91	122.24
1	A	217	ALA	N-CA-C	5.02	117.01	109.23
1	A	99	VAL	N-CA-C	5.01	112.51	107.55

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3039	0	2905	16	0
1	B	3039	0	2905	14	0
2	C	38	0	33	0	0
2	D	38	0	33	0	0
3	A	423	0	0	0	0
3	B	418	0	0	0	0
All	All	6995	0	5876	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:196:ASN:H	1:B:201:GLN:HE22	1.20	0.88
1:A:196:ASN:H	1:A:201:GLN:HE22	1.22	0.83
1:A:133:ASP:OD1	1:A:338:LYS:NZ	2.22	0.72
1:A:237:ARG:NH2	1:A:341:ASN:HD21	1.94	0.65
1:B:133:ASP:OD1	1:B:338:LYS:NZ	2.33	0.62
1:A:200:VAL:HG22	1:A:347:GLU:HG2	1.85	0.58
1:B:29:ARG:HH12	1:B:176:ASN:ND2	2.05	0.54
1:A:40:ILE:HD12	1:A:164:TYR:HE2	1.73	0.54
1:B:40:ILE:HD12	1:B:164:TYR:HE2	1.73	0.53
1:B:176:ASN:O	1:B:180:PRO:HA	2.09	0.53
1:B:237:ARG:NH2	1:B:341:ASN:HD21	2.06	0.53
1:B:312[B]:ARG:HH12	1:B:347:GLU:CD	2.18	0.52
1:B:269:PRO:HA	1:B:273:ASN:HD21	1.80	0.47
1:A:19:ALA:O	1:A:59:GLY:HA3	2.15	0.47
1:B:19:ALA:O	1:B:59:GLY:HA3	2.15	0.47
1:A:269:PRO:HA	1:A:273:ASN:HD21	1.80	0.46
1:A:196:ASN:N	1:A:201:GLN:HE22	2.02	0.46
1:A:14:THR:H	1:A:17:ASN:HD22	1.64	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:237:ARG:HH22	1:A:341:ASN:HD21	1.60	0.45
1:A:14:THR:HB	1:A:15:PRO:HD2	1.98	0.45
1:B:208:VAL:HG22	1:B:222:ASN:HD22	1.82	0.45
1:B:270:ALA:H	1:B:273:ASN:HD21	1.65	0.44
1:A:208:VAL:HG22	1:A:222:ASN:HD22	1.82	0.44
1:A:312[A]:ARG:NH2	1:A:347:GLU:OE1	2.36	0.43
1:A:144:ARG:HG2	1:A:144:ARG:HH11	1.83	0.43
1:A:270:ALA:H	1:A:273:ASN:HD21	1.66	0.43
1:B:14:THR:HB	1:B:15:PRO:HD2	2.02	0.41
1:B:38:ILE:HD11	1:B:48:LEU:HD11	2.02	0.41
1:B:179:ASN:HD22	1:B:182:TYR:HB2	1.86	0.40
1:A:325:GLU:OE2	1:A:370:ARG:HD3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	387/393 (98%)	375 (97%)	12 (3%)	0	100	100
1	B	387/393 (98%)	376 (97%)	10 (3%)	1 (0%)	36	36
All	All	774/786 (98%)	751 (97%)	22 (3%)	1 (0%)	48	50

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	178	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	328/332 (99%)	314 (96%)	14 (4%)	26	27
1	B	328/332 (99%)	314 (96%)	14 (4%)	26	27
All	All	656/664 (99%)	628 (96%)	28 (4%)	26	27

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	ILE
1	A	18	GLU
1	A	34	ASP
1	A	38	ILE
1	A	50	LEU
1	A	75	LEU
1	A	117	ARG
1	A	145	VAL
1	A	146	ILE
1	A	173	ASP
1	A	179	ASN
1	A	273	ASN
1	A	323	MET
1	A	370	ARG
1	B	6	ILE
1	B	18	GLU
1	B	34	ASP
1	B	38	ILE
1	B	50	LEU
1	B	75	LEU
1	B	117	ARG
1	B	144	ARG
1	B	145	VAL
1	B	173	ASP
1	B	179	ASN
1	B	273	ASN
1	B	323	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	370	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (27) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	17	ASN
1	A	28	ASN
1	A	52	HIS
1	A	83	ASN
1	A	157	GLN
1	A	179	ASN
1	A	201	GLN
1	A	222	ASN
1	A	273	ASN
1	A	341	ASN
1	A	345	GLN
1	A	371	ASN
1	A	383	ASN
1	B	28	ASN
1	B	52	HIS
1	B	83	ASN
1	B	157	GLN
1	B	176	ASN
1	B	179	ASN
1	B	196	ASN
1	B	201	GLN
1	B	222	ASN
1	B	273	ASN
1	B	341	ASN
1	B	345	GLN
1	B	371	ASN
1	B	383	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	GYP	C	1	2	13,13,13	0.66	0	18,18,18	2.13	6 (33%)
2	GZL	C	2	2	11,11,12	0.39	0	13,15,17	1.25	0
2	NAG	C	3	2	14,14,15	0.73	0	17,19,21	1.32	2 (11%)
2	GYP	D	1	2	13,13,13	0.75	0	18,18,18	1.67	4 (22%)
2	GZL	D	2	2	11,11,12	0.37	0	13,15,17	0.97	0
2	NAG	D	3	2	14,14,15	0.67	0	17,19,21	1.13	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GYP	C	1	2	-	2/4/24/24	0/1/1/1
2	GZL	C	2	2	-	4/6/19/22	0/1/1/1
2	NAG	C	3	2	-	0/6/23/26	0/1/1/1
2	GYP	D	1	2	-	2/4/24/24	0/1/1/1
2	GZL	D	2	2	-	4/6/19/22	0/1/1/1
2	NAG	D	3	2	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1	GYP	O1-C1-C2	5.47	114.45	108.14
2	C	3	NAG	C1-C2-N2	-4.29	103.67	110.43
2	D	1	GYP	O1-C1-C2	3.78	112.50	108.14
2	C	1	GYP	C3-C4-C5	-3.45	103.98	110.23
2	D	3	NAG	C1-C2-N2	-3.26	105.29	110.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	1	GYP	O6-C6-C5	2.68	120.47	111.33
2	C	1	GYP	O5-C1-C2	2.57	115.64	110.37
2	C	1	GYP	C7-O1-C1	-2.49	109.48	113.26
2	D	1	GYP	C3-C4-C5	-2.30	106.06	110.23
2	C	1	GYP	O2-C2-C3	2.15	115.45	110.38
2	C	3	NAG	O4-C4-C5	2.08	114.44	109.32
2	C	1	GYP	C1-O5-C5	2.05	117.72	113.72
2	D	1	GYP	C1-O5-C5	2.04	117.71	113.72

There are no chirality outliers.


All (12) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	1	GYP	C2-C1-O1-C7
2	D	1	GYP	C2-C1-O1-C7
2	D	2	GZL	C4-C5-C6-O6
2	D	2	GZL	O5-C5-C6-O6
2	C	1	GYP	O5-C1-O1-C7
2	D	1	GYP	O5-C1-O1-C7
2	C	2	GZL	C4-C5-C6-O6
2	C	2	GZL	O4-C4-C5-O5
2	D	2	GZL	O4-C4-C5-O5
2	C	2	GZL	O5-C5-C6-O6
2	C	2	GZL	O4-C4-C5-C6
2	D	2	GZL	O4-C4-C5-C6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	390/393 (99%)	0.52	55 (14%) 6 6	20, 40, 90, 132	1 (0%)
1	B	390/393 (99%)	0.49	58 (14%) 5 5	23, 40, 85, 115	1 (0%)
All	All	780/786 (99%)	0.50	113 (14%) 6 6	20, 40, 88, 132	2 (0%)

All (113) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	174	GLY	7.8
1	A	68	ALA	7.2
1	B	68	ALA	7.1
1	A	275	TRP	6.4
1	B	176	ASN	6.2
1	A	176	ASN	5.8
1	B	175	GLY	5.8
1	B	105	GLY	5.8
1	B	275	TRP	5.7
1	B	178	GLY	5.3
1	B	104	VAL	5.3
1	B	18	GLU	5.2
1	B	270	ALA	5.2
1	A	174	GLY	5.1
1	B	268	PHE	5.0
1	A	266	TYR	5.0
1	B	106	THR	5.0
1	A	71	ALA	4.8
1	A	18	GLU	4.8
1	A	2	VAL	4.3
1	B	140	ILE	4.2
1	B	177	SER	4.2
1	A	155	PHE	4.2
1	B	67	ALA	4.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	277	VAL	4.1
1	A	140	ILE	4.0
1	A	67	ALA	3.9
1	A	268	PHE	3.9
1	B	2	VAL	3.9
1	A	106	THR	3.9
1	B	15	PRO	3.8
1	A	20	SER	3.7
1	B	7	ASN	3.7
1	A	173	ASP	3.7
1	A	175	GLY	3.7
1	A	40	ILE	3.6
1	A	276	ARG	3.5
1	B	16	GLU	3.5
1	A	160	THR	3.5
1	A	104	VAL	3.5
1	B	97	VAL	3.5
1	A	323	MET	3.4
1	A	272	GLY	3.4
1	B	40	ILE	3.4
1	B	271	SER	3.4
1	A	44	THR	3.4
1	B	173	ASP	3.4
1	B	276	ARG	3.3
1	B	13	VAL	3.3
1	B	71	ALA	3.2
1	A	85	THR	3.2
1	A	16	GLU	3.2
1	A	158	THR	3.1
1	A	78	THR	3.1
1	A	161	THR	3.0
1	A	177	SER	3.0
1	A	105	GLY	2.9
1	A	159	GLY	2.8
1	A	7	ASN	2.7
1	A	19	ALA	2.7
1	A	274	SER	2.7
1	B	44	THR	2.7
1	A	17	ASN	2.7
1	A	273	ASN	2.7
1	B	73	ASP	2.7
1	B	46	ASN	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	139	LEU	2.6
1	A	269	PRO	2.6
1	B	79	THR	2.6
1	B	269	PRO	2.6
1	B	48	LEU	2.6
1	B	66	ARG	2.5
1	B	277	VAL	2.5
1	B	14	THR	2.5
1	B	159	GLY	2.5
1	A	157	GLN	2.5
1	B	179	ASN	2.5
1	B	6	ILE	2.5
1	B	107	ASN	2.4
1	A	49	ALA	2.4
1	A	15	PRO	2.4
1	B	41	LEU	2.3
1	A	21	ILE	2.3
1	B	43	THR	2.3
1	B	160	THR	2.3
1	B	65	THR	2.3
1	A	179	ASN	2.3
1	A	271	SER	2.2
1	B	20	SER	2.2
1	A	142	PRO	2.2
1	B	45	SER	2.2
1	B	155	PHE	2.2
1	A	43	THR	2.2
1	B	21	ILE	2.2
1	B	83	ASN	2.2
1	B	74	VAL	2.2
1	B	28	ASN	2.2
1	A	24	LEU	2.1
1	A	46	ASN	2.1
1	B	154	THR	2.1
1	B	331	VAL	2.1
1	A	73	ASP	2.1
1	B	238	ASP	2.1
1	A	312[A]	ARG	2.1
1	B	103	THR	2.1
1	B	163	LYS	2.1
1	B	84	GLU	2.0
1	A	139	LEU	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	42	ASN	2.0
1	B	42	ASN	2.0
1	A	81	GLN	2.0
1	A	154	THR	2.0
1	A	83	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GZL	C	2	11/12	0.78	0.15	65,71,76,77	0
2	GZL	D	2	11/12	0.78	0.16	66,76,83,83	0
2	GYP	D	1	13/13	0.88	0.13	39,52,63,67	0
2	GYP	C	1	13/13	0.88	0.15	33,48,60,66	0
2	NAG	C	3	14/15	0.90	0.13	41,55,61,61	0
2	NAG	D	3	14/15	0.90	0.15	39,58,61,62	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.