



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 4, 2026 – 06:06 PM EDT

PDB ID : 3RNT / pdb_00003rnt
Title : CRYSTAL STRUCTURE OF GUANOSINE-FREE RIBONUCLEASE T1,
COMPLEXED WITH VANADATE(V), SUGGESTS CONFORMATIONAL
CHANGE UPON SUBSTRATE BINDING
Authors : Kostrewa, D.; Choe, H.-W.; Heinemann, U.; Saenger, W.
Deposited on : 1989-05-31
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	104	 47% 48% . .

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 954 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called RIBONUCLEASE T1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	104	Total	C	N	O	S	0	2	0
			786	483	128	171	4			

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ca	0	0
			1	1		

- Molecule 3 is VANADATE ION (CCD ID: VO4) (formula: O₄V).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	V	0	0
			5	4	1		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	160	Total 162	O 162	0	2

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: RIBONUCLEASE T1

Chain A: 



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	48.82Å 46.53Å 41.20Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 1.80	Depositor
% Data completeness (in resolution range)	(Not available) (10.00-1.80)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	PROFFT	Depositor
R, R_{free}	0.137 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	954	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, VO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.81	16/817 (2.0%)	2.92	85/1113 (7.6%)

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	29	ASP	C-N	-7.03	1.26	1.33
1	A	59	TRP	CA-C	6.18	1.59	1.52
1	A	78[A]	VAL	CA-C	6.12	1.60	1.52
1	A	78[B]	VAL	CA-C	6.12	1.60	1.52
1	A	70	GLY	C-O	5.96	1.28	1.23
1	A	61	ILE	CA-C	5.93	1.60	1.52
1	A	90	ILE	CA-CB	5.59	1.63	1.54
1	A	91	THR	CA-CB	5.57	1.63	1.53
1	A	20	GLN	C-O	5.43	1.30	1.24
1	A	89	VAL	C-N	-5.25	1.27	1.33
1	A	69	SER	C-O	5.17	1.31	1.24
1	A	70	GLY	N-CA	5.14	1.51	1.45
1	A	24	TYR	C-O	5.13	1.30	1.24
1	A	57	TYR	CA-C	5.11	1.58	1.52
1	A	35	SER	C-O	5.06	1.30	1.24
1	A	10	CYS	C-O	5.01	1.30	1.24

All (85) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	84	ASN	OD1-CG-ND2	-17.42	105.18	122.60
1	A	46	GLU	CB-CG-CD	16.01	139.82	112.60
1	A	31	GLU	CG-CD-OE1	11.31	144.42	118.40
1	A	53	SER	CA-C-N	10.92	138.50	121.03
1	A	53	SER	C-N-CA	10.92	138.50	121.03

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	44	ASN	OD1-CG-ND2	10.51	133.11	122.60
1	A	50	PHE	CA-CB-CG	10.34	124.14	113.80
1	A	25	LYS	CA-CB-CG	9.35	132.81	114.10
1	A	76	ASP	O-C-N	9.35	133.85	123.10
1	A	59	TRP	O-C-N	9.28	131.68	121.94
1	A	99	ASN	N-CA-CB	9.26	123.68	110.44
1	A	17	SER	CA-CB-OG	-9.20	92.70	111.10
1	A	93	THR	CA-C-O	-8.90	110.77	120.92
1	A	93	THR	CA-CB-OG1	-8.72	96.52	109.60
1	A	58	GLU	CA-C-O	8.04	129.38	120.70
1	A	70	GLY	N-CA-C	-7.96	102.48	111.63
1	A	85	GLN	CB-CG-CD	-7.72	99.47	112.60
1	A	61	ILE	N-CA-C	-7.60	97.47	108.11
1	A	31	GLU	O-C-N	7.58	133.05	123.19
1	A	52	VAL	CB-CA-C	7.50	123.03	110.95
1	A	99	ASN	OD1-CG-ND2	7.40	130.00	122.60
1	A	31	GLU	CG-CD-OE2	-7.09	102.10	118.40
1	A	38	TYR	O-C-N	7.02	129.68	121.39
1	A	54	SER	N-CA-CB	-6.97	100.01	110.05
1	A	84	ASN	O-C-N	-6.90	113.16	122.41
1	A	75	ALA	CB-CA-C	-6.88	97.68	109.65
1	A	99	ASN	N-CA-C	-6.88	101.66	110.53
1	A	49	ASP	CA-CB-CG	-6.73	105.87	112.60
1	A	89	VAL	CB-CA-C	6.71	119.49	110.42
1	A	55	PRO	O-C-N	6.70	133.83	121.10
1	A	65	GLY	N-CA-C	-6.68	105.86	115.27
1	A	75	ALA	N-CA-CB	-6.66	100.61	110.53
1	A	83	ASN	CB-CG-OD1	-6.65	107.50	120.80
1	A	26	LEU	CB-CG-CD1	6.58	130.43	110.70
1	A	20	GLN	OE1-CD-NE2	-6.45	116.15	122.60
1	A	10	CYS	N-CA-CB	-6.41	100.50	110.57
1	A	103	CYS	N-CA-C	-6.33	101.11	110.48
1	A	58	GLU	CB-CG-CD	6.29	123.30	112.60
1	A	37	SER	N-CA-C	6.28	118.87	111.02
1	A	98[A]	ASN	CB-CA-C	6.21	121.09	111.28
1	A	98[B]	ASN	CB-CA-C	6.21	121.09	111.28
1	A	5	THR	N-CA-CB	6.13	121.40	110.80
1	A	88	GLY	O-C-N	6.12	130.71	123.61
1	A	76	ASP	CA-C-O	-6.09	114.10	121.28
1	A	52	VAL	N-CA-CB	-6.08	97.46	111.81
1	A	59	TRP	CA-C-N	-6.04	113.72	119.76
1	A	59	TRP	C-N-CA	-6.04	113.72	119.76

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	83	ASN	CB-CG-ND2	6.03	125.45	116.40
1	A	13	SER	N-CA-C	-6.00	104.74	111.28
1	A	17	SER	CA-C-N	5.97	128.21	120.44
1	A	17	SER	C-N-CA	5.97	128.21	120.44
1	A	28	GLU	CB-CA-C	-5.97	99.21	110.67
1	A	59	TRP	CA-C-O	-5.95	113.60	120.03
1	A	44	ASN	CA-CB-CG	5.95	118.55	112.60
1	A	47	GLY	N-CA-C	5.81	125.78	114.16
1	A	85	GLN	OE1-CD-NE2	-5.74	116.86	122.60
1	A	89	VAL	CA-CB-CG2	5.72	120.12	110.40
1	A	29	ASP	CA-C-N	5.67	131.37	120.66
1	A	29	ASP	C-N-CA	5.67	131.37	120.66
1	A	88	GLY	CA-C-O	-5.65	116.26	121.63
1	A	91	THR	CA-CB-OG1	-5.64	101.14	109.60
1	A	78[A]	VAL	N-CA-C	-5.63	100.32	108.71
1	A	78[B]	VAL	N-CA-C	-5.63	100.32	108.71
1	A	96	SER	O-C-N	5.60	129.28	122.96
1	A	26	LEU	O-C-N	5.59	128.53	122.15
1	A	13	SER	CA-C-O	-5.54	114.68	120.55
1	A	89	VAL	N-CA-C	-5.51	99.62	107.77
1	A	84	ASN	CA-C-O	5.50	127.82	121.28
1	A	25	LYS	CB-CG-CD	-5.43	98.81	111.30
1	A	84	ASN	CB-CG-OD1	5.42	131.63	120.80
1	A	1	ALA	N-CA-CB	5.42	118.52	110.40
1	A	8	SER	N-CA-CB	-5.41	102.47	110.53
1	A	32	THR	O-C-N	5.37	129.67	123.17
1	A	17	SER	CA-C-O	-5.36	115.19	120.82
1	A	66	ASP	O-C-N	5.34	128.94	122.85
1	A	104	THR	CA-CB-OG1	-5.31	101.64	109.60
1	A	8	SER	CB-CA-C	-5.29	100.45	109.65
1	A	48	PHE	O-C-N	5.27	129.32	122.89
1	A	74	GLY	N-CA-C	-5.22	101.50	112.45
1	A	46	GLU	CB-CA-C	-5.21	99.57	109.95
1	A	84	ASN	CA-C-N	5.17	129.86	122.72
1	A	84	ASN	C-N-CA	5.17	129.86	122.72
1	A	44	ASN	CB-CG-OD1	-5.14	110.52	120.80
1	A	2	CYS	CA-CB-SG	5.11	126.16	114.40
1	A	58	GLU	N-CA-CB	5.10	119.71	110.83

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	786	0	685	2	0
2	A	1	0	0	0	0
3	A	5	0	0	0	0
4	A	162	0	0	1	0
All	All	954	0	685	2	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (2) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:25:LYS:HE2	1:A:26:LEU:HD13	1.81	0.63
1:A:5:THR:HG21	4:A:127:HOH:O	2.08	0.52

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	104/104 (100%)	103 (99%)	1 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	87/85 (102%)	80 (92%)	7 (8%)	11 3

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	THR
1	A	25	LYS
1	A	26	LEU
1	A	52	VAL
1	A	54	SER
1	A	86	LEU
1	A	102	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	20	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	VO4	A	106	-	0,4,4	-	-	-		

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.