



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 03:47 PM UTC

PDB ID : 4S3Q / pdb_00004s3q
Title : Amylomaltase MalQ from Escherichia coli in complex with maltose
Authors : Weiss, S.C.; Schiefner, A.
Deposited on : 2015-03-26
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	696	3% 92% 6%
1	B	696	3% 91% 8%
1	C	696	3% 92% 6%
2	D	2	100%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 16868 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 4-alpha-glucanotransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	682	Total	C	N	O	S	0	0	0
			5440	3458	945	1005	32			
1	B	686	Total	C	N	O	S	0	0	0
			5480	3482	957	1009	32			
1	C	681	Total	C	N	O	S	0	0	0
			5430	3452	942	1004	32			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	689	SER	-	expression tag	UNP P15977
A	690	ALA	-	expression tag	UNP P15977
A	691	HIS	-	expression tag	UNP P15977
A	692	HIS	-	expression tag	UNP P15977
A	693	HIS	-	expression tag	UNP P15977
A	694	HIS	-	expression tag	UNP P15977
A	695	HIS	-	expression tag	UNP P15977
A	696	HIS	-	expression tag	UNP P15977
B	689	SER	-	expression tag	UNP P15977
B	690	ALA	-	expression tag	UNP P15977
B	691	HIS	-	expression tag	UNP P15977
B	692	HIS	-	expression tag	UNP P15977
B	693	HIS	-	expression tag	UNP P15977
B	694	HIS	-	expression tag	UNP P15977
B	695	HIS	-	expression tag	UNP P15977
B	696	HIS	-	expression tag	UNP P15977
C	689	SER	-	expression tag	UNP P15977
C	690	ALA	-	expression tag	UNP P15977
C	691	HIS	-	expression tag	UNP P15977
C	692	HIS	-	expression tag	UNP P15977
C	693	HIS	-	expression tag	UNP P15977
C	694	HIS	-	expression tag	UNP P15977
C	695	HIS	-	expression tag	UNP P15977

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
C	696	HIS	-	expression tag	UNP P15977

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	D	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	C	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is IODIDE ION (CCD ID: IOD) (formula: I).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	3	Total I 3 3	0	0
4	B	3	Total I 3 3	0	0
4	C	2	Total I 2 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	174	Total O 174 174	0	0
5	B	156	Total O 156 156	0	0
5	C	137	Total O 137 137	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: 4-alpha-glucanotransferase



• Molecule 1: 4-alpha-glucanotransferase



• Molecule 1: 4-alpha-glucanotransferase



HIS
HIS
HIS
HIS
HIS
HIS

- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain D:

100%

GLC1
GLC2

4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	75.13Å 77.02Å 128.40Å 76.11° 75.48° 66.31°	Depositor
Resolution (Å)	34.23 – 2.10 34.23 – 2.10	Depositor EDS
% Data completeness (in resolution range)	94.3 (34.23-2.10) 94.3 (34.23-2.10)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.90 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.221 , 0.258 0.225 , 0.262	Depositor DCC
R_{free} test set	6970 reflections (4.74%)	wwPDB-VP
Wilson B-factor (Å ²)	29.8	Xtriage
Anisotropy	0.829	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 39.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.030 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	16868	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.25% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, IOD, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.68	0/5586	0.84	0/7592
1	B	0.64	0/5630	0.83	0/7652
1	C	0.66	0/5575	0.83	0/7577
All	All	0.66	0/16791	0.83	0/22821

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5440	0	5292	30	0
1	B	5480	0	5320	34	0
1	C	5430	0	5285	23	0
2	D	23	0	21	0	0
3	A	8	0	12	0	0
3	B	8	0	12	1	0
3	C	4	0	6	3	0
4	A	3	0	0	1	0
4	B	3	0	0	0	0
4	C	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	174	0	0	3	0
5	B	156	0	0	1	0
5	C	137	0	0	2	0
All	All	16868	0	15948	85	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (85) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:283:GLN:NE2	1:B:287:GLU:OE2	2.26	0.65
1:B:137:LEU:HD22	1:B:631:ALA:HB1	1.81	0.61
1:C:123:ARG:NH1	1:C:429:GLU:OE2	2.30	0.60
1:A:294:TRP:CH2	1:A:325:VAL:HG22	2.35	0.60
1:A:690:ALA:O	1:A:691:HIS:C	2.45	0.59
1:A:268:LYS:O	1:A:272:GLN:NE2	2.36	0.59
1:A:574:VAL:HA	1:A:577:ARG:HD2	1.83	0.58
4:A:706:IOD:I	5:A:880:HOH:O	2.87	0.57
1:A:130:ARG:NH2	5:A:832:HOH:O	2.37	0.57
1:C:190:LEU:N	3:C:701:EDO:H12	2.19	0.57
1:C:210:ASN:HA	3:C:701:EDO:O1	2.05	0.57
1:C:192:PRO:HD3	3:C:701:EDO:H11	1.86	0.57
1:A:294:TRP:CZ3	1:A:325:VAL:CG2	2.88	0.56
1:C:429:GLU:O	1:C:433:GLU:HG2	2.06	0.56
1:C:451:MET:HE1	1:C:502:PRO:HD3	1.88	0.55
1:B:130:ARG:NH2	5:B:939:HOH:O	2.39	0.55
1:B:192:PRO:HB3	1:B:260:MET:HE3	1.90	0.54
1:B:42:HIS:O	1:B:43:GLN:HG2	2.08	0.54
1:B:454:LEU:HD22	1:B:505:ILE:HG13	1.90	0.53
1:A:68:MET:HE3	1:A:111:LEU:HD22	1.90	0.53
1:A:335:GLU:HG2	1:B:534:LYS:HE3	1.91	0.53
1:A:335:GLU:CG	1:B:534:LYS:HE3	2.38	0.53
1:B:541:MET:HE2	1:B:634:GLY:HA3	1.92	0.52
1:A:68:MET:HE3	1:A:111:LEU:CD2	2.39	0.52
1:C:372:TYR:CE1	1:C:494:ILE:HD13	2.45	0.51
1:C:404:ASP:O	1:C:408:PRO:HA	2.11	0.51
1:B:313:TRP:CZ3	1:B:382:GLY:HA3	2.45	0.51
1:B:640:TRP:CE2	1:B:680:LEU:HD21	2.46	0.50
1:A:26:PRO:HG2	5:C:860:HOH:O	2.12	0.49
1:B:262:ALA:HA	1:B:265:MET:CE	2.43	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:262:ALA:HA	1:B:265:MET:HE3	1.94	0.49
1:A:315:TRP:HA	1:A:318:TRP:CD2	2.47	0.49
1:C:453:MET:CE	1:C:479:LEU:HD23	2.43	0.49
1:B:453:MET:CE	1:B:479:LEU:HD23	2.43	0.48
1:B:165:GLY:HA2	3:B:701:EDO:C2	2.44	0.48
1:C:541:MET:HE3	1:C:632:LEU:HD13	1.95	0.47
1:A:154:ARG:NH2	1:A:248:ASP:HA	2.29	0.47
1:B:69:VAL:HG22	1:B:97:ALA:HB2	1.95	0.47
1:A:326:ASP:O	1:A:331:ARG:NH1	2.47	0.47
1:A:404:ASP:OD1	1:A:407:GLY:N	2.46	0.47
1:A:333:PHE:CD1	1:A:333:PHE:C	2.93	0.46
1:A:294:TRP:CH2	1:A:325:VAL:CG2	2.99	0.46
1:A:574:VAL:HG22	1:A:577:ARG:NH1	2.31	0.46
1:B:372:TYR:CE1	1:B:446:ARG:HB2	2.51	0.46
1:C:315:TRP:HA	1:C:318:TRP:CD2	2.49	0.46
1:A:335:GLU:HG2	1:B:534:LYS:CE	2.46	0.46
1:B:154:ARG:NH2	1:B:248:ASP:OD1	2.48	0.46
1:C:183:GLY:HA2	1:C:372:TYR:HB3	1.98	0.46
1:B:183:GLY:HA2	1:B:372:TYR:HB3	1.98	0.46
1:B:196:GLU:OE1	1:B:660:ARG:NH2	2.41	0.45
1:C:372:TYR:CE1	1:C:446:ARG:HB2	2.51	0.45
1:B:321:MET:HE3	1:B:322:TYR:CZ	2.52	0.45
1:A:154:ARG:HB2	1:A:666:THR:HA	1.99	0.45
1:B:123:ARG:HD2	1:B:427:ALA:HB1	2.00	0.44
1:B:150:LEU:O	1:B:162:GLY:HA3	2.17	0.44
1:B:633:LEU:HD21	1:B:635:LEU:HD21	2.00	0.44
1:B:321:MET:HE3	1:B:322:TYR:CE2	2.53	0.44
1:B:321:MET:SD	1:B:329:GLU:HG2	2.58	0.44
1:C:454:LEU:HD22	1:C:505:ILE:CG1	2.48	0.44
1:C:163:ASP:HB2	1:C:213:TYR:O	2.18	0.43
1:A:18:ASN:C	1:A:18:ASN:OD1	2.61	0.43
1:A:201:TYR:C	1:A:203:PRO:HD3	2.43	0.43
1:A:283:GLN:HE21	1:A:287:GLU:CD	2.26	0.43
1:C:188:HIS:HB2	5:C:903:HOH:O	2.17	0.43
1:A:352:TYR:CD1	1:A:352:TYR:C	2.97	0.43
1:B:679:LEU:O	1:B:683:LEU:HG	2.18	0.43
1:C:453:MET:HE3	1:C:478:LEU:HB3	1.98	0.43
1:A:315:TRP:N	1:A:316:PRO:CD	2.82	0.43
1:C:321:MET:HE3	1:C:322:TYR:CE2	2.54	0.43
1:B:694:HIS:CG	1:B:695:HIS:H	2.36	0.43
1:A:315:TRP:CG	1:A:316:PRO:HD3	2.54	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:313:TRP:CZ3	1:C:382:GLY:HA3	2.54	0.42
1:B:392:LEU:HG	1:B:418:MET:HG3	2.01	0.42
5:A:915:HOH:O	1:B:685:ARG:HD3	2.18	0.42
1:B:284:PHE:O	1:B:288:GLN:HG2	2.19	0.42
1:B:453:MET:HE3	1:B:478:LEU:HB3	2.01	0.42
1:A:454:LEU:HD22	1:A:505:ILE:HG13	2.02	0.42
1:C:527:GLU:O	1:C:528:LYS:HB2	2.20	0.41
1:A:163:ASP:HB2	1:A:213:TYR:O	2.20	0.41
1:A:187:ILE:HD12	1:A:209:LEU:HD22	2.03	0.41
1:C:623:ARG:HG2	1:C:686:ARG:HG3	2.02	0.41
1:C:565:LYS:HD3	1:C:576:LEU:CD2	2.50	0.41
1:B:454:LEU:HA	1:B:475:VAL:HB	2.02	0.41
1:C:133:GLU:OE2	1:C:515:TYR:OH	2.26	0.40
1:A:183:GLY:HA2	1:A:372:TYR:HB3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	678/696 (97%)	660 (97%)	18 (3%)	0	100	100
1	B	682/696 (98%)	666 (98%)	16 (2%)	0	100	100
1	C	677/696 (97%)	662 (98%)	15 (2%)	0	100	100
All	All	2037/2088 (98%)	1988 (98%)	49 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	570/582 (98%)	567 (100%)	3 (0%)	81	88
1	B	574/582 (99%)	572 (100%)	2 (0%)	86	91
1	C	569/582 (98%)	564 (99%)	5 (1%)	70	78
All	All	1713/1746 (98%)	1703 (99%)	10 (1%)	78	86

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	148	VAL
1	A	239	GLN
1	A	581	GLN
1	B	117	ASP
1	B	377	VAL
1	C	73	SER
1	C	148	VAL
1	C	233	GLN
1	C	342	ASP
1	C	689	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	295	GLN
1	A	337	HIS
1	A	606	HIS
1	A	648	ASN
1	B	42	HIS
1	B	524	ASN
1	B	622	GLN
1	C	42	HIS
1	C	239	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLC	D	1	2	12,12,12	0.59	0	17,17,17	0.94	1 (5%)
2	GLC	D	2	2	11,11,12	0.61	0	15,15,17	1.13	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	D	1	2	-	0/2/22/22	0/1/1/1
2	GLC	D	2	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	1	GLC	C4-C3-C2	-2.26	106.86	110.83
2	D	2	GLC	C1-O5-C5	2.19	115.12	112.19


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry ⓘ

Of 13 ligands modelled in this entry, 8 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	EDO	C	701	-	3,3,3	0.44	0	2,2,2	0.37	0
3	EDO	A	702	-	3,3,3	0.47	0	2,2,2	0.28	0
3	EDO	B	701	-	3,3,3	0.29	0	2,2,2	0.31	0
3	EDO	B	702	-	3,3,3	0.43	0	2,2,2	0.46	0
3	EDO	A	701	-	3,3,3	0.55	0	2,2,2	0.13	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	C	701	-	-	1/1/1/1	-
3	EDO	A	702	-	-	1/1/1/1	-
3	EDO	B	701	-	-	1/1/1/1	-
3	EDO	B	702	-	-	1/1/1/1	-
3	EDO	A	701	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	701	EDO	O1-C1-C2-O2
3	B	702	EDO	O1-C1-C2-O2
3	A	701	EDO	O1-C1-C2-O2
3	A	702	EDO	O1-C1-C2-O2
3	C	701	EDO	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	701	EDO	3	0
3	B	701	EDO	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	682/696 (97%)	0.34	24 (3%)	47 49	24, 36, 62, 81	0
1	B	686/696 (98%)	0.34	20 (2%)	53 56	27, 37, 58, 87	0
1	C	681/696 (97%)	0.36	24 (3%)	47 49	25, 37, 63, 98	0
All	All	2049/2088 (98%)	0.34	68 (3%)	49 51	24, 37, 62, 98	0

All (68) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	405	ILE	4.4
1	B	22	ALA	4.4
1	A	120	ALA	3.7
1	C	3	SER	3.4
1	A	574	VAL	3.4
1	A	691	HIS	3.4
1	B	694	HIS	3.2
1	C	117	ASP	3.2
1	A	406	LEU	3.1
1	C	52	THR	3.1
1	A	577	ARG	3.1
1	A	409	LEU	3.0
1	C	689	SER	2.9
1	B	117	ASP	2.9
1	C	406	LEU	2.8
1	A	5	ARG	2.8
1	A	324	ASN	2.8
1	A	403	PRO	2.8
1	A	404	ASP	2.7
1	B	406	LEU	2.7
1	C	23	HIS	2.7
1	C	405	ILE	2.7
1	C	42	HIS	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	313	TRP	2.6
1	B	313	TRP	2.6
1	B	20	ILE	2.5
1	B	311	MET	2.5
1	C	4	LYS	2.5
1	B	695	HIS	2.5
1	A	575	VAL	2.4
1	B	16	SER	2.4
1	B	25	LYS	2.4
1	C	22	ALA	2.4
1	C	31	ALA	2.4
1	C	2	GLU	2.4
1	B	24	GLY	2.4
1	B	405	ILE	2.4
1	A	52	THR	2.4
1	A	234	LEU	2.3
1	C	72	GLY	2.3
1	A	311	MET	2.3
1	C	690	ALA	2.3
1	B	409	LEU	2.3
1	C	116	ASP	2.3
1	A	3	SER	2.3
1	B	603	ARG	2.2
1	B	74	GLY	2.2
1	C	74	GLY	2.2
1	C	461	TYR	2.2
1	C	119	ARG	2.2
1	A	114	THR	2.2
1	A	2	GLU	2.2
1	B	97	ALA	2.2
1	A	365	TYR	2.1
1	C	43	GLN	2.1
1	B	73	SER	2.1
1	A	582	ASP	2.1
1	A	229	GLN	2.1
1	A	571	PRO	2.1
1	B	19	TYR	2.1
1	A	70	VAL	2.1
1	C	409	LEU	2.0
1	A	53	PRO	2.0
1	C	114	THR	2.0
1	B	85	GLY	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	399	VAL	2.0
1	C	73	SER	2.0
1	B	72	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GLC	D	2	11/12	0.72	0.18	50,56,66,66	0
2	GLC	D	1	12/12	0.87	0.12	45,51,55,56	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	EDO	C	701	4/4	0.86	0.19	29,30,30,31	0
3	EDO	A	701	4/4	0.89	0.10	39,39,39,41	0
3	EDO	B	702	4/4	0.92	0.10	35,35,35,37	0
3	EDO	B	701	4/4	0.93	0.08	31,33,36,40	0
4	IOD	C	702	1/1	0.93	0.10	69,69,69,69	0
4	IOD	B	705	1/1	0.94	0.23	69,69,69,69	0
3	EDO	A	702	4/4	0.94	0.07	26,28,28,29	0
4	IOD	C	703	1/1	0.95	0.19	83,83,83,83	0
4	IOD	B	703	1/1	0.96	0.12	52,52,52,52	0
4	IOD	B	704	1/1	0.96	0.21	76,76,76,76	0
4	IOD	A	704	1/1	0.96	0.07	46,46,46,46	0
4	IOD	A	705	1/1	0.96	0.16	69,69,69,69	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	IOD	A	706	1/1	0.96	0.21	75,75,75,75	0

6.5 Other polymers [i](#)

There are no such residues in this entry.