



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 5, 2026 – 09:34 AM EDT

PDB ID : 7SB8 / pdb_00007sb8
Title : d(GA(CGA)5) parallel-stranded homo-duplex
Authors : Luteran, E.M.; Paukstelis, P.J.
Deposited on : 2021-09-24
Resolution : 1.32 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.015 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	E	83	Total 86	O 86	0	3
5	F	77	Total 79	O 79	0	2
5	C	83	Total 87	O 87	0	4

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: GA(CG A)5

Chain A:  100%

There are no outlier residues recorded for this chain.


- Molecule 1: GA(CG A)5

Chain B:  94% 6%



- Molecule 1: GA(CG A)5

Chain D:  94% 6%



- Molecule 1: GA(CG A)5

Chain E:  100%

There are no outlier residues recorded for this chain.

- Molecule 1: GA(CG A)5

Chain F:  88% 12%



- Molecule 1: GA(CG A)5

Chain C:  100%

There are no outlier residues recorded for this chain.

[illegible]

Xtriage’s analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 66.87 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.6442e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*



WORLD WIDE
PDB
PROTEIN DATA BANK

5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 35 ligands modelled in this entry, 24 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NCO	B	104[B]	-	6,6,6	0.60	0	-		
4	NCO	D	105[B]	-	6,6,6	0.55	0	-		
4	NCO	B	105[A]	-	6,6,6	0.69	0	-		
4	NCO	B	104[A]	-	6,6,6	0.48	0	-		
4	NCO	F	103[A]	-	6,6,6	0.59	0	-		
4	NCO	D	103[A]	-	6,6,6	0.53	0	-		
4	NCO	D	105[A]	-	6,6,6	0.50	0	-		
4	NCO	D	104	-	6,6,6	0.65	0	-		
4	NCO	B	105[B]	-	6,6,6	0.52	0	-		
4	NCO	F	103[B]	-	6,6,6	0.55	0	-		
4	NCO	D	103[B]	-	6,6,6	0.51	0	-		

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	104[A]	NCO	1	0
4	F	103[A]	NCO	1	0
4	D	105[A]	NCO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

