



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 12, 2026 – 06:08 AM UTC

PDB ID : 6TBI / pdb_00006tbi
Title : Structure of a beta galactosidase with inhibitor
Authors : Offen, W.; Davies, G.
Deposited on : 2019-11-01
Resolution : 1.46 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.46 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1756 (1.46-1.46)
Clashscore	190562	1795 (1.46-1.46)
Ramachandran outliers	187476	1776 (1.46-1.46)
Sidechain outliers	187428	1776 (1.46-1.46)
RSRZ outliers	180081	1756 (1.46-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	550	
1	B	550	
1	C	550	
1	D	550	
1	E	550	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	550	2% 92% 5%
1	G	550	3% 91% 7%
1	H	550	3% 90% 8%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 40015 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-galactosidase, putative, bgl35A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	540	Total	C	N	O	S	0	8	0
			4289	2745	725	803	16			
1	B	534	Total	C	N	O	S	0	9	0
			4266	2729	725	795	17			
1	C	536	Total	C	N	O	S	0	10	0
			4313	2757	741	799	16			
1	D	539	Total	C	N	O	S	0	11	0
			4317	2761	730	810	16			
1	E	538	Total	C	N	O	S	0	9	0
			4305	2758	727	803	17			
1	F	539	Total	C	N	O	S	0	20	0
			4406	2814	754	822	16			
1	G	540	Total	C	N	O	S	0	17	0
			4382	2800	745	820	17			
1	H	538	Total	C	N	O	S	0	16	0
			4356	2785	742	812	17			

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	26	MET	-	initiating methionine	UNP B3PBE0
A	27	GLY	-	expression tag	UNP B3PBE0
A	28	SER	-	expression tag	UNP B3PBE0
A	29	SER	-	expression tag	UNP B3PBE0
A	30	HIS	-	expression tag	UNP B3PBE0
A	31	HIS	-	expression tag	UNP B3PBE0
A	32	HIS	-	expression tag	UNP B3PBE0
A	33	HIS	-	expression tag	UNP B3PBE0
A	34	HIS	-	expression tag	UNP B3PBE0
A	35	HIS	-	expression tag	UNP B3PBE0
B	26	MET	-	initiating methionine	UNP B3PBE0
B	27	GLY	-	expression tag	UNP B3PBE0
B	28	SER	-	expression tag	UNP B3PBE0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	29	SER	-	expression tag	UNP B3PBE0
B	30	HIS	-	expression tag	UNP B3PBE0
B	31	HIS	-	expression tag	UNP B3PBE0
B	32	HIS	-	expression tag	UNP B3PBE0
B	33	HIS	-	expression tag	UNP B3PBE0
B	34	HIS	-	expression tag	UNP B3PBE0
B	35	HIS	-	expression tag	UNP B3PBE0
C	26	MET	-	initiating methionine	UNP B3PBE0
C	27	GLY	-	expression tag	UNP B3PBE0
C	28	SER	-	expression tag	UNP B3PBE0
C	29	SER	-	expression tag	UNP B3PBE0
C	30	HIS	-	expression tag	UNP B3PBE0
C	31	HIS	-	expression tag	UNP B3PBE0
C	32	HIS	-	expression tag	UNP B3PBE0
C	33	HIS	-	expression tag	UNP B3PBE0
C	34	HIS	-	expression tag	UNP B3PBE0
C	35	HIS	-	expression tag	UNP B3PBE0
D	26	MET	-	initiating methionine	UNP B3PBE0
D	27	GLY	-	expression tag	UNP B3PBE0
D	28	SER	-	expression tag	UNP B3PBE0
D	29	SER	-	expression tag	UNP B3PBE0
D	30	HIS	-	expression tag	UNP B3PBE0
D	31	HIS	-	expression tag	UNP B3PBE0
D	32	HIS	-	expression tag	UNP B3PBE0
D	33	HIS	-	expression tag	UNP B3PBE0
D	34	HIS	-	expression tag	UNP B3PBE0
D	35	HIS	-	expression tag	UNP B3PBE0
E	26	MET	-	initiating methionine	UNP B3PBE0
E	27	GLY	-	expression tag	UNP B3PBE0
E	28	SER	-	expression tag	UNP B3PBE0
E	29	SER	-	expression tag	UNP B3PBE0
E	30	HIS	-	expression tag	UNP B3PBE0
E	31	HIS	-	expression tag	UNP B3PBE0
E	32	HIS	-	expression tag	UNP B3PBE0
E	33	HIS	-	expression tag	UNP B3PBE0
E	34	HIS	-	expression tag	UNP B3PBE0
E	35	HIS	-	expression tag	UNP B3PBE0
F	26	MET	-	initiating methionine	UNP B3PBE0
F	27	GLY	-	expression tag	UNP B3PBE0
F	28	SER	-	expression tag	UNP B3PBE0
F	29	SER	-	expression tag	UNP B3PBE0
F	30	HIS	-	expression tag	UNP B3PBE0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	31	HIS	-	expression tag	UNP B3PBE0
F	32	HIS	-	expression tag	UNP B3PBE0
F	33	HIS	-	expression tag	UNP B3PBE0
F	34	HIS	-	expression tag	UNP B3PBE0
F	35	HIS	-	expression tag	UNP B3PBE0
G	26	MET	-	initiating methionine	UNP B3PBE0
G	27	GLY	-	expression tag	UNP B3PBE0
G	28	SER	-	expression tag	UNP B3PBE0
G	29	SER	-	expression tag	UNP B3PBE0
G	30	HIS	-	expression tag	UNP B3PBE0
G	31	HIS	-	expression tag	UNP B3PBE0
G	32	HIS	-	expression tag	UNP B3PBE0
G	33	HIS	-	expression tag	UNP B3PBE0
G	34	HIS	-	expression tag	UNP B3PBE0
G	35	HIS	-	expression tag	UNP B3PBE0
H	26	MET	-	initiating methionine	UNP B3PBE0
H	27	GLY	-	expression tag	UNP B3PBE0
H	28	SER	-	expression tag	UNP B3PBE0
H	29	SER	-	expression tag	UNP B3PBE0
H	30	HIS	-	expression tag	UNP B3PBE0
H	31	HIS	-	expression tag	UNP B3PBE0
H	32	HIS	-	expression tag	UNP B3PBE0
H	33	HIS	-	expression tag	UNP B3PBE0
H	34	HIS	-	expression tag	UNP B3PBE0
H	35	HIS	-	expression tag	UNP B3PBE0

- Molecule 2 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	4	Total Na 4 4	0	0
2	B	4	Total Na 4 4	0	0
2	C	4	Total Na 4 4	0	0
2	D	6	Total Na 6 6	0	0
2	E	6	Total Na 6 6	0	0
2	F	4	Total Na 4 4	0	0
2	G	5	Total Na 5 5	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	H	5	Total	Na	0	0
			5	5		

- Molecule 3 is ACETATE ION (CCD ID: ACT) (formula: $C_2H_3O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	C	1	Total	C	O	0	0
			4	2	2		
3	D	1	Total	C	O	0	0
			4	2	2		
3	D	1	Total	C	O	0	0
			4	2	2		
3	E	1	Total	C	O	0	0
			4	2	2		
3	E	1	Total	C	O	0	0
			4	2	2		
3	E	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	F	1	Total C O 4 2 2	0	0
3	F	1	Total C O 4 2 2	0	0
3	G	1	Total C O 4 2 2	0	0
3	G	1	Total C O 4 2 2	0	0
3	G	1	Total C O 4 2 2	0	0
3	G	1	Total C O 4 2 2	0	0
3	H	1	Total C O 4 2 2	0	0

- Molecule 4 is (1S,2S,3S,6R)-4-(hydroxymethyl)-6-(octylamino)cyclohex-4-ene-1,2,3-triol (CCD ID: N8V) (formula: C₁₅H₂₉NO₄) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C N O 20 15 1 4	0	0
4	B	1	Total C N O 20 15 1 4	0	0
4	C	1	Total C N O 16 11 1 4	0	0
4	D	1	Total C N O 20 15 1 4	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	E	1	Total	C	N	O	0	0
			14	9	1	4		
4	F	1	Total	C	N	O	0	0
			20	15	1	4		
4	G	1	Total	C	N	O	0	0
			20	15	1	4		
4	H	1	Total	C	N	O	0	0
			20	15	1	4		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	504	Total	O	0	1
			505	505		
5	B	590	Total	O	0	0
			590	590		
5	C	606	Total	O	0	2
			608	608		
5	D	709	Total	O	0	4
			713	713		
5	E	715	Total	O	0	5
			720	720		
5	F	669	Total	O	0	3
			672	672		
5	G	719	Total	O	0	2
			721	721		
5	H	595	Total	O	0	1
			596	596		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Beta-galactosidase, putative, bgl35A



- Molecule 1: Beta-galactosidase, putative, bgl35A




- Molecule 1: Beta-galactosidase, putative, bgl35A



- Molecule 1: Beta-galactosidase, putative, bgl35A





4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	99.12Å 115.91Å 116.10Å 90.20° 89.96° 90.08°	Depositor
Resolution (Å)	116.10 – 1.46 116.10 – 1.46	Depositor EDS
% Data completeness (in resolution range)	96.3 (116.10-1.46) 96.4 (116.10-1.46)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.98 (at 1.46Å)	Xtriage
Refinement program	REFMAC 5.8.0253	Depositor
R, R_{free}	0.126 , 0.172 0.135 , 0.176	Depositor DCC
R_{free} test set	43376 reflections (4.62%)	wwPDB-VP
Wilson B-factor (Å ²)	19.8	Xtriage
Anisotropy	0.219	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 61.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.000 for h,-l,k 0.000 for h,l,-k 0.000 for h,-k,-l 0.000 for -h,k,-l 0.000 for -h,-k,l 0.000 for -h,-l,-k 0.000 for -h,l,k	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	40015	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, N8V, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.21	8/4403 (0.2%)	1.29	6/6002 (0.1%)
1	B	1.19	9/4378 (0.2%)	1.30	13/5965 (0.2%)
1	C	1.17	8/4428 (0.2%)	1.28	7/6026 (0.1%)
1	D	1.22	7/4433 (0.2%)	1.31	10/6039 (0.2%)
1	E	1.24	11/4418 (0.2%)	1.34	19/6014 (0.3%)
1	F	1.21	4/4523 (0.1%)	1.31	12/6153 (0.2%)
1	G	1.22	8/4495 (0.2%)	1.31	14/6116 (0.2%)
1	H	1.21	8/4475 (0.2%)	1.27	10/6094 (0.2%)
All	All	1.21	63/35553 (0.2%)	1.30	91/48409 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	F	0	1

The worst 5 of 63 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	34	HIS	CE1-NE2	9.06	1.41	1.32
1	H	326	ARG	C-O	7.83	1.34	1.24
1	A	139	HIS	CE1-NE2	7.78	1.40	1.32
1	F	525	LYS	CG-CD	-7.57	1.29	1.52
1	G	255	HIS	CE1-NE2	7.46	1.40	1.32

The worst 5 of 91 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	525	LYS	CB-CG-CD	7.56	128.69	111.30
1	B	238	LYS	CB-CA-C	7.51	120.10	108.61
1	E	159	ASP	CA-CB-CG	7.42	120.02	112.60
1	G	537	GLU	CB-CG-CD	7.30	125.01	112.60
1	H	227	VAL	N-CA-C	-6.96	102.55	109.02

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	F	71	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4289	0	4100	10	0
1	B	4266	0	4088	13	0
1	C	4313	0	4151	15	0
1	D	4317	0	4139	14	0
1	E	4305	0	4150	12	0
1	F	4406	0	4232	13	0
1	G	4382	0	4206	19	0
1	H	4356	0	4161	21	0
2	A	4	0	0	0	0
2	B	4	0	0	0	0
2	C	4	0	0	0	0
2	D	6	0	0	0	0
2	E	6	0	0	0	0
2	F	4	0	0	0	0
2	G	5	0	0	0	0
2	H	5	0	0	0	0
3	A	4	0	3	0	0
3	B	8	0	6	0	0
3	C	4	0	3	0	0
3	D	8	0	6	0	0
3	E	16	0	12	0	0
3	F	8	0	6	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	G	16	0	12	1	0
3	H	4	0	3	0	0
4	A	20	0	29	0	0
4	B	20	0	29	0	0
4	C	16	0	18	0	0
4	D	20	0	29	1	0
4	E	14	0	14	0	0
4	F	20	0	29	0	0
4	G	20	0	29	1	0
4	H	20	0	29	0	0
5	A	505	0	0	2	0
5	B	590	0	0	4	0
5	C	608	0	0	6	0
5	D	713	0	0	12	0
5	E	720	0	0	6	0
5	F	672	0	0	4	0
5	G	721	0	0	9	0
5	H	596	0	0	12	0
All	All	40015	0	33484	113	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 113 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:503:GLU:OE2	1:F:566[A]:ARG:HD3	1.46	1.10
1:G:421:GLN:HA	5:G:703:HOH:O	1.52	1.09
1:C:452:LYS:HE2	5:C:1086:HOH:O	1.56	1.06
1:B:105:GLN:HG3	5:B:1230:HOH:O	1.77	0.85
1:F:503:GLU:OE2	1:F:566[A]:ARG:CD	2.25	0.84

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	546/550 (99%)	530 (97%)	15 (3%)	1 (0%)	43	21
1	B	539/550 (98%)	519 (96%)	19 (4%)	1 (0%)	43	21
1	C	542/550 (98%)	520 (96%)	22 (4%)	0	100	100
1	D	546/550 (99%)	527 (96%)	19 (4%)	0	100	100
1	E	543/550 (99%)	527 (97%)	15 (3%)	1 (0%)	43	21
1	F	558/550 (102%)	537 (96%)	20 (4%)	1 (0%)	43	21
1	G	555/550 (101%)	533 (96%)	22 (4%)	0	100	100
1	H	550/550 (100%)	530 (96%)	18 (3%)	2 (0%)	30	11
All	All	4379/4400 (100%)	4223 (96%)	150 (3%)	6 (0%)	48	22

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	491	ALA
1	H	491	ALA
1	B	491	ALA
1	E	491	ALA
1	F	491	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	438/461 (95%)	436 (100%)	2 (0%)	81	64
1	B	439/461 (95%)	436 (99%)	3 (1%)	76	54
1	C	445/461 (96%)	442 (99%)	3 (1%)	76	54
1	D	445/461 (96%)	443 (100%)	2 (0%)	84	70
1	E	445/461 (96%)	442 (99%)	3 (1%)	76	54
1	F	453/461 (98%)	446 (98%)	7 (2%)	57	25

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	450/461 (98%)	446 (99%)	4 (1%)	70	46
1	H	448/461 (97%)	446 (100%)	2 (0%)	84	70
All	All	3563/3688 (97%)	3537 (99%)	26 (1%)	78	54

5 of 26 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	169[A]	LEU
1	F	537	GLU
1	H	136	ASN
1	F	525	LYS
1	F	566[A]	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 22 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	451	HIS
1	G	64	GLN
1	F	498	GLN
1	G	275	ASN
1	C	233	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 63 ligands modelled in this entry, 38 are monoatomic - leaving 25 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ACT	E	605	-	3,3,3	1.07	0	3,3,3	0.74	0
4	N8V	A	606	-	20,20,20	1.35	3 (15%)	18,25,25	1.33	1 (5%)
3	ACT	F	604	-	3,3,3	1.24	0	3,3,3	0.95	0
3	ACT	A	602	-	3,3,3	1.37	0	3,3,3	0.58	0
3	ACT	H	602	-	3,3,3	1.46	1 (33%)	3,3,3	0.92	0
4	N8V	G	610	-	20,20,20	1.68	3 (15%)	18,25,25	1.74	3 (16%)
3	ACT	G	603	-	3,3,3	1.38	0	3,3,3	0.59	0
4	N8V	D	609	-	20,20,20	1.87	4 (20%)	18,25,25	1.54	3 (16%)
4	N8V	F	607	-	20,20,20	2.14	3 (15%)	18,25,25	1.58	3 (16%)
3	ACT	E	604	-	3,3,3	1.40	0	3,3,3	0.93	0
3	ACT	G	606	-	3,3,3	1.09	0	3,3,3	1.13	0
3	ACT	E	607	-	3,3,3	1.38	0	3,3,3	0.62	0
3	ACT	G	604	-	3,3,3	0.62	0	3,3,3	1.20	0
3	ACT	D	605	-	3,3,3	0.86	0	3,3,3	1.02	0
3	ACT	F	605	-	3,3,3	0.81	0	3,3,3	0.56	0
3	ACT	G	605	-	3,3,3	1.24	0	3,3,3	0.74	0
3	ACT	B	603	-	3,3,3	0.83	0	3,3,3	1.11	0
4	N8V	B	607	-	20,20,20	1.84	5 (25%)	18,25,25	1.37	2 (11%)
4	N8V	H	607	-	20,20,20	1.49	5 (25%)	18,25,25	1.31	3 (16%)
4	N8V	E	611	-	14,14,20	2.32	3 (21%)	12,19,25	1.42	2 (16%)
3	ACT	B	602	-	3,3,3	0.67	0	3,3,3	1.15	0
3	ACT	D	604	-	3,3,3	1.13	0	3,3,3	1.19	0
3	ACT	E	606	-	3,3,3	1.14	0	3,3,3	0.81	0
4	N8V	C	606	-	16,16,20	1.65	3 (18%)	14,21,25	1.92	2 (14%)
3	ACT	C	603	-	3,3,3	1.38	0	3,3,3	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	N8V	H	607	-	-	1/11/31/31	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	N8V	E	611	-	-	0/5/25/31	0/1/1/1
4	N8V	A	606	-	-	1/11/31/31	0/1/1/1
4	N8V	D	609	-	-	0/11/31/31	0/1/1/1
4	N8V	F	607	-	-	2/11/31/31	0/1/1/1
4	N8V	C	606	-	-	0/7/27/31	0/1/1/1
4	N8V	G	610	-	-	3/11/31/31	0/1/1/1
4	N8V	B	607	-	-	1/11/31/31	0/1/1/1

The worst 5 of 30 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	F	607	N8V	C4-C5	-7.86	1.45	1.51
4	E	611	N8V	C4-C5	-5.94	1.46	1.51
4	D	609	N8V	C4-C5	-5.88	1.46	1.51
4	B	607	N8V	C4-C5	-4.96	1.47	1.51
4	C	606	N8V	C2-C1	4.61	1.58	1.53

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	606	N8V	C3-C4-C5	-5.96	102.30	111.67
4	D	609	N8V	C3-C4-C5	-4.97	103.86	111.67
4	G	610	N8V	C3-C4-C5	-4.95	103.89	111.67
4	F	607	N8V	C7-C1-N8	-4.27	104.63	110.71
4	E	611	N8V	C7-C1-N8	-3.69	105.46	110.71

There are no chirality outliers.

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	F	607	N8V	C11-C10-C9-N8
4	G	610	N8V	C11-C10-C9-N8
4	A	606	N8V	C11-C12-C13-C14
4	G	610	N8V	C9-C10-C11-C12
4	F	607	N8V	C9-C10-C11-C12

There are no ring outliers.

3 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	G	610	N8V	1	0


Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	609	N8V	1	0
3	G	604	ACT	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.
















Ligand N8V E 611





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	540/550 (98%)	0.01	23 (4%)	40	42	10, 24, 40, 60	50 (9%)
1	B	534/550 (97%)	-0.21	11 (2%)	63	66	10, 22, 36, 58	30 (5%)
1	C	536/550 (97%)	-0.19	13 (2%)	59	62	9, 22, 36, 65	35 (6%)
1	D	539/550 (98%)	-0.35	13 (2%)	59	62	9, 19, 33, 65	35 (6%)
1	E	538/550 (97%)	-0.41	7 (1%)	75	78	9, 19, 34, 49	33 (6%)
1	F	539/550 (98%)	-0.28	10 (1%)	66	68	7, 20, 39, 72	46 (8%)
1	G	540/550 (98%)	-0.32	16 (2%)	52	55	8, 19, 35, 63	43 (7%)
1	H	538/550 (97%)	-0.10	16 (2%)	52	55	10, 22, 35, 63	56 (10%)
All	All	4304/4400 (97%)	-0.23	109 (2%)	58	60	7, 21, 37, 72	328 (7%)

The worst 5 of 109 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	438	ILE	12.1
1	A	438	ILE	11.7
1	G	438	ILE	11.2
1	F	438	ILE	10.0
1	B	439	TRP	9.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	ACT	E	607	4/4	0.86	0.19	20,24,25,26	4
3	ACT	F	604	4/4	0.86	0.16	30,32,32,33	4
3	ACT	F	605	4/4	0.88	0.15	22,27,27,28	4
3	ACT	G	604	4/4	0.88	0.21	43,47,48,53	4
3	ACT	D	604	4/4	0.90	0.13	31,32,38,40	0
3	ACT	H	602	4/4	0.90	0.12	26,27,29,31	4
3	ACT	G	606	4/4	0.91	0.10	31,31,31,34	4
3	ACT	E	604	4/4	0.92	0.14	20,21,22,24	4
3	ACT	G	605	4/4	0.92	0.12	30,34,35,36	4
3	ACT	E	605	4/4	0.93	0.12	31,32,34,37	4
3	ACT	C	603	4/4	0.93	0.19	18,20,21,22	4
3	ACT	D	605	4/4	0.94	0.10	25,25,40,46	4
3	ACT	E	606	4/4	0.94	0.10	29,43,45,47	0
3	ACT	A	602	4/4	0.94	0.09	27,29,29,34	4
3	ACT	G	603	4/4	0.95	0.17	21,22,26,27	4
3	ACT	B	602	4/4	0.96	0.07	32,32,33,34	4
3	ACT	B	603	4/4	0.96	0.10	29,31,33,36	4
2	NA	C	602	1/1	0.97	0.06	35,35,35,35	0
4	N8V	D	609	20/20	0.97	0.09	14,17,39,40	6
2	NA	B	605	1/1	0.98	0.07	42,42,42,42	0
2	NA	H	605	1/1	0.98	0.04	35,35,35,35	1
2	NA	B	606	1/1	0.98	0.08	35,35,35,35	0
4	N8V	B	607	20/20	0.98	0.08	16,19,35,36	6
4	N8V	C	606	16/20	0.98	0.08	16,18,35,40	3
2	NA	C	601	1/1	0.98	0.06	43,43,43,43	0
4	N8V	E	611	14/20	0.98	0.05	15,16,29,35	0
4	N8V	F	607	20/20	0.98	0.06	14,17,38,41	5
4	N8V	G	610	20/20	0.98	0.07	13,16,31,32	6
4	N8V	H	607	20/20	0.98	0.07	15,20,48,51	5
2	NA	H	606	1/1	0.99	0.04	39,39,39,39	0
2	NA	A	604	1/1	0.99	0.10	36,36,36,36	0
2	NA	A	605	1/1	0.99	0.06	31,31,31,31	0
2	NA	B	604	1/1	0.99	0.08	25,25,25,25	0
2	NA	C	604	1/1	0.99	0.10	25,25,25,25	0
2	NA	D	601	1/1	0.99	0.07	22,22,22,22	0
2	NA	D	606	1/1	0.99	0.12	23,23,23,23	0
2	NA	D	607	1/1	0.99	0.11	27,27,27,27	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NA	D	608	1/1	0.99	0.05	24,24,24,24	0
2	NA	E	603	1/1	0.99	0.03	22,22,22,22	0
2	NA	E	608	1/1	0.99	0.09	28,28,28,28	0
2	NA	E	609	1/1	0.99	0.04	40,40,40,40	0
2	NA	E	610	1/1	0.99	0.04	22,22,22,22	0
2	NA	F	601	1/1	0.99	0.06	25,25,25,25	0
2	NA	F	602	1/1	0.99	0.04	25,25,25,25	0
2	NA	F	603	1/1	0.99	0.03	20,20,20,20	0
2	NA	G	601	1/1	0.99	0.07	25,25,25,25	0
2	NA	G	602	1/1	0.99	0.05	20,20,20,20	0
4	N8V	A	606	20/20	0.99	0.06	17,20,35,37	7
2	NA	G	607	1/1	0.99	0.04	27,27,27,27	0
2	NA	G	608	1/1	0.99	0.05	23,23,23,23	0
2	NA	G	609	1/1	0.99	0.03	25,25,25,25	0
2	NA	H	601	1/1	0.99	0.04	32,32,32,32	0
2	NA	H	603	1/1	0.99	0.09	24,24,24,24	0
2	NA	H	604	1/1	0.99	0.06	27,27,27,27	0
2	NA	A	601	1/1	0.99	0.06	32,32,32,32	0
2	NA	C	605	1/1	1.00	0.08	27,27,27,27	0
2	NA	A	603	1/1	1.00	0.08	25,25,25,25	0
2	NA	E	601	1/1	1.00	0.03	21,21,21,21	0
2	NA	E	602	1/1	1.00	0.03	20,20,20,20	0
2	NA	F	606	1/1	1.00	0.05	26,26,26,26	0
2	NA	D	602	1/1	1.00	0.04	22,22,22,22	0
2	NA	D	603	1/1	1.00	0.04	20,20,20,20	0
2	NA	B	601	1/1	1.00	0.10	32,32,32,32	0



The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around N8V D 609:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around N8V B 607:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around N8V C 606:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around N8V E 611:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around N8V F 607:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around N8V G 610:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around N8V H 607:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around N8V A 606:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.