



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 7, 2026 – 12:29 AM UTC

PDB ID : 1TNB / pdb_00001tnb
Title : Rat Protein Geranylgeranyltransferase Type-I Complexed with a GGPP analog and a substrate KKSKTKCVIF Peptide Derived from TC21
Authors : Reid, T.S.; Terry, K.L.; Casey, P.J.; Beese, L.S.
Deposited on : 2004-06-11
Resolution : 2.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1407 (2.88-2.84)
Clashscore	190562	1446 (2.88-2.84)
Ramachandran outliers	187476	1406 (2.88-2.84)
Sidechain outliers	187428	1407 (2.88-2.84)
RSRZ outliers	180081	1408 (2.88-2.84)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	377	2% 58% 23% • 17%
1	C	377	3% 62% 21% • 17%
1	E	377	2% 59% 23% • 17%
1	G	377	3% 62% 20% • 17%
1	I	377	4% 62% 20% • 17%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	K	377	
2	B	377	
2	D	377	
2	F	377	
2	H	377	
2	J	377	
2	L	377	
3	M	10	
3	N	10	
3	O	10	
3	P	10	
3	Q	10	
3	R	10	

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 33504 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called geranylgeranyltransferase type I alpha subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	314	Total	C	N	O	S	0	0	0
			2636	1682	463	486	5			
1	C	314	Total	C	N	O	S	0	0	0
			2655	1695	465	490	5			
1	E	314	Total	C	N	O	S	0	0	0
			2664	1698	466	495	5			
1	G	314	Total	C	N	O	S	0	0	0
			2651	1694	465	487	5			
1	I	314	Total	C	N	O	S	0	0	0
			2648	1691	461	491	5			
1	K	314	Total	C	N	O	S	0	0	0
			2675	1705	468	497	5			

- Molecule 2 is a protein called Geranylgeranyl transferase type I beta subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	346	Total	C	N	O	S	0	0	0
			2703	1709	468	502	24			
2	D	346	Total	C	N	O	S	0	0	0
			2706	1713	467	502	24			
2	F	346	Total	C	N	O	S	0	0	0
			2717	1716	473	504	24			
2	H	346	Total	C	N	O	S	0	0	0
			2694	1705	464	501	24			
2	J	346	Total	C	N	O	S	0	0	0
			2708	1711	471	502	24			
2	L	346	Total	C	N	O	S	0	0	0
			2719	1718	473	504	24			


- Molecule 3 is a protein called Fusion protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	M	5	Total	C	N	O	S	0	0	0
			42	29	6	6	1			
3	N	5	Total	C	N	O	S	0	0	0
			42	29	6	6	1			
3	O	5	Total	C	N	O	S	0	0	0
			42	29	6	6	1			
3	P	5	Total	C	N	O	S	0	0	0
			42	29	6	6	1			
3	Q	5	Total	C	N	O	S	0	0	0
			42	29	6	6	1			
3	R	5	Total	C	N	O	S	0	0	0
			42	29	6	6	1			

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Zn	0	0
			1	1		
4	D	1	Total	Zn	0	0
			1	1		
4	F	1	Total	Zn	0	0
			1	1		
4	H	1	Total	Zn	0	0
			1	1		
4	J	1	Total	Zn	0	0
			1	1		
4	L	1	Total	Zn	0	0
			1	1		

- Molecule 5 is 2-[METHYL-(5-GERANYL-4-METHYL-PENT-3-ENYL)-AMINO]-ETHYL-DIPHOSPHATE (CCD ID: MGM) (formula: C₁₉H₃₇NO₇P₂).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	B	1	Total	C	N	O	P	0	0
			29	19	1	7	2		
5	D	1	Total	C	N	O	P	0	0
			29	19	1	7	2		
5	F	1	Total	C	N	O	P	0	0
			29	19	1	7	2		
5	H	1	Total	C	N	O	P	0	0
			29	19	1	7	2		
5	J	1	Total	C	N	O	P	0	0
			29	19	1	7	2		
5	L	1	Total	C	N	O	P	0	0
			29	19	1	7	2		

- Molecule 6 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	D	1	Total	Cl	0	0
			1	1		
6	F	1	Total	Cl	0	0
			1	1		
6	H	1	Total	Cl	0	0
			1	1		

- Molecule 7 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	F	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 8 is water.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	56	Total	O	0	0
			56	56		
8	B	55	Total	O	0	0
			55	55		
8	C	64	Total	O	0	0
			64	64		
8	D	70	Total	O	0	0
			70	70		
8	E	60	Total	O	0	0
			60	60		
8	F	77	Total	O	0	0
			77	77		
8	G	57	Total	O	0	0
			57	57		
8	H	49	Total	O	0	0
			49	49		
8	I	62	Total	O	0	0
			62	62		
8	J	59	Total	O	0	0
			59	59		
8	K	128	Total	O	0	0
			128	128		


Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	L	110	Total 110	O 110	0	0
8	M	9	Total 9	O 9	0	0
8	N	7	Total 7	O 7	0	0
8	O	2	Total 2	O 2	0	0
8	P	4	Total 4	O 4	0	0
8	Q	6	Total 6	O 6	0	0
8	R	6	Total 6	O 6	0	0

- Molecule 1: geranylgeranyltransferase type I alpha subunit







- Molecule 1: geranylgeranyltransferase type I alpha subunit




- Molecule 1: geranylgeranyltransferase type I alpha subunit




- Molecule 1: geranylgeranyltransferase type I alpha subunit




• Molecule 2: Geranylgeranyl transferase type I beta subunit




• Molecule 2: Geranylgeranyl transferase type I beta subunit




• Molecule 2: Geranylgeranyl transferase type I beta subunit



- Molecule 3: Fusion protein




- Molecule 3: Fusion protein




- Molecule 3: Fusion protein



- Molecule 3: Fusion protein



- Molecule 3: Fusion protein



- Molecule 3: Fusion protein



LYS	LYS	LYS	SER	LYS	THR	K6	F10
-----	-----	-----	-----	-----	-----	----	-----

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	271.13Å 266.72Å 185.14Å 90.00° 131.68° 90.00°	Depositor
Resolution (Å)	27.66 – 2.85 27.66 – 2.85	Depositor EDS
% Data completeness (in resolution range)	93.6 (27.66-2.85) 93.6 (27.66-2.85)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.12 (at 2.85Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.189 , 0.210 0.188 , 0.209	Depositor DCC
R_{free} test set	10767 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	42.8	Xtriage
Anisotropy	0.049	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 60.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.087 for -h-2*1,-k,l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	33504	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MGM, CL, ZN, MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.41	0/2702	0.87	6/3677 (0.2%)
1	C	0.41	0/2721	0.86	6/3698 (0.2%)
1	E	0.42	0/2730	0.87	8/3710 (0.2%)
1	G	0.43	0/2717	0.88	7/3692 (0.2%)
1	I	0.42	0/2714	0.85	8/3690 (0.2%)
1	K	0.46	0/2741	0.87	8/3722 (0.2%)
2	B	0.41	0/2765	0.90	4/3741 (0.1%)
2	D	0.43	0/2768	0.90	9/3743 (0.2%)
2	F	0.44	0/2779	0.91	7/3757 (0.2%)
2	H	0.41	0/2755	0.90	6/3727 (0.2%)
2	J	0.41	0/2769	0.90	6/3744 (0.2%)
2	L	0.45	0/2781	0.91	5/3759 (0.1%)
3	M	0.55	0/42	0.75	0/53
3	N	0.57	0/42	0.74	0/53
3	O	0.59	0/42	0.84	0/53
3	P	0.61	0/42	0.78	0/53
3	Q	0.58	0/42	0.83	0/53
3	R	0.64	0/42	0.77	0/53
All	All	0.43	0/33194	0.89	80/44978 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	E	0	1
2	B	0	1
2	D	0	1
2	H	0	1
2	J	0	1

Continued on next page...

Continued from previous page...

Mol	Chain	#Chirality outliers	#Planarity outliers
2	L	0	1
All	All	0	6

There are no bond length outliers.

The worst 5 of 80 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	J	259	GLY	N-CA-C	-10.58	97.48	112.60
2	D	259	GLY	N-CA-C	-10.38	97.75	112.60
2	H	259	GLY	N-CA-C	-10.27	97.92	112.60
2	L	259	GLY	N-CA-C	-10.22	97.98	112.60
2	F	259	GLY	N-CA-C	-10.18	98.04	112.60

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	297	TYR	Sidechain
2	D	272	TYR	Sidechain
1	E	102	TYR	Sidechain
2	H	272	TYR	Sidechain
2	J	272	TYR	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2636	0	2526	79	0
1	C	2655	0	2562	58	0
1	E	2664	0	2570	70	0
1	G	2651	0	2563	59	0
1	I	2648	0	2547	52	0
1	K	2675	0	2594	57	0
2	B	2703	0	2606	58	0
2	D	2706	0	2616	70	0
2	F	2717	0	2635	52	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	H	2694	0	2589	81	0
2	J	2708	0	2613	73	0
2	L	2719	0	2639	57	0
3	M	42	0	45	0	0
3	N	42	0	45	0	0
3	O	42	0	45	0	0
3	P	42	0	45	0	0
3	Q	42	0	45	0	0
3	R	42	0	45	0	0
4	B	1	0	0	0	0
4	D	1	0	0	0	0
4	F	1	0	0	0	0
4	H	1	0	0	0	0
4	J	1	0	0	0	0
4	L	1	0	0	0	0
5	B	29	0	34	2	0
5	D	29	0	34	0	0
5	F	29	0	34	2	0
5	H	29	0	34	1	0
5	J	29	0	34	2	0
5	L	29	0	34	0	0
6	D	1	0	0	0	0
6	F	1	0	0	0	0
6	H	1	0	0	0	0
7	F	12	0	13	0	0
8	A	56	0	0	4	0
8	B	55	0	0	0	0
8	C	64	0	0	2	0
8	D	70	0	0	1	0
8	E	60	0	0	1	0
8	F	77	0	0	3	0
8	G	57	0	0	1	0
8	H	49	0	0	2	0
8	I	62	0	0	2	0
8	J	59	0	0	0	0
8	K	128	0	0	5	0
8	L	110	0	0	3	0
8	M	9	0	0	0	0
8	N	7	0	0	0	0
8	O	2	0	0	0	0
8	P	4	0	0	0	0
8	Q	6	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	R	6	0	0	0	0
All	All	33504	0	31547	731	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 731 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:156:ILE:HG12	1:A:172:ARG:HH12	1.12	1.14
1:G:156:ILE:HG12	1:G:172:ARG:HH12	1.15	1.08
1:K:156:ILE:HG12	1:K:172:ARG:HH12	0.98	1.08
1:I:156:ILE:HG12	1:I:172:ARG:HH12	1.12	1.07
1:E:156:ILE:HG12	1:E:172:ARG:HH12	1.17	1.04

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	312/377 (83%)	288 (92%)	24 (8%)	0	100	100
1	C	312/377 (83%)	290 (93%)	22 (7%)	0	100	100
1	E	312/377 (83%)	289 (93%)	23 (7%)	0	100	100
1	G	312/377 (83%)	289 (93%)	23 (7%)	0	100	100
1	I	312/377 (83%)	290 (93%)	22 (7%)	0	100	100
1	K	312/377 (83%)	293 (94%)	19 (6%)	0	100	100
2	B	344/377 (91%)	326 (95%)	16 (5%)	2 (1%)	21	38
2	D	344/377 (91%)	329 (96%)	12 (4%)	3 (1%)	14	28

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	F	344/377 (91%)	327 (95%)	16 (5%)	1 (0%)	36	54
2	H	344/377 (91%)	326 (95%)	17 (5%)	1 (0%)	36	54
2	J	344/377 (91%)	322 (94%)	20 (6%)	2 (1%)	21	38
2	L	344/377 (91%)	328 (95%)	14 (4%)	2 (1%)	21	38
3	M	3/10 (30%)	3 (100%)	0	0	100	100
3	N	3/10 (30%)	3 (100%)	0	0	100	100
3	O	3/10 (30%)	3 (100%)	0	0	100	100
3	P	3/10 (30%)	3 (100%)	0	0	100	100
3	Q	3/10 (30%)	3 (100%)	0	0	100	100
3	R	3/10 (30%)	3 (100%)	0	0	100	100
All	All	3954/4584 (86%)	3715 (94%)	228 (6%)	11 (0%)	36	54

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	38	GLU
2	B	258	ASN
2	D	258	ASN
2	F	258	ASN
2	H	258	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	282/338 (83%)	271 (96%)	11 (4%)	28	54
1	C	286/338 (85%)	277 (97%)	9 (3%)	35	60
1	E	289/338 (86%)	280 (97%)	9 (3%)	35	60
1	G	285/338 (84%)	279 (98%)	6 (2%)	47	70
1	I	285/338 (84%)	280 (98%)	5 (2%)	51	73
1	K	292/338 (86%)	287 (98%)	5 (2%)	53	75

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	291/326 (89%)	280 (96%)	11 (4%)	29	54
2	D	291/326 (89%)	276 (95%)	15 (5%)	21	43
2	F	295/326 (90%)	281 (95%)	14 (5%)	23	47
2	H	288/326 (88%)	277 (96%)	11 (4%)	29	54
2	J	291/326 (89%)	277 (95%)	14 (5%)	23	46
2	L	295/326 (90%)	281 (95%)	14 (5%)	23	47
3	M	5/10 (50%)	5 (100%)	0	100	100
3	N	5/10 (50%)	5 (100%)	0	100	100
3	O	5/10 (50%)	5 (100%)	0	100	100
3	P	5/10 (50%)	5 (100%)	0	100	100
3	Q	5/10 (50%)	5 (100%)	0	100	100
3	R	5/10 (50%)	5 (100%)	0	100	100
All	All	3500/4044 (86%)	3376 (96%)	124 (4%)	32	57

5 of 124 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	F	33	LEU
2	L	21	LEU
1	G	194	ASN
1	K	357	ARG
2	L	284	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 75 such sidechains are listed below:

Mol	Chain	Res	Type
1	I	367	HIS
1	K	326	GLN
2	J	34	GLN
1	K	81	ASN
2	D	208	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 9 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	MES	F	380	-	12,12,12	9.23	8 (66%)	15,16,16	2.15	5 (33%)
5	MGM	D	380	-	27,28,28	0.95	2 (7%)	34,37,37	2.03	6 (17%)
5	MGM	L	379	-	27,28,28	1.02	2 (7%)	34,37,37	2.04	6 (17%)
5	MGM	J	379	-	27,28,28	0.98	2 (7%)	34,37,37	2.02	6 (17%)
5	MGM	F	381	-	27,28,28	1.03	1 (3%)	34,37,37	2.09	6 (17%)
5	MGM	B	379	-	27,28,28	1.01	2 (7%)	34,37,37	2.02	7 (20%)
5	MGM	H	380	-	27,28,28	0.97	2 (7%)	34,37,37	2.00	6 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	MES	F	380	-	-	4/6/14/14	0/1/1/1
5	MGM	D	380	-	-	10/31/31/31	-
5	MGM	L	379	-	-	10/31/31/31	-
5	MGM	J	379	-	-	10/31/31/31	-
5	MGM	F	381	-	-	10/31/31/31	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	MGM	B	379	-	-	10/31/31/31	-
5	MGM	H	380	-	-	10/31/31/31	-

The worst 5 of 19 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	F	380	MES	C8-S	-24.24	1.43	1.77
7	F	380	MES	O2S-S	12.77	1.81	1.45
7	F	380	MES	O1S-S	12.13	1.79	1.45
7	F	380	MES	O3S-S	8.56	1.79	1.47
7	F	380	MES	C7-C8	-4.81	1.39	1.52

The worst 5 of 42 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	F	381	MGM	C1-C2-N3	7.49	131.28	113.30
5	B	379	MGM	C1-C2-N3	7.21	130.61	113.30
5	J	379	MGM	C1-C2-N3	7.20	130.58	113.30
5	L	379	MGM	C1-C2-N3	7.15	130.47	113.30
5	H	380	MGM	C1-C2-N3	7.09	130.33	113.30

There are no chirality outliers.

5 of 64 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	379	MGM	C1-C2-N3-C4
5	B	379	MGM	O1-C1-C2-N3
5	B	379	MGM	PA-O3A-PB-O1B
5	D	380	MGM	C1-C2-N3-C4
5	D	380	MGM	O1-C1-C2-N3


There are no ring outliers.


4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	J	379	MGM	2	0
5	F	381	MGM	2	0
5	B	379	MGM	2	0
5	H	380	MGM	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	314/377 (83%)	0.18	7 (2%) 62 53	31, 53, 86, 100	0
1	C	314/377 (83%)	0.09	10 (3%) 50 41	28, 49, 77, 98	0
1	E	314/377 (83%)	0.14	6 (1%) 66 59	28, 51, 77, 95	0
1	G	314/377 (83%)	0.14	12 (3%) 44 35	29, 49, 76, 98	0
1	I	314/377 (83%)	0.27	14 (4%) 38 29	27, 52, 82, 96	0
1	K	314/377 (83%)	-0.33	4 (1%) 75 68	21, 37, 63, 75	0
2	B	346/377 (91%)	-0.09	11 (3%) 50 41	30, 45, 67, 91	0
2	D	346/377 (91%)	-0.36	7 (2%) 65 57	26, 39, 62, 76	0
2	F	346/377 (91%)	-0.32	10 (2%) 53 45	24, 38, 64, 85	0
2	H	346/377 (91%)	0.02	15 (4%) 40 31	30, 50, 74, 96	0
2	J	346/377 (91%)	-0.03	10 (2%) 53 45	28, 48, 74, 94	0
2	L	346/377 (91%)	-0.48	6 (1%) 69 63	21, 34, 54, 77	0
3	M	5/10 (50%)	-0.25	0 100 100	35, 35, 43, 53	0
3	N	5/10 (50%)	-0.33	1 (20%) 3 3	35, 39, 41, 53	0
3	O	5/10 (50%)	-0.11	1 (20%) 3 3	34, 35, 41, 52	0
3	P	5/10 (50%)	0.18	1 (20%) 3 3	40, 41, 47, 59	0
3	Q	5/10 (50%)	0.11	1 (20%) 3 3	38, 39, 42, 54	0
3	R	5/10 (50%)	-0.28	0 100 100	33, 34, 39, 51	0
All	All	3990/4584 (87%)	-0.07	116 (2%) 53 45	21, 45, 73, 100	0

The worst 5 of 116 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	55	PHE	7.7
1	C	55	PHE	6.3
1	E	55	PHE	5.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	H	18	LEU	5.7
1	I	55	PHE	5.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
7	MES	F	380	12/12	0.92	0.17	72,77,79,79	0
5	MGM	H	380	29/29	0.95	0.12	30,38,53,54	0
5	MGM	B	379	29/29	0.95	0.12	33,40,53,54	0
5	MGM	D	380	29/29	0.96	0.10	29,35,47,47	0
5	MGM	L	379	29/29	0.96	0.10	18,29,42,44	0
5	MGM	F	381	29/29	0.96	0.10	27,39,47,47	0
5	MGM	J	379	29/29	0.97	0.10	24,34,43,44	0
6	CL	F	379	1/1	0.99	0.04	35,35,35,35	0
6	CL	H	379	1/1	0.99	0.04	47,47,47,47	0
6	CL	D	379	1/1	0.99	0.05	37,37,37,37	0
4	ZN	J	378	1/1	1.00	0.01	33,33,33,33	0
4	ZN	L	378	1/1	1.00	0.01	29,29,29,29	0
4	ZN	B	378	1/1	1.00	0.02	33,33,33,33	0
4	ZN	D	378	1/1	1.00	0.02	31,31,31,31	0
4	ZN	F	378	1/1	1.00	0.02	32,32,32,32	0
4	ZN	H	378	1/1	1.00	0.01	43,43,43,43	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around MGM H 380:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around MGM B 379:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MGM D 380:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around MGM L 379:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MGM F 381:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around MGM J 379:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.