



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 05:42 PM UTC

PDB ID : 5U6C / pdb_00005u6c
Title : Crystal structure of the Mer kinase domain in complex with a macrocyclic inhibitor
Authors : Gajiwala, K.S.; Ferre, R.A.
Deposited on : 2016-12-07
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	315	7% 77% 9% 14%
1	B	315	9% 71% 10% 18%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4502 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tyrosine-protein kinase Mer.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	272	Total	C	N	O	S	0	0	0
			2189	1399	363	406	21			
1	B	257	Total	C	N	O	S	0	1	0
			2075	1329	346	378	22			

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	550	MET	-	initiating methionine	UNP Q12866
A	551	GLY	-	expression tag	UNP Q12866
A	552	SER	-	expression tag	UNP Q12866
A	553	SER	-	expression tag	UNP Q12866
A	554	HIS	-	expression tag	UNP Q12866
A	555	HIS	-	expression tag	UNP Q12866
A	556	HIS	-	expression tag	UNP Q12866
A	557	HIS	-	expression tag	UNP Q12866
A	558	HIS	-	expression tag	UNP Q12866
A	559	HIS	-	expression tag	UNP Q12866
A	560	SER	-	expression tag	UNP Q12866
A	561	SER	-	expression tag	UNP Q12866
A	562	GLY	-	expression tag	UNP Q12866
A	563	GLU	-	expression tag	UNP Q12866
A	564	ASN	-	expression tag	UNP Q12866
A	565	LEU	-	expression tag	UNP Q12866
A	566	TYR	-	expression tag	UNP Q12866
A	567	PHE	-	expression tag	UNP Q12866
A	568	GLN	-	expression tag	UNP Q12866
A	569	GLY	-	expression tag	UNP Q12866
A	650	MET	ILE	conflict	UNP Q12866
B	550	MET	-	initiating methionine	UNP Q12866
B	551	GLY	-	expression tag	UNP Q12866
B	552	SER	-	expression tag	UNP Q12866
B	553	SER	-	expression tag	UNP Q12866

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	554	HIS	-	expression tag	UNP Q12866
B	555	HIS	-	expression tag	UNP Q12866
B	556	HIS	-	expression tag	UNP Q12866
B	557	HIS	-	expression tag	UNP Q12866
B	558	HIS	-	expression tag	UNP Q12866
B	559	HIS	-	expression tag	UNP Q12866
B	560	SER	-	expression tag	UNP Q12866
B	561	SER	-	expression tag	UNP Q12866
B	562	GLY	-	expression tag	UNP Q12866
B	563	GLU	-	expression tag	UNP Q12866
B	564	ASN	-	expression tag	UNP Q12866
B	565	LEU	-	expression tag	UNP Q12866
B	566	TYR	-	expression tag	UNP Q12866
B	567	PHE	-	expression tag	UNP Q12866
B	568	GLN	-	expression tag	UNP Q12866
B	569	GLY	-	expression tag	UNP Q12866
B	650	MET	ILE	conflict	UNP Q12866

- Molecule 2 is (10R)-7-amino-11-chloro-12-fluoro-1-(2-hydroxyethyl)-3,10,16-trimethyl-16,17-dihydro-1H-8,4-(azeno)pyrazolo[4,3-h][2,5,11]benzoxadiazacyclotetradecin-15(10H)-one (CCD ID: 7YS) (formula: C₂₁H₂₂ClFN₆O₃).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	Cl	F	N	O	0	0
			32	21	1	1	6	3		
2	B	1	Total	C	Cl	F	N	O	0	0
			32	21	1	1	6	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	90	Total 90	O 90	0	0
3	B	84	Total 84	O 84	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosine-protein kinase Mer



- Molecule 1: Tyrosine-protein kinase Mer



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	51.69Å 92.33Å 69.27Å 90.00° 100.80° 90.00°	Depositor
Resolution (Å)	25.34 – 2.10 25.34 – 2.10	Depositor EDS
% Data completeness (in resolution range)	93.7 (25.34-2.10) 93.9 (25.34-2.10)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.00 (at 1.86Å)	Xtriage
Refinement program	BUSTER 2.11.7	Depositor
R, R_{free}	0.211 , 0.257 (Not available) , 0.260	Depositor DCC
R_{free} test set	2585 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	31.9	Xtriage
Anisotropy	0.600	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 55.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4502	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 7YS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.83	0/2233	1.30	8/3014 (0.3%)
1	B	0.80	0/2117	1.25	5/2858 (0.2%)
All	All	0.82	0/4350	1.27	13/5872 (0.2%)

There are no bond length outliers.

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	763	PRO	CA-C-N	7.55	130.34	120.60
1	A	763	PRO	C-N-CA	7.55	130.34	120.60
1	A	775	ARG	CA-C-N	5.87	128.60	120.49
1	A	775	ARG	C-N-CA	5.87	128.60	120.49
1	A	585	ASN	CA-CB-CG	5.76	118.36	112.60
1	A	644	PHE	CA-CB-CG	5.66	119.46	113.80
1	B	775	ARG	CA-C-N	5.63	128.91	120.69
1	B	775	ARG	C-N-CA	5.63	128.91	120.69
1	A	840	ASP	CA-CB-CG	5.41	118.01	112.60
1	A	703	PHE	CA-CB-CG	5.24	119.04	113.80
1	B	644	PHE	CA-CB-CG	5.16	118.96	113.80
1	B	822	PRO	CA-C-N	5.16	127.51	120.54
1	B	822	PRO	C-N-CA	5.16	127.51	120.54

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2189	0	2196	8	0
1	B	2075	0	2090	11	0
2	A	32	0	0	0	0
2	B	32	0	0	0	0
3	A	90	0	0	0	0
3	B	84	0	0	0	0
All	All	4502	0	4286	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:652:LEU:HD13	1:B:669:VAL:HG13	1.84	0.59
1:A:621:MET:HE1	1:A:744:LEU:HG	1.87	0.56
1:A:823:GLU:HA	1:B:689:GLU:HG2	1.90	0.52
1:A:582:ILE:HG21	1:A:606:LEU:HD11	1.92	0.52
1:A:620:THR:HG22	1:A:668:MET:HG2	1.94	0.49
1:B:651[A]:ARG:HH12	1:B:732:ARG:HH12	1.59	0.48
1:B:621:MET:HB2	1:B:667:PRO:HG2	1.96	0.46
1:A:764:VAL:HA	1:A:767:ILE:HD12	1.97	0.46
1:A:685:TYR:HB3	1:A:692:PRO:HD2	1.98	0.45
1:B:620:THR:HG22	1:B:668:MET:HG2	1.98	0.44
1:B:727:ARG:HH22	1:B:763:PRO:HB3	1.81	0.44
1:B:601:VAL:HG22	1:B:619:LYS:HD3	1.99	0.43
1:B:701:LEU:HD21	1:B:829:LEU:HD13	1.99	0.43
1:B:580:VAL:HG23	1:B:642:LYS:HD2	2.02	0.41
1:A:731:LEU:HD23	1:A:737:VAL:HG22	2.01	0.41
1:A:687:ARG:HD2	1:A:796:ARG:HB3	2.03	0.41
1:B:711:MET:HE1	1:B:721:HIS:HB2	2.03	0.40
1:B:651[A]:ARG:NH1	1:B:732:ARG:HH12	2.18	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	266/315 (84%)	261 (98%)	4 (2%)	1 (0%)	30	28
1	B	250/315 (79%)	244 (98%)	6 (2%)	0	100	100
All	All	516/630 (82%)	505 (98%)	10 (2%)	1 (0%)	43	44

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	744	LEU

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	246/282 (87%)	235 (96%)	11 (4%)	24	25
1	B	233/282 (83%)	219 (94%)	14 (6%)	17	15
All	All	479/564 (85%)	454 (95%)	25 (5%)	22	20

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	573	LEU
1	A	577	LEU
1	A	585	ASN
1	A	603	GLU
1	A	608	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	629	ARG
1	A	633	GLU
1	A	652	LEU
1	A	693	LYS
1	A	720	LEU
1	A	774	ASP
1	B	577	LEU
1	B	579	ASP
1	B	592	ILE
1	B	603	GLU
1	B	613	SER
1	B	651[A]	ARG
1	B	651[B]	ARG
1	B	656	CYS
1	B	693	LYS
1	B	717	ARG
1	B	720	LEU
1	B	741	ASP
1	B	774	ASP
1	B	811	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	628	GLN
1	A	694	HIS
1	A	805	GLN
1	B	605	ASN
1	B	805	GLN
1	B	807	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	7YS	B	9001	-	35,35,35	0.89	3 (8%)	38,52,52	0.98	1 (2%)
2	7YS	A	9001	-	35,35,35	1.08	3 (8%)	38,52,52	1.05	3 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	7YS	B	9001	-	-	2/27/27/27	0/3/4/4
2	7YS	A	9001	-	-	2/27/27/27	0/3/4/4

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	9001	7YS	C12-C11	4.35	1.54	1.49
2	A	9001	7YS	C7-C8	3.35	1.48	1.42
2	B	9001	7YS	C7-C8	3.17	1.47	1.42
2	B	9001	7YS	C12-C11	2.91	1.52	1.49
2	B	9001	7YS	O18-C17	-2.15	1.43	1.45
2	A	9001	7YS	O18-C17	-2.10	1.43	1.45

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	9001	7YS	C1-O18-C17	3.09	121.51	117.01
2	B	9001	7YS	C1-O18-C17	2.78	121.07	117.01
2	A	9001	7YS	C12-N13-C14	2.22	125.78	119.97
2	A	9001	7YS	C27-N13-C14	-2.22	117.43	123.14

There are no chirality outliers.


All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	9001	7YS	C11-C12-N13-C14
2	B	9001	7YS	C11-C12-N13-C14
2	B	9001	7YS	C11-C12-N13-C27
2	A	9001	7YS	C11-C12-N13-C27

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	272/315 (86%)	0.65	21 (7%)	19	21	20, 37, 77, 110	0
1	B	257/315 (81%)	0.67	28 (10%)	10	11	19, 37, 78, 104	1 (0%)
All	All	529/630 (83%)	0.66	49 (9%)	14	15	19, 37, 77, 110	1 (0%)

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	631	ILE	6.2
1	A	573	LEU	5.7
1	B	777	TYR	4.4
1	A	777	TYR	4.2
1	A	665	PRO	4.2
1	A	577	LEU	4.2
1	B	636	SER	4.1
1	A	598	PHE	4.1
1	A	864	VAL	4.1
1	A	745	SER	3.8
1	A	664	ILE	3.8
1	B	635	LEU	3.3
1	A	597	GLU	3.3
1	B	742	PHE	3.2
1	B	595	GLU	3.2
1	A	659	MET	3.2
1	B	579	ASP	3.1
1	B	577	LEU	3.1
1	B	773	ALA	2.9
1	A	576	LYS	2.9
1	B	690	THR	2.9
1	A	776	VAL	2.8
1	B	634	PHE	2.8
1	B	862	PRO	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	576	LYS	2.7
1	A	656	CYS	2.6
1	A	690	THR	2.6
1	A	772	LEU	2.6
1	B	814	LEU	2.6
1	B	651[A]	ARG	2.4
1	A	743	GLY	2.4
1	B	580	VAL	2.4
1	B	621	MET	2.4
1	A	693	LYS	2.4
1	B	861	LEU	2.3
1	B	776	VAL	2.3
1	B	772	LEU	2.3
1	B	762	MET	2.3
1	B	640	CYS	2.3
1	B	689	GLU	2.1
1	A	575	ASN	2.1
1	A	689	GLU	2.1
1	A	763	PRO	2.1
1	B	695	ILE	2.1
1	B	664	ILE	2.1
1	B	575	ASN	2.0
1	A	691	GLY	2.0
1	B	691	GLY	2.0
1	B	643	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	7YS	A	9001	32/32	0.92	0.08	21,28,30,39	0
2	7YS	B	9001	32/32	0.93	0.08	23,29,32,41	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 7YS A 9001:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.