wwPDB EM Validation Summary Report

Oct 19, 2026 – 05:00 PM JST

PDB ID	:	9UEU / pdb_00009ueu
EMDB ID	:	EMD-64092
Title	:	A helper NLR structure of Pii-1
Authors	:	Xiong, Y.; Zhang, X.; Xiao, Y.
Deposited on	:	2025-04-09
Resolution	:	3.31 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


EMDB validation analysis	:	0.0.1.dev133
Mogul	:	1.8.5 (274361), CSD as541be (2020)
MolProbity	:	4-5-2 with Phenix2.2
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics	:	202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ	:	1.9.13
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51


1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	14550 (2.81 - 3.81)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1025	5% 19% 13% · 66%
1	B	1025	6% 19% 13% · 66%
1	C	1025	5% 19% 13% · 66%
1	D	1025	5% 19% 13% · 66%
1	E	1025	5% 19% 13% · 66%

2 Entry composition


There are 2 unique types of molecules in this entry. The entry contains 14160 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pii-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	E	348	Total	C	N	O	S	0	0
			2801	1789	475	520	17		
1	A	348	Total	C	N	O	S	0	0
			2801	1789	475	520	17		
1	B	348	Total	C	N	O	S	0	0
			2801	1789	475	520	17		
1	C	348	Total	C	N	O	S	0	0
			2801	1789	475	520	17		
1	D	348	Total	C	N	O	S	0	0
			2801	1789	475	520	17		

- Molecule 2 is ADENOSINE-5'-TRIPHOSPHATE (CCD ID: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					AltConf
2	E	1	Total	C	N	O	P	0
			31	10	5	13	3	

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf
2	A	1	Total	C	N	O	P	0
			31	10	5	13	3	
2	B	1	Total	C	N	O	P	0
			31	10	5	13	3	
2	C	1	Total	C	N	O	P	0
			31	10	5	13	3	
2	D	1	Total	C	N	O	P	0
			31	10	5	13	3	


3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pii-1

Chain E:









• Molecule 1: Pii-1









• Molecule 1: Pii-1

Chain B:









• Molecule 1: Pii-1









• Molecule 1: Pii-1

Chain D: 5% 19% 13% 66%




LEU SER LYS CYS LEU LYS LEU GLU VAL LEU PRO GLN SER PHE GLY GLN LEU ALA TYR LEU LYS SER LEU ASN LEU SER TYR CYS SER ASP LEU LYS LEU LEU GLU SER PHE GLU CYS LEU THR SER LEU ARG PHE LEU ASN LEU SER ASN CYS SER ARG LEU GLU TYR LEU PRO SER CYS
PHE ASP LYS LEU ASN ASN LEU GLU SER LEU ASN LEU SER GLN CYS LEU GLY LEU LYS ALA LEU PRO GLU SER LEU GLN ASN LEU LYS ASN LEU GLN LEU ASP VAL SER GLY CYS GLN ASP CYS ILE VAL GLN SER PHE SER LEU SER THR ARG SER SER GLN SER CYS GLN ARG SER GLU
LYS ALA GLU GLN VAL ARG SER ARG ASN SER GLU ILE SER GLU ILE THR TYR GLU GLU PRO ALA GLU ILE GLU LEU LEU LYS ASN ASN PRO SER LYS ASP LEU ALA SER ILE SER HIS LEU ASN GLU ASP ARG ILE GLU GLU PRO GLU VAL VAL THR GLU PRO SER ALA THR ARG GLY MET
VAL GLN GLN ILE PRO GLY ASN GLN LEU SER SER PRO SER SER HIS LEU SER SER PHE ALA SER SER SER ALA PRO PHE ALA SER SER SER SER ASP THR SER THR SER GLU HIS PRO VAL PRO ASN GLU GLU ALA ALA ALA LEU THR VAL PRO ARG SER ASN GLU LYS CYS ASP ASN THR
PRO MET PRO VAL LYS ASP GLY LEU ILE SER GLU ASP ASP ALA PRO VAL HIS LEU HIS GLN LYS PRO LEU GLN ALA THR ALA MET ALA ALA ILE

4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	126423	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI POLARA 300	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/Å^2$)	50	Depositor
Minimum defocus (nm)	1300	Depositor
Maximum defocus (nm)	1800	Depositor
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.095	Depositor
Minimum map value	-0.045	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.003	Depositor
Recommended contour level	0.0219	Depositor
Map size (Å)	218.0, 218.0, 218.0	wwPDB
Map dimensions	200, 200, 200	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.09, 1.09, 1.09	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| > 5$	RMSZ	$\#\|Z\| > 5$
1	A	0.39	0/2853	0.53	1/3853 (0.0%)
1	B	0.39	0/2853	0.53	1/3853 (0.0%)
1	C	0.39	0/2853	0.53	1/3853 (0.0%)
1	D	0.39	0/2853	0.53	1/3853 (0.0%)
1	E	0.39	0/2853	0.53	1/3853 (0.0%)
All	All	0.39	0/14265	0.53	5/19265 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
1	C	0	2
1	D	0	2
1	E	0	2
All	All	0	10

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	C	171	ILE	CA-CB-CG1	5.11	119.09	110.40
1	D	171	ILE	CA-CB-CG1	5.10	119.06	110.40
1	B	171	ILE	CA-CB-CG1	5.09	119.06	110.40
1	E	171	ILE	CA-CB-CG1	5.08	119.04	110.40
1	A	171	ILE	CA-CB-CG1	5.08	119.04	110.40

There are no chirality outliers.

5 of 10 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	280	SER	Peptide
1	A	401	TYR	Peptide
1	B	280	SER	Peptide
1	E	280	SER	Peptide
1	E	401	TYR	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2801	0	2839	110	0
1	B	2801	0	2839	115	0
1	C	2801	0	2837	114	0
1	D	2801	0	2839	116	0
1	E	2801	0	2837	117	0
2	A	31	0	12	4	0
2	B	31	0	12	6	0
2	C	31	0	11	5	0
2	D	31	0	12	8	0
2	E	31	0	12	6	0
All	All	14160	0	14250	551	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 551 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:292:LYS:O	1:B:296:LEU:HB2	1.75	0.87
1:E:292:LYS:O	1:E:296:LEU:HB2	1.75	0.86
1:C:292:LYS:O	1:C:296:LEU:HB2	1.75	0.85
1:C:219:THR:O	1:C:225:ARG:NH1	2.09	0.85
1:C:172:VAL:H	2:C:1101:ATP:HN62	1.24	0.85

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	344/1025 (34%)	292 (85%)	52 (15%)	0	100	100
1	B	344/1025 (34%)	292 (85%)	52 (15%)	0	100	100
1	C	344/1025 (34%)	293 (85%)	51 (15%)	0	100	100
1	D	344/1025 (34%)	294 (86%)	50 (14%)	0	100	100
1	E	344/1025 (34%)	293 (85%)	51 (15%)	0	100	100
All	All	1720/5125 (34%)	1464 (85%)	256 (15%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	312/926 (34%)	273 (88%)	39 (12%)	4	19
1	B	312/926 (34%)	273 (88%)	39 (12%)	4	19
1	C	312/926 (34%)	273 (88%)	39 (12%)	4	19
1	D	312/926 (34%)	273 (88%)	39 (12%)	4	19
1	E	312/926 (34%)	273 (88%)	39 (12%)	4	19
All	All	1560/4630 (34%)	1365 (88%)	195 (12%)	7	19

5 of 195 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	467	THR
1	C	385	ILE
1	C	166	ASN
1	C	261	ILE
1	C	461	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 35 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	187	GLN
1	D	236	ASN
1	D	387	GLN
1	B	187	GLN
1	A	504	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\|>2$	Counts	RMSZ	$\#\|Z\|>2$
2	ATP	E	1101	-	29,33,33	1.31	6 (20%)	44,52,52	2.26	13 (29%)
2	ATP	C	1101	-	29,33,33	1.30	4 (13%)	44,52,52	2.08	10 (22%)
2	ATP	D	1101	-	29,33,33	1.33	6 (20%)	44,52,52	1.87	11 (25%)
2	ATP	A	1101	-	29,33,33	1.33	4 (13%)	44,52,52	2.08	9 (20%)
2	ATP	B	1101	-	29,33,33	1.32	6 (20%)	44,52,52	2.02	12 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ATP	E	1101	-	-	3/22/38/38	0/3/3/3
2	ATP	C	1101	-	-	5/22/38/38	0/3/3/3
2	ATP	D	1101	-	-	4/22/38/38	0/3/3/3
2	ATP	A	1101	-	-	4/22/38/38	0/3/3/3
2	ATP	B	1101	-	-	5/22/38/38	0/3/3/3

The worst 5 of 26 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1101	ATP	C5-C4	3.86	1.46	1.39
2	C	1101	ATP	C5-C4	3.83	1.46	1.39
2	B	1101	ATP	C5-C4	3.72	1.46	1.39
2	E	1101	ATP	C5-C4	3.69	1.45	1.39
2	D	1101	ATP	C5-C4	3.69	1.45	1.39

The worst 5 of 55 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	E	1101	ATP	PB-O3B-PG	-6.33	111.11	132.83
2	C	1101	ATP	C5-C4-N3	-6.22	118.64	126.75
2	A	1101	ATP	PA-O3A-PB	-6.18	111.63	132.83
2	A	1101	ATP	C5-C4-N3	-6.10	118.79	126.75
2	C	1101	ATP	PA-O3A-PB	-5.76	113.07	132.83

There are no chirality outliers.

5 of 21 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	E	1101	ATP	C5'-O5'-PA-O1A
2	A	1101	ATP	C5'-O5'-PA-O2A
2	A	1101	ATP	C5'-O5'-PA-O3A
2	B	1101	ATP	PB-O3B-PG-O3G
2	C	1101	ATP	C5'-O5'-PA-O3A


There are no ring outliers.

5 monomers are involved in 29 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	1101	ATP	6	0
2	C	1101	ATP	5	0
2	D	1101	ATP	8	0
2	A	1101	ATP	4	0
2	B	1101	ATP	6	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand ATP E 1101




Bond lengths



Bond angles




Torsions




Rings


Ligand ATP C 1101




Bond lengths



Bond angles




Torsions




Rings


Ligand ATP D 1101




Bond lengths



Bond angles




Torsions




Rings


Ligand ATP A 1101




Bond lengths



Bond angles




Torsions



Rings

Ligand ATP B 1101



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.


6 Map visualisation

This section contains visualisations of the EMDB entry EMD-64092. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)


6.2.1 Primary map



X Index: 100




Y Index: 100




Z Index: 100


6.2.2 Raw map



X Index: 100



Y Index: 100




Z Index: 100


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)


6.3.1 Primary map



X Index: 80




Y Index: 73




Z Index: 100


6.3.2 Raw map



X Index: 80



Y Index: 73




Z Index: 100


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) (i)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views (i)


6.5.1 Primary map



X Y Z

The images above show the 3D surface view of the map at the recommended contour level 0.0219. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



X Y Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation (i)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_64092_msk_1.map (i)



X



Y




Z

7 Map analysis


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate (i)



The volume at the recommended contour level is 64 nm^3; this corresponds to an approximate mass of 58 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum (i)




*Reported resolution corresponds to spatial frequency of 0.302 Å^{-1}

8 Fourier-Shell correlation

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC



*Reported resolution corresponds to spatial frequency of 0.302 Å^{-1}

8.2 Resolution estimates (i)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.31	-	-
Author-provided FSC curve	3.30	3.77	3.35
Unmasked-calculated*	3.64	4.17	3.71

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.


9 Map-model fit

This section contains information regarding the fit between EMDB map EMD-64092 and PDB model 9UEU. Per-residue inclusion information can be found in section 3 on page 5.


9.1 Map-model overlay



X




Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.0219 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.


9.2 Q-score mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0219).

9.4 Atom inclusion (i)



At the recommended contour level, 79% of all backbone atoms, 62% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.0219) and Q-score for the entire model and for each chain.



Chain	Atom inclusion	Q-score
All	0.6250	0.4800
A	0.6280	0.4830
B	0.6170	0.4730
C	0.6280	0.4810
D	0.6240	0.4800
E	0.6280	0.4820